



COMPOSITIO MATHEMATICA

Invariants of reflection groups, arrangements, and normality of decomposition classes in Lie algebras

J. Matthew Douglass and Gerhard Röhrle

Compositio Math. **148** (2012), 921–930.

[doi:10.1112/S0010437X11007512](https://doi.org/10.1112/S0010437X11007512)



FOUNDATION
COMPOSITIO
MATHEMATICA



LONDON
MATHEMATICAL
SOCIETY



Invariants of reflection groups, arrangements, and normality of decomposition classes in Lie algebras

J. Matthew Douglass and Gerhard Röhrle

ABSTRACT

Suppose that W is a finite, unitary, reflection group acting on the complex vector space V and X is a subspace of V . Define N to be the setwise stabilizer of X in W , Z to be the pointwise stabilizer, and $C = N/Z$. Then restriction defines a homomorphism from the algebra of W -invariant polynomial functions on V to the algebra of C -invariant functions on X . In this note we consider the special case when W is a Coxeter group, V is the complexified reflection representation of W , and X is in the lattice of the arrangement of W , and give a simple, combinatorial characterization of when the restriction mapping is surjective in terms of the exponents of W and C . As an application of our result, in the case when W is the Weyl group of a semisimple, complex Lie algebra, we complete a calculation begun by Richardson in 1987 and obtain a simple combinatorial characterization of regular decomposition classes whose closure is a normal variety.

1. Introduction

Suppose that W is a finite, complex reflection group acting on the complex vector space $V = \mathbb{C}^l$ and X is a subspace of V . Define $N_X = \{w \in W \mid w(X) = X\}$, the setwise stabilizer of X in W , and $Z_X = \{w \in W \mid w(x) = x \forall x \in X\}$, the pointwise stabilizer of X in V . Then Z_X is a normal subgroup of N_X and we set $C_X = N_X/Z_X$. It is easy to see that restriction defines a homomorphism from the algebra of W -invariant polynomial functions on V to the algebra of C_X -invariant functions on X , say $\rho : \mathbb{C}[V]^W \rightarrow \mathbb{C}[X]^{C_X}$. In this note we consider the special case when W is a Coxeter group, V is the complexified reflection representation of W , and X is in the lattice of the arrangement of W . Our main result is a simple combinatorial characterization in terms of the exponents of W and C_X of when the map ρ is surjective.

As an application, our main result combined with a theorem of Richardson [Ric87] leads immediately to a complete, and easily computable, classification of the regular decomposition classes in a semisimple, complex Lie algebra whose closure is a normal variety.

2. Statement of the main results

By a *hyperplane arrangement* we mean a pair (V, \mathcal{A}) , where \mathcal{A} is a finite set of hyperplanes in V . The arrangement of a subgroup $C \subseteq \mathrm{GL}(V)$ consists of the reflecting hyperplanes of the elements

Received 13 March 2011, accepted in final form 19 October 2011, published online 19 March 2012.

2010 Mathematics Subject Classification 20F55, 14N20 (primary), 13A50 (secondary).

Keywords: arrangements, Coxeter groups, decomposition classes, invariants.

This journal is © [Foundation Compositio Mathematica](http://www.cambridge.org/core) 2012.

in C that act on V as reflections. We denote the arrangement of C in V by $\mathcal{A}(V, C)$. Define C^{ref} to be the subgroup generated by the reflections in C . Then obviously $\mathcal{A}(V, C) = \mathcal{A}(V, C^{\text{ref}})$.

For general information about arrangements and reflection groups, we refer the reader to [Bou68] and [OT92].

Suppose from now on that W is a finite subgroup of $\text{GL}(V)$ generated by reflections. Unless otherwise specified, we allow the case when the generators of W are ‘pseudo-reflections’, that is, elements in $\text{GL}(V)$ with finite order whose 1-eigenspace is a hyperplane in V . For a subspace X of V , we have two natural hyperplane arrangements in X :

- the restricted arrangement $\mathcal{A}(V, W)^X$ consisting of intersections $H \cap X$ for H in $\mathcal{A}(V, W)$ with $X \not\subseteq H$; and
- the reflection arrangement $\mathcal{A}(X, C_X) = \mathcal{A}(X, C_X^{\text{ref}})$ consisting of the reflecting hyperplanes of elements in C_X that act on X as reflections.

For a free hyperplane arrangement \mathcal{A} , we denote the multiset of exponents of \mathcal{A} by $\text{exp}(\mathcal{A})$. Terao [Ter80] has shown that reflection arrangements are free and that $\text{exp}(\mathcal{A}(V, W)) = \text{coexp}(W)$, where $\text{coexp}(W)$ denotes the multiset of coexponents of W .

The lattice of a hyperplane arrangement is the set of subspaces of V of the form $H_1 \cap \dots \cap H_n$, where $\{H_1, \dots, H_n\}$ is a subset of \mathcal{A} . It is known that $\mathcal{A}(V, W)^X$ is free when W is a Coxeter group and X is a subspace in the lattice of $\mathcal{A}(V, W)$ (see [OT93, Dou99]). Thus, in this case, we have that:

- (1) $\text{exp}(\mathcal{A}(X, C_X))$, $\text{exp}(\mathcal{A}(V, W)^X)$, and $\text{exp}(\mathcal{A}(V, W))$ are all defined;
- (2) $\text{exp}(\mathcal{A}(X, C_X)) = \text{exp}(C_X^{\text{ref}})$; and
- (3) $\text{exp}(\mathcal{A}(V, W)) = \text{exp}(W)$.

We can now state our main result.

THEOREM 2.1. *Suppose W is a finite Coxeter group, V affords the reflection representation of W , and X is in the lattice of the arrangement $\mathcal{A}(V, W)$. Then the restriction mapping $\rho : \mathbb{C}[V]^W \rightarrow \mathbb{C}[X]^{C_X}$ is surjective if and only if*

$$\text{exp}(\mathcal{A}(X, C_X)) = \text{exp}(\mathcal{A}(V, W)^X) \subseteq \text{exp}(\mathcal{A}(V, W)).$$

To simplify the notation, in the rest of this paper we denote the arrangements $\mathcal{A}(X, C_X)$, $\mathcal{A}(V, W)^X$, and $\mathcal{A}(V, W)$ by $\mathcal{A}(C_X)$, \mathcal{A}^X , and \mathcal{A} , respectively.

In the next section, using a modification of an argument of Denef and Loeser [DL95], we show in Proposition 3.1 that if W is any complex reflection group, X is in the lattice of \mathcal{A} , $C_X = C_X^{\text{ref}}$, and ρ is surjective, then $\mathcal{A}(C_X) = \mathcal{A}^X$ and $\text{exp}(C_X) \subseteq \text{exp}(W)$. It then follows that in this case \mathcal{A}^X is a free arrangement and $\text{exp}(\mathcal{A}(C_X)) = \text{exp}(\mathcal{A}^X)$ and $\text{exp}(C_X) \subseteq \text{exp}(W)$. In particular, the forward implication in the theorem holds whenever C_X acts on X as a reflection group.

In § 4, we complete the proof of Theorem 2.1 by:

- (1) showing in Proposition 4.1 that if W is a Coxeter group and C_X does not act on X as a reflection group, then ρ is not surjective; and
- (2) computing all cases in which $\text{exp}(\mathcal{A}(C_X)) = \text{exp}(\mathcal{A}^X) \subseteq \text{exp}(\mathcal{A})$ for a Coxeter group W and showing that ρ is surjective in these cases.

Notice that the conditions $\text{exp}(\mathcal{A}(C_X)) = \text{exp}(\mathcal{A}^X) \subseteq \text{exp}(\mathcal{A})$ are not that easy to satisfy. In case W is a Coxeter group of type A_{r-1} , up to the action of W , the subspaces X in the lattice

of \mathcal{A} are parametrized by partitions of r . The conditions $\exp(\mathcal{A}(C_X)) = \exp(\mathcal{A}^X) \subseteq \exp(\mathcal{A})$ hold if and only if the corresponding partition of r has equal parts. For W a Coxeter group of type E_8 , up to the action of W , there are 41 possibilities for X , eight of which have the property that $\exp(\mathcal{A}(C_X)) = \exp(\mathcal{A}^X) \subseteq \exp(\mathcal{A})$. All cases in which $\exp(\mathcal{A}(C_X)) = \exp(\mathcal{A}^X) \subseteq \exp(\mathcal{A})$ when W is a finite, irreducible Coxeter group are listed in Tables 1 and 2 in § 4.

In the rest of this section, we explain how our main result leads to a characterization of regular decomposition classes in a complex, semisimple Lie algebra whose closure is a normal variety. The classification of these decomposition classes was completed, case-by-case, for classical Weyl groups by Richardson in 1987 [Ric87] and extended by Broer in 1998 [Bro98], again using case-by-case arguments, to exceptional Weyl groups.

Suppose that \mathfrak{g} is a semisimple, complex Lie algebra and G is the adjoint group of \mathfrak{g} . Motivated by a question of De Concini and Procesi about the normality of the closure of the G -saturation of a Cartan subspace for an involution of \mathfrak{g} , Richardson proved the following theorem.

THEOREM 2.2 [Ric87, Theorem B]. *Suppose that \mathfrak{t} is a Cartan subalgebra of \mathfrak{g} , W is the Weyl group of $(\mathfrak{g}, \mathfrak{t})$, and X is a subspace of \mathfrak{t} with the property that C_X acts on X as a reflection group. Let Y denote the closure of the set of elements in \mathfrak{g} whose semisimple part is in $\text{Ad}(G)X$. Then Y is a normal, Cohen–Macaulay variety if and only if $\rho : \mathbb{C}[\mathfrak{t}]^W \rightarrow \mathbb{C}[X]^{C_X}$ is surjective.*

When $V = \mathfrak{t}$ is a Cartan subalgebra of \mathfrak{g} , a subspace X of \mathfrak{t} is in the lattice of $\mathcal{A}(\mathfrak{t}, W)$ if and only if there are a parabolic subalgebra \mathfrak{p} of \mathfrak{g} and a Levi subalgebra \mathfrak{l} of \mathfrak{p} with $\mathfrak{t} \subseteq \mathfrak{l}$ so that $X = \mathfrak{z}$ is the centre of \mathfrak{l} .

Now let $\mathfrak{g}_{\text{reg}}$ denote the set of regular elements in \mathfrak{g} . Then $\mathfrak{g}_{\text{reg}}$ is the disjoint union of decomposition classes of \mathfrak{g} (see [Bor81, § 3]). A decomposition class contained in $\mathfrak{g}_{\text{reg}}$ is a *regular decomposition class*. Suppose that \mathfrak{l} and \mathfrak{z} are as in the last paragraph, \mathfrak{z}_0 is the subspace of elements in \mathfrak{z} whose centralizer in \mathfrak{g} is \mathfrak{l} , and \mathcal{O} is the regular, nilpotent, adjoint orbit in \mathfrak{l} . Then $\text{Ad}(G)(\mathfrak{z}_0 + \mathcal{O})$ is a regular decomposition class. Moreover, every regular decomposition class is of this form for some \mathfrak{l} (see [Bor81, § 3]). Therefore, combining Theorems 2.1 and 2.2, we obtain the following characterization of regular decomposition classes in \mathfrak{g} that have normal closure.

THEOREM 2.3. *With the notation above, suppose that $D = \text{Ad}(G)(\mathfrak{z}_0 + \mathcal{O})$ is a regular decomposition class in \mathfrak{g} . Then \overline{D} is a normal variety if and only if*

$$\exp(\mathcal{A}(\mathfrak{z}, C_{\mathfrak{z}})) = \exp(\mathcal{A}(\mathfrak{t}, W)^{\mathfrak{z}}) \subseteq \exp(\mathcal{A}(\mathfrak{t}, W)).$$

Using case-by-case arguments, Richardson [Ric87] determined all cases in which $\rho : \mathbb{C}[\mathfrak{t}]^W \rightarrow \mathbb{C}[\mathfrak{z}]^{C_{\mathfrak{z}}}$ is surjective when W is a Weyl group of classical type. Broer [Bro98] computed almost all of the additional cases for exceptional Weyl groups. The statement of [Bro98, Theorem 3.1 (e7)] is missing one case: if \mathfrak{g} is of type E_7 and \mathfrak{l} is of type $(A_1^3)'$ (with simple roots $\alpha_2, \alpha_5, \alpha_7$, where the labelling is as in [Bou68]), then the restriction map ρ is surjective.

3. A preliminary result

In this section, we prove the following result.

PROPOSITION 3.1. *Suppose $W \subseteq \text{GL}(V)$ is a complex reflection group, X is in the lattice of \mathcal{A} , C_X acts on X as a reflection group, and the restriction mapping $\rho : \mathbb{C}[V]^W \rightarrow \mathbb{C}[X]^{C_X}$ is surjective. Then $\exp(C_X) \subseteq \exp(W)$ and $\mathcal{A}(C_X) = \mathcal{A}^X$. Thus, \mathcal{A}^X is a free arrangement and if W is a Coxeter group, then $\exp(\mathcal{A}(C_X)) = \exp(\mathcal{A}^X) \subseteq \exp(\mathcal{A})$.*

The proof shows that if X is any subspace of V , C_X acts on X as a reflection group, and ρ is surjective, then $\exp(C_X) \subseteq \exp(W)$ and $\mathcal{A}(C_X) \subseteq \mathcal{A}^X$. The assumption that X is in the lattice of \mathcal{A} is only used to conclude that $\mathcal{A}^X \subseteq \mathcal{A}(C_X)$.

By assumption, the restriction mapping $\rho : \mathbb{C}[V]^W \rightarrow \mathbb{C}[X]^{C_X}$ is a degree-preserving, surjective homomorphism of graded polynomial algebras and so, by a result of Richardson [Ric87, §4], we may choose algebraically independent, homogeneous polynomials f_1, \dots, f_r in $\mathbb{C}[V]^W$ so that $\mathbb{C}[V]^W = \mathbb{C}[f_1, \dots, f_r]$ and $\mathbb{C}[X]^{C_X} = \mathbb{C}[\rho(f_1), \dots, \rho(f_r)]$. Since $\exp(C_X) = \{\deg f_1 - 1, \dots, \deg f_l - 1\}$ and $\exp(W) = \{\deg f_1 - 1, \dots, \deg f_r - 1\}$, we have $\exp(C_X) \subseteq \exp(W)$.

We next show that $\mathcal{A}(C_X) \subseteq \mathcal{A}^X$. Suppose K is in $\mathcal{A}(C_X)$. By assumption, there is a w in N_X so that $\text{Fix}(w) \cap X = K$. It was shown in [OT92, Theorem 6.27] that $\text{Fix}(w)$ is in the lattice of \mathcal{A} , say $\text{Fix}(w) = H_1 \cap \dots \cap H_n$, where H_1, \dots, H_n are in \mathcal{A} . Then $K = H_1 \cap \dots \cap H_n \cap X$. Since $\dim K = \dim X - 1$, it follows that $K = H_i \cap X$ for some i and so K is in \mathcal{A}^X .

It remains to show that $\mathcal{A}^X \subseteq \mathcal{A}(C_X)$. We use a variant of an argument given by Denef and Loeser [DL95] (see also [LS99]).

Suppose that homogeneous polynomial invariants $\{f_1, \dots, f_r\}$ have been chosen as above. Let J denote the $r \times r$ matrix whose (i, j) th entry is $(\partial f_i / \partial x_j)$ and let J_1 denote the $l \times l$ submatrix of J consisting of the first l rows and columns. Then J and J_1 are matrices of functions on V . For v in V , let $J(v)$ and $J_1(v)$ be the matrices obtained from J and J_1 , respectively, by evaluating each entry at v .

Then $\det J_1$ is in $\mathbb{C}[V]$ and, by a result of Steinberg (see [OT92, §6.2]), the zero set of $\rho(\det J_1) = \det \rho(J_1)$ in X is precisely $\bigcup_{K \in \mathcal{A}(C_X)} K$. Thus, to show that $\mathcal{A}^X \subseteq \mathcal{A}(C_X)$, it is enough to show that if K is in \mathcal{A}^X , then $\rho(\det J_1)$ vanishes on K .

Denef and Loeser have shown that if w is in W , v_1 and v_2 are eigenvectors for w with eigenvalues λ_1 and λ_2 , respectively, and f in $\mathbb{C}[V]^W$ is homogeneous with degree d , then $\lambda_2 D_{v_2}(f)(v_1) = \lambda_1^{1-d} D_{v_2}(f)(v_1)$, where $D_v(f)$ denotes the directional derivative of f in the direction of v . This proves the following lemma.

LEMMA 3.2. *Suppose w is in W , x is in $\text{Fix}(w)$, and v in V is an eigenvector of w with eigenvalue $\lambda \neq 1$. Then $D_v(f)(x) = 0$ for every f in $\mathbb{C}[V]^W$.*

Suppose H is in \mathcal{A} , s is a reflection in W that fixes H , and v is orthogonal to H with respect to some W -invariant inner product on V . Since H is the full 1-eigenspace of s in V , Lemma 3.2 shows that

$$D_v(f) \text{ vanishes on } H \text{ for every } f \text{ in } \mathbb{C}[V]^W. \tag{3.3}$$

By [OT92, Theorem 6.27], we may find w in W with $\text{Fix}(w) = X$. Choose a basis $\{b_1, \dots, b_r\}$ of V consisting of eigenvectors for w so that $\{b_1, \dots, b_l\}$ is a basis of X . Let $\{x_1, \dots, x_r\}$ denote the dual basis of V^* . Since X is the full 1-eigenspace of w in V , Lemma 3.2 shows that

$$\text{for } j > l, \quad D_{b_j}(f) = \frac{\partial f}{\partial x_j} \text{ vanishes on } X \text{ for every } f \text{ in } \mathbb{C}[V]^W. \tag{3.4}$$

Now suppose K is in \mathcal{A}^X . Say $K = H \cap X$, where H is in \mathcal{A} with $X \not\subseteq H$. Choose v in V orthogonal to H with respect to a W -invariant inner product. Say $v = \sum_{i=1}^r \xi_i b_i$. Define $[v]$ to be the column vector whose i th entry is ξ_i for $1 \leq i \leq r$ and $[v_1]$ to be the column vector whose i th entry is ξ_i for $1 \leq i \leq l$. It follows from (3.3) that $J(h) \cdot [v] = 0$ for every h in H . Therefore, it follows from (3.4) that $J_1(k) \cdot [v_1] = 0$ for every k in K . Since $X \not\subseteq H$, we have $[v_1] \neq 0$ and

so it must be the case that for k in K , the matrix $J_1(k)$ is not invertible. Therefore, $\det J_1$ vanishes on K and so $\rho(\det J_1)$ vanishes on K . Thus, K is in $\mathcal{A}(C_X)$. This completes the proof of Proposition 3.1.

4. Completion of the proof of Theorem 2.1

In this section, we complete the proof of Theorem 2.1 and show that if W is a Coxeter group, V affords the reflection representation of W , and X is in the lattice of \mathcal{A} , then $\rho : \mathbb{C}[V]^W \rightarrow \mathbb{C}[X]^{C_X}$ is surjective if and only if $\exp(\mathcal{A}(C_X)) = \exp(\mathcal{A}^X) \subseteq \exp(\mathcal{A})$.

In the arguments below, ‘degree’ means with respect to the natural grading on $\mathbb{C}[V]$. For an integer d , let $\mathbb{C}[V]_d$ denote the subspace of elements of degree d . For a subalgebra R of $\mathbb{C}[V]$, we set $R_d = R \cap \mathbb{C}[V]_d$. After choosing an appropriate basis of V , we may consider $\mathbb{C}[X]$, $\mathbb{C}[X]^{C_X}$, and $\mathbb{C}[X]^{C_X^{\text{ref}}}$ as subalgebras of $\mathbb{C}[V]$.

Also, we use the conventions that in type A , A_{-1} and A_0 are to be interpreted as the trivial group; in type B , B_0 is to be interpreted as the trivial group and B_1 is to be interpreted as a component of type A_1 supported on a short root; and in type D , D_1 is to be interpreted as the trivial group and D_2 is to be interpreted as a component of type $A_1 \times A_1$ supported on the two distinguished end nodes in the Coxeter graph.

It is easy to see that if $W = W_1 \times W_2$ is reducible, then Theorem 2.1 holds for W if and only if it holds for W_1 and W_2 . Thus, we may assume that W is an irreducible Coxeter group.

Fix a generating set S in W so that (W, S) is a Coxeter system. For a subset I of S , define $X_I = \bigcap_{s \in I} \text{Fix}(s)$ and $W_I = \langle I \rangle$, the subgroup of W generated by I . Orlik and Solomon [OS83] have shown that there are a w in W and a subset I of S so that $w(X) = X_I$, $wZ_X w^{-1} = W_I$, and $wN_X w^{-1} = N_W(W_I)$. Howlett [How80] has shown that W_I has a canonical complement, C_I , in $N_W(W_I)$.

We say that C_I acts on X_I as a Coxeter group with full rank if $C_I = C_I^{\text{ref}}$ and the Coxeter rank of C_I equals the dimension of X_I . For example, if W is of type E_6 and W_I is of type $A_1 \times A_2$, then $C_I = C_I^{\text{ref}}$ is of type A_2 and $\dim X_I = 3$, so C_I does not act on X_I as a Coxeter group with full rank. Another example is when W is of type $I_2(r)$ with r odd and I is a one-element subset of S . In this case, C_I is the trivial group and X_I is one dimensional.

Suppose now that the restriction mapping ρ is surjective. It follows from the next proposition that C_X acts on X as a Coxeter group with full rank. In particular, we may apply Proposition 3.1 and conclude that $\exp(\mathcal{A}(C_X)) = \exp(\mathcal{A}^X) \subseteq \exp(\mathcal{A})$. This proves the forward implication of Theorem 2.1.

PROPOSITION 4.1. *Suppose W is a Coxeter group, X is in the lattice of \mathcal{A} , and C_X does not act on X as a Coxeter group with full rank. Then the restriction mapping $\rho : \mathbb{C}[V]^W \rightarrow \mathbb{C}[X]^{C_X}$ is not surjective.*

Proof. We may assume that W is irreducible and that $X = X_I$ for some subset I of S . Then $W_X = W_I$, $N_X = N_W(W_I)$, and $C_X = C_I$. To show that ρ is not surjective, in each case when C_I does not act on X_I as a Coxeter group with full rank, we find an integer d so that $\dim \mathbb{C}[V]_d^W < \dim \mathbb{C}[X_I]_d^{C_I}$. It then follows that $\mathbb{C}[X_I]_d^{C_I}$ is not contained in the image of ρ .

If $I = \emptyset$ or $I = S$, then C_I acts on X_I as a Coxeter group with full rank. Thus, we may assume that I is a non-empty, proper subset of S .

Howlett [How80] has computed C_I, C_I^{ref} , and the representation of C_I on X_I for all Coxeter groups with rank greater than two. When W has rank two, W is of type $I_2(r)$ for some r . It is easy to see that in this case C_I acts on X_I as a Coxeter group with full rank unless r is odd and $|I| = 1$. Then, as noted above, C_I is the trivial group acting on the one-dimensional vector space X_I .

The subgroup C_I^{ref} is always a normal subgroup of C_I and it turns out that if $C_I^{\text{ref}} \neq C_I$, then C_I is the semidirect product of C_I^{ref} with an elementary abelian 2-group. Notice that if w is any element in C_I with order two, then w acts on X_I with eigenvalues ± 1 , and so w fixes every even-degree, homogeneous polynomial function on X_I . Therefore,

$$\mathbb{C}[X_I]_{2n}^{C_I} = \mathbb{C}[X_I]_{2n}^{C_I^{\text{ref}}}$$

for all n . Consequently, if either C_I^{ref} is reducible or C_I^{ref} is irreducible and the Coxeter rank of C_I^{ref} is strictly less than the dimension of X_I , then $\dim \mathbb{C}[X_I]_2^{C_I} > 1 = \dim \mathbb{C}[V]_2^W$ and so ρ is not surjective.

It remains to consider the cases when $C_I \neq C_I^{\text{ref}}, C_I^{\text{ref}}$ is irreducible, and the Coxeter rank of C_I^{ref} equals $\dim X_I$.

If W is a dihedral group, then $C_I = C_I^{\text{ref}}$ for all I .

If W is of classical type and $C_I \neq C_I^{\text{ref}}$, then W is of type D_r and W_I has only components of type A . Suppose that this is the case. Then it follows from Howlett’s computations [How80] that whenever $C_I \neq C_I^{\text{ref}}$, either C_I^{ref} is reducible or the Coxeter rank of C_I^{ref} is strictly less than the dimension of X_I .

There are four cases when $C_I \neq C_I^{\text{ref}}, C_I^{\text{ref}}$ is irreducible, and the Coxeter rank of C_I^{ref} equals $\dim X_I$: either W is of type E_7 and W_I is of type A_2 , or W is of type E_8 and W_I is of type $A_2, A_1 \times A_2$, or A_4 .

Suppose W is of type E_7 and W_I is of type A_2 , or that W is of type E_8 and W_I is of type $A_1 \times A_2$. We show that $\dim \mathbb{C}[V]_4^W < \dim \mathbb{C}[X_I]_4^{C_I}$. Fix $f_2 \neq 0$ in $\mathbb{C}[V]_2^W$. Because the two smallest exponents of W are 1, 5 and 1, 7, respectively, it follows that $\mathbb{C}[V]_4^W$ is one dimensional with basis $\{f_2^2\}$. Since C_I^{ref} is of type A_5 in both cases, we have $\dim \mathbb{C}[X_I]_4^{C_I} = \dim \mathbb{C}[X_I]_4^{C_I^{\text{ref}}} = 2$.

Finally, suppose W is of type E_8 and W_I is of type A_2 or A_4 . We show that $\dim \mathbb{C}[V]_6^W < \dim \mathbb{C}[X_I]_6^{C_I}$. Fix $f_2 \neq 0$ in $\mathbb{C}[V]_2^W$. Since the two smallest exponents of W are 1 and 7, it follows that $\mathbb{C}[V]_6^W$ is one dimensional with basis $\{f_2^3\}$. Because C_I^{ref} is of type E_6 when W_I is of type A_2 and C_I^{ref} is of type A_4 when W_I is of type A_4 , we have $\dim \mathbb{C}[X_I]_6^{C_I} = 2$ in the first case and $\dim \mathbb{C}[X_I]_6^{C_I} = 3$ in the second. This completes the proof of the proposition. \square

To complete the proof of Theorem 2.1, we suppose that $\exp(C_X) = \exp(\mathcal{A}^X) \subseteq \exp(\mathcal{A})$ and show that $\rho: \mathbb{C}[V]^W \rightarrow \mathbb{C}[X]^{C_X}$ is surjective. Our argument is case-by-case, using the computation of $\exp(\mathcal{A}^X)$ by Orlik and Solomon [OS83], Howlett’s results in [How80], and some computer-aided computations using GAP [Sch97] for six cases when W is of exceptional type. For W of classical type, our argument is similar to that of Richardson [Ric87], but more streamlined, especially when W is of type D_r , because of our assumptions on \mathcal{A}^X .

As above, we may assume that W is irreducible and that $X = X_I$ for some proper, non-empty subset I of S . Then $W_X = W_I, N_X = N_W(W_I)$, and $C_X = C_I$. Notice that it follows from the assumption $\exp(C_I) \subseteq \exp(\mathcal{A})$ that C_I^{ref} is irreducible.

Suppose first that W is classical of type A_r, B_r , or D_r with $r \geq 1, r \geq 2$, and $r \geq 4$, respectively. Say W_I has m_i components of type A_i and a component of type B_j or D_j ,

where $j \geq 0$. In type A , we set $j = -1$. Set $k = j + \sum_i (i + 1)m_i$. Then k is minimal so that W_I may be embedded in a Coxeter group of type A_k, B_k , or D_k . The group C_I^{ref} is given as follows:

- $\prod_i A_{m_{i-1}} \times A_{r-k-1}$ if W is of type A_r ;
- $\prod_i B_{m_i} \times B_{r-k}$ if W is of type B_r ;
- $\prod_i B_{m_i} \times B_{r-k}$ if W is of type D_r and $j \neq 0$; and
- $\prod_{i \text{ even}} D_{m_i} \times \prod_{i \text{ odd}} B_{m_i} \times D_{r-k}$ if W is of type D_r and $j = 0$.

The exponents of \mathcal{A}^{X_I} have been computed by Orlik and Solomon in [OS83]. Set $l = \dim X_I$. Then $\exp(\mathcal{A}^{X_I})$ is given as follows:

- $\{1, 2, 3, \dots, l\}$ if W is of type A_r ;
- $\{1, 3, 5, \dots, 2l - 1\}$ if W is of type B_r ;
- $\{1, 3, 5, \dots, 2l - 1\}$ if W is of type D_r and $j \neq 0$; and
- $\{1, 3, 5, \dots, 2l - 3, l - 1 + \sum_i m_i\}$ if W is of type D_r and $j = 0$.

Type A_r . Suppose W is of type A_r . If $r - k - 1 > 0$, then, since C_I is irreducible, it must be that $m_i \leq 1$ for all i . Then $\exp(C_I) = \{1, 2, \dots, r - \sum_i (i + 1)\}$ and $\exp(\mathcal{A}^{X_I}) = \{1, 2, \dots, r - \sum_i i\}$, and so $r - \sum_i (i + 1) = r - \sum_i i$, which is absurd. Therefore, $r - k - 1 \leq 0$. Thus, $r \leq k + 1$ and W_I is of type A_d^m . In this case, $\exp(C_I) = \{1, 2, \dots, m - 1\}$, $\dim X_I = r - dm$, and $\exp(\mathcal{A}^{X_I}) = \{1, 2, \dots, r - dm\}$. Therefore, $m - 1 = r - dm$. We conclude that $\exp(C_I) = \exp(\mathcal{A}^{X_I}) \subseteq \exp(\mathcal{A})$ if and only if W_I is of type A_d^m , where r, d , and m are related by the equation $r + 1 = (d + 1)m$.

Now suppose that W_I is of type A_d^m with $r + 1 = (d + 1)m$. Then identifying W with the symmetric group S_{r+1} acting on \mathbb{C}^{r+1} , V with the subspace of \mathbb{C}^{r+1} consisting of all vectors whose components sum to zero, and W_I with the Young subgroup $S_{d+1}^m \subseteq S_{r+1}$ and taking the power sums as a set of fundamental polynomial invariants for S_{r+1} , it is straightforward to check that ρ is surjective.

Type B_r . Suppose that W is of type B_r with $r \geq 2$. Since C_I is irreducible, there is at most one value of i with $m_i > 0$. Suppose first that there is a value of i with $m_i > 0$. Say W_I has type $A_d^m \times B_j$. Then we must have $r - k = 0$ and so r, j, d , and m are related by $r = j + (d + 1)m$. In this case, C_I has type B_m and $\dim X_I = r - j - dm = m$. Thus, $\exp(C_I) = \{1, 3, \dots, 2m - 1\} = \exp(\mathcal{A}^{X_I})$. On the other hand, if $m_i = 0$ for all i , then W_I is of type B_j , C_I is of type B_{r-j} , $\dim X_I = r - j$, and $\exp(C_I) = \{1, 3, \dots, 2(r - j) - 1\} = \exp(\mathcal{A}^{X_I})$. We conclude that $\exp(C_I) = \exp(\mathcal{A}^{X_I}) \subseteq \exp(\mathcal{A})$ if and only if W_I is of type $A_d^m \times B_j$, where, if $m > 0$, then r, d, j , and m satisfy $r = j + (d + 1)m$.

Now suppose that W_I is of type $A_d^m \times B_j$ with $r = j + (d + 1)m$ if $m > 0$. We may consider W as signed permutation matrices acting on \mathbb{C}^r . Let x_1, \dots, x_r denote the coordinate functions on \mathbb{C}^r . Then $\mathbb{C}[V]^W = \mathbb{C}[x_1, \dots, x_r]^W = \mathbb{C}[f_2, f_4, \dots, f_{2r}]$, where f_{2p} is the p th elementary symmetric function in $\{x_1^2, \dots, x_r^2\}$. In case $m > 0$, we may choose coordinate functions $\{y_1, \dots, y_m\}$ on X_I so that C_I acts as signed permutations on the coordinates and the restriction map $\mathbb{C}[V] \rightarrow \mathbb{C}[X_I]$ is given by mapping $x_{p(d+1)+q}$ to y_p for $0 \leq p \leq m - 1$ and $1 \leq q \leq d + 1$, and x_t to zero for $t > r - j = (d + 1)m$. It is then easily checked that $\rho : \mathbb{C}[x_1, \dots, x_r]^W \rightarrow \mathbb{C}[y_1, \dots, y_m]^{C_I}$ is surjective. In case $m = 0$, we may take C_I to act on the first $r - j$ components

TABLE 1. Pairs (W, W_I) with W of classical or dihedral type, $\emptyset \neq I \neq S$, and $\exp(C_I) = \exp(\mathcal{A}^I) \subseteq \exp(\mathcal{A})$.

W	W_I	
A_r	A_d^m	$r + 1 = (d + 1)m$
B_r	$A_d^m B_j$	$m > 0 \Rightarrow r = j + (d + 1)m$
D_r	$A_d^m D_j$	$[j, m > 0 \Rightarrow r = j + (d + 1)m]$ or $[j = 0 \Rightarrow m \text{ odd} \wedge r = (d + 1)m]$
$I_2(r)$	A_1, \tilde{A}_1	$r \text{ even}$

of \mathbb{C}^r and so the restriction map $\mathbb{C}[V] \rightarrow \mathbb{C}[X_I]$ is given by evaluating x_{r-j+1}, \dots, x_r at zero. It is now easily checked that $\rho : \mathbb{C}[x_1, \dots, x_r]^W \rightarrow \mathbb{C}[x_1, \dots, x_{r-j}]^{C_I}$ is surjective.

Type D_r . Suppose that W is of type D_r with $r \geq 4$. In case $j \neq 0$, the argument for type B applies almost verbatim (B_j is replaced by D_j) and shows that $\exp(C_I) = \exp(\mathcal{A}^{X_I}) \subseteq \exp(\mathcal{A})$ if and only if W_I is of type $A_d^m \times D_j$, where, if $m > 0$, then r, d, j , and m satisfy $r = j + (d + 1)m$. In the case when $j = 0$, the arrangement \mathcal{A}^{X_I} is a Coxeter arrangement if and only if either $\sum_i m_i = 0$, in which case it is a Coxeter arrangement of type D_l , or $\sum_i m_i = l$, in which case it is a Coxeter arrangement of type B_l . Since $\sum_i m_i \neq 0$, we must have that $\sum_i m_i = l = r - \sum_i i m_i$ and \mathcal{A}^{X_I} is of type B_l . Thus, C_I^{ref} must be of type B_l and so W_I must be of type A_d^m , where d is odd and $r = (d + 1)m$. We conclude that if $j = 0$, then $\exp(C_I) = \exp(\mathcal{A}^{X_I}) \subseteq \exp(\mathcal{A})$ if and only if W_I is of type A_d^m , where d is odd and $r = (d + 1)m$.

Now suppose that W_I is of type $A_d^m \times D_j$, where, if $j, m > 0$, then $r = j + (d + 1)m$ and if $j = 0$, then d is odd and $r = (d + 1)m$. We may consider W as signed permutation matrices with determinant 1 acting on \mathbb{C}^r . Then $\mathbb{C}[V]^W = \mathbb{C}[x_1, \dots, x_r]^W = \mathbb{C}[f_2, f_4, \dots, f_{2r-2}, g_r]$, where f_{2p} is the p th elementary symmetric function in $\{x_1^2, \dots, x_r^2\}$ and $g_r = x_1 \cdots x_r$. The argument showing that ρ is surjective when W is of type B applies word for word to show that ρ is surjective in this case as well.

In order to determine the remaining cases when $\exp(C_I) = \exp(\mathcal{A}^I) \subseteq \exp(\mathcal{A})$, we fix a root system Φ for W . Then $\Phi \subseteq V^*$ and the choices of S and I determine a positive system and a closed parabolic subsystem denoted by Φ^+ and Φ_I , respectively. For α in Φ , we have $\alpha|_{X_I} \neq 0$ if and only if $\alpha \notin \Phi_I$.

If W_I is a maximal parabolic subgroup of W and $\exp(C_I) = \exp(\mathcal{A}^I) \subseteq \exp(\mathcal{A})$, then C_I is of type A_1 and acts as -1 on the one-dimensional space X_I . By [Bou68, ch. VI, § 1.1], $f_2 = \sum_{\alpha \in \Phi} \alpha^2$ is a non-zero polynomial in $\mathbb{C}[V]_2^W$. Fix β in $\Phi^+ \setminus \Phi_I$. Then $\{\beta|_{X_I}\}$ is a basis of X_I^* . If $g_2 = \beta|_{X_I}^2$, then $\mathbb{C}[X_I]^{C_I} = \mathbb{C}[g_2]$. Since $\alpha|_{X_I}$ is a non-zero multiple of $\beta|_{X_I}$ for α in $\Phi^+ \setminus \Phi_I$, it follows that $\rho(f_2)$ is a non-zero multiple of g_2 and so ρ is surjective.

Suppose that W is of type $I_2(r)$ and $|I| = 1$. We have observed above that if r is odd, then C_I is the trivial group, so $\exp(C_I) = \{0\}$ and $\exp(\mathcal{A}^I) = \{1\}$. On the other hand, if r is even, then $\exp(C_I) = \exp(\mathcal{A}^I) = \{1\}$ and $\exp(\mathcal{A}) = \{1, m - 1\}$ and so $\exp(C_I) = \exp(\mathcal{A}^I) \subseteq \exp(\mathcal{A})$.

Our computations when W is of classical or dihedral type are summarized in Table 1.

Finally, suppose that W is of exceptional type. The pairs (W, W_I) for which $\exp(C_I) = \exp(\mathcal{A}^I) \subseteq \exp(\mathcal{A})$ are given in Table 2. The notation is as in [OS83].

We have seen above that if W_I is maximal and $\exp(C_I) = \exp(\mathcal{A}^I) \subseteq \exp(\mathcal{A})$, then ρ is surjective. For the remaining six cases, A_2^2 in E_6 ; $(A_1^3)'$, $A_1^3 \times A_2$, and A_5' in E_7 ; and A_2 and \tilde{A}_2 in F_4 , the type of C_I is given in Table 3.

TABLE 2. Pairs (W, W_I) with W of exceptional type, $\emptyset \neq I \neq S$, and $\exp(C_I) = \exp(\mathcal{A}^I) \subseteq \exp(\mathcal{A})$.

W	W_I										
E_6	A_2^2	$A_1A_2^2$	A_5								
E_7	$(A_1^3)'$	$A_1^3A_2$	A_5'	$A_1A_2A_3$	A_2A_4	A_1A_5	A_6	A_1D_5	D_6	E_6	
E_8	$A_1A_2A_4$	A_3A_4	A_1A_6	A_7	A_2D_5	D_7	A_1E_6	E_7			
F_4	A_2	\tilde{A}_2	C_3	B_3	$A_1\tilde{A}_2$	\tilde{A}_1A_2					
G_2	A_1	\tilde{A}_1									
H_3	A_1A_1	A_2	$I_2(5)$								
H_4	A_1A_2	A_3	$A_1I_2(5)$	H_3							

TABLE 3. Triples (W, W_I, C_I) with W of exceptional type, $\emptyset \neq I$, $|I| < r - 1$, and $\exp(C_I) = \exp(\mathcal{A}^I) \subseteq \exp(\mathcal{A})$.

W	E_6		E_7		F_4	
W_I	A_2^2	$(A_1^3)'$	$A_1^3A_2$	A_5'	A_2	\tilde{A}_2
C_I	G_2	F_4	G_2	G_2	G_2	G_2

For these six cases, the fact that ρ is surjective was checked directly by implementing the following argument using GAP [Sch97] and the CHEVIE package [GHLMP96].

(1) For s in S , let α_s and ω_s denote the simple root in V^* and the fundamental dominant weight in V^* determined by s , respectively. Then $\{\omega_s \mid s \notin I\}$ is a basis of X_I^* and $\{\omega_s \mid s \notin I\} \cup \{\alpha_s \mid s \in I\}$ is a basis of V^* . This basis can be computed from the basis consisting of simple roots using the Cartan matrix of W . The restriction mapping $\mathbb{C}[V] \rightarrow \mathbb{C}[X_I]$ is then given by evaluating α_s at zero for s in I .

(2) Suppose that the exponents of W are $\{d_1 - 1, d_2 - 1, \dots, d_r - 1\}$, where $\{d_1 - 1, d_2 - 1, \dots, d_l - 1\}$ are the exponents of C_I . For $i = 1, 2, \dots, l$, define $f_i = \sum_{\alpha \in \Phi^+} \alpha^{d_i}$. Even though $\{f_1, \dots, f_l\}$ is not obviously algebraically independent, each f_i is a non-zero element in $\mathbb{C}[V]_{d_i}^W$.

(3) For $i = 1, 2, \dots, l$, express each f_i as a polynomial in $\{\omega_s \mid s \notin I\} \cup \{\alpha_s \mid s \in I\}$. Then set $\alpha_s = 0$ for s in I to get a polynomial $\rho(f_i)$ in $\mathbb{C}[X_I]_{d_i}^{C_I}$.

(4) Compute the Jacobian determinant of $\{\rho(f_1), \rho(f_2), \dots, \rho(f_l)\}$.

It turns out that in all cases, the Jacobian determinant above is non-zero and so it follows from [Spr74, Proposition 2.3] that $\mathbb{C}[X_I]^{C_I} = \mathbb{C}[\rho(f_1), \rho(f_2), \dots, \rho(f_l)]$. Therefore, ρ is surjective. This completes the proof of Theorem 2.1.

ACKNOWLEDGEMENTS

The authors acknowledge the financial support of a DFG grant for the enhancement of bilateral cooperation and the DFG-priority program SPP1388 ‘Representation Theory’. Parts of this paper were written during a stay of the authors at the Isaac Newton Institute for Mathematical Sciences in Cambridge during the ‘Algebraic Lie Theory’ Programme in 2009, and during a visit of the first author at the University of Bochum in 2010.

REFERENCES

- Bor81 W. Borho, *Über Schichten halbeinfacher Lie-Algebren*, Invent. Math. **65** (1981), 283–317.
- Bou68 N. Bourbaki, *Éléments de mathématique. Groupes et algèbres de Lie. Chapitres IV–VI*, in *Actualités scientifiques et industrielles, no. 1337* (Hermann, Paris, 1968).
- Bro98 A. Broer, *Decomposition varieties in semisimple Lie algebras*, Canad. J. Math. **50** (1998), 929–971.
- DL95 J. Denef and F. Loeser, *Regular elements and monodromy of discriminants of finite reflection groups*, Indag. Math. (N.S.) **6** (1995), 129–143.
- Dou99 J. M. Douglass, *The adjoint representation of a reductive group and hyperplane arrangements*, Represent. Theory **3** (1999), 444–456.
- GHLMP96 M. Geck, G. Hiß, F. Lübeck, G. Malle and G. Pfeiffer, *CHEVIE – A system for computing and processing generic character tables*, Appl. Algebra Engrg. Comm. Comput. **7** (1996), 175–210.
- How80 R. B. Howlett, *Normalizers of parabolic subgroups of reflection groups*, J. Lond. Math. Soc. (2) **21** (1980), 62–80.
- LS99 G. I. Lehrer and T. A. Springer, *Intersection multiplicities and reflection subquotients of unitary reflection groups. I*, Geometric Group Theory Down Under (Canberra, 1996) (de Gruyter, Berlin, 1999), 181–193.
- OS83 P. Orlik and L. Solomon, *Coxeter arrangements*, in *Singularities*, Proceedings in Symposium in Pure Mathematics, vol. 40 (American Mathematical Society, Providence, RI, 1983), 269–292.
- OT92 P. Orlik and H. Terao, *Arrangements of hyperplanes* (Springer, Berlin, 1992).
- OT93 P. Orlik and H. Terao, *Coxeter arrangements are hereditarily free*, Tôhoku Math. J. (2) **45** (1993), 369–383.
- Ric87 R. W. Richardson, *Normality of G -stable subvarieties of a semisimple Lie algebra*, in *Algebraic groups: Utrecht 1986*, Lecture Notes in Mathematics, vol. 1271 (Springer, Berlin, 1987), 243–264.
- Sch97 M. Schönert *et al.*, *GAP – Groups, Algorithms, and Programming – version 3 release 4*, RWTH Aachen (1997).
- Spr74 T. A. Springer, *Regular elements of finite reflection groups*, Invent. Math. **25** (1974), 159–198.
- Ter80 H. Terao, *Arrangements of hyperplanes and their freeness I*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **27** (1980), 293–320.

J. Matthew Douglass douglass@unt.edu

Department of Mathematics, University of North Texas, Denton, TX 76203, USA

Gerhard Röhrle gerhard.roehrle@rub.de

Fakultät für Mathematik, Ruhr-Universität Bochum, D-44780 Bochum, Germany