



Effect of using hydroxychloride as a copper source on performance, eggshell quality, tibia properties, mineral excretion, and antioxidant capacity of yolk in layer quails

Animal Research Paper

Cite this article: Gül ET, Olgun O, Kılınç G, Gökmen F, Yıldız A, Uygur V, Sevim B, Sarmiento-García A (2024). Effect of using hydroxychloride as a copper source on performance, eggshell quality, tibia properties, mineral excretion, and antioxidant capacity of yolk in layer quails. *The Journal of Agricultural Science* 1–8. <https://doi.org/10.1017/S0021859624000236>

Received: 4 December 2023

Revised: 27 March 2024



Accepted: 12 April 2024

Keywords:

antioxidant; copper; hydroxychloride; mineral excretion; quail

Corresponding author:

Ainhoa Sarmiento-García;
Email: asarmg00@usal.es

Esra Tuğçe Gül¹ , Osman Olgun¹, Gözde Kılınç², Fatih Gökmen³, Alpönder Yıldız¹, Veli Uygur⁴, Behlül Sevim⁵ and Ainhoa Sarmiento-García^{6,7} 

¹Department of Animal Science, Agriculture Faculty, Selcuk University, 42130, Konya, Türkiye; ²Department of Food Processing, Suluova Vocational Schools, Amasya University, 05500 Amasya, Türkiye; ³Department of Soil Science and Plant Nutrition, Agriculture Faculty, Iğdır University, Iğdır, Türkiye; ⁴Department of Soil Science and Plant Nutrition, Agriculture Faculty, Applied Sciences University of Isparta, Isparta, Türkiye; ⁵Department of Food Processing, Aksaray Technical Sciences Vocational School, Aksaray University, 68100, Aksaray, Türkiye; ⁶Área de Producción Animal, Departamento de Construcción y Agronomía, Facultad de Ciencias Agrarias y Ambientales, Universidad de Salamanca, 37007, Salamanca, Spain and ⁷Estación Tecnológica de la Carne, Instituto Tecnológico Agrario de Castilla y León (ITACyL), 37770, Guijuelo, Salamanca, Spain

Abstract

This research aimed to examine the impact of varying levels of dietary copper (Cu) hydroxychloride on the performance, egg quality, yolk antioxidant capacity, tibia traits, and mineral excretion in laying quails. 125 female 10-week-old quails were randomly distributed into five experimental groups with five replicates, each consisting of five quails. Five experimental iso-nitrogenous and isoenergetic diets were designed to contain different Cu hydroxychloride (54% Cu) levels at 7.20 (basal diet), 15, 30, 45 and 60 mg/kg respectively. Quails were fed with trial diets for 12 weeks. Performance, egg production, eggshell quality, and biomechanical traits of the tibia were not impacted ($P > 0.005$) by variations in dietary Cu levels. Yolk antioxidant capacity, measured as yolk DPPH value, exhibited an increase ($P < 0.01$) in the high-dose group (60 mg/kg). Regarding tibia mineral concentration, Cu concentration decreased linearly ($P < 0.001$) with increasing Cu level, manganese and zinc content recorded the highest values in quails that had received 60 mg/kg Cu in the diet ($P < 0.01$), while the lowest phosphorus content was described for 45 and 60 mg/kg. Contrarily, increases ($P < 0.01$) in dietary Cu resulted in raised faecal Cu content, while phosphorus, manganese, and zinc, were reduced when Cu was added. It can be inferred that adding Cu to the diet of laying quails would not be necessary, which in turn decreases Cu excretion and prevents substantial environmental harm.

Introduction

Copper (Cu) is an indispensable and crucial trace mineral for livestock (Olukosi *et al.*, 2019). Cu is involved in several physiological processes that influence growth development, stimulate the immune system against infections, red blood cell production, repair damaged tissues, and eliminate reactive oxygen species which enhances antioxidant capacity (Mroczek-Sosnowska *et al.*, 2016; Hill and Shannon, 2019; Abdullah *et al.*, 2022). These improvements are linked to Cu's role in the development and activity of certain metalloenzymes (such as superoxide dismutase, ceruloplasmin or cytochrome oxidase), which are involved in several metabolism processes including cellular respiration, red blood formation or glucose and cholesterol metabolism (Scott *et al.*, 2018). For these reasons, poultry diets frequently incorporate Cu supplementation (Olukosi *et al.*, 2019; El Sabry *et al.*, 2021).

The National Research Council (1994) suggests incorporating Cu into the diet of laying quails at a rate of 5 mg/kg. However, standard poultry feeding practices include high doses of Cu (150–200 mg/kg) to improve certain aspects of avian production such as performance (Olgun and Aygun, 2017; Wu *et al.*, 2019; Deo *et al.*, 2023), eggshell quality (Kaya *et al.*, 2018; Cufadar *et al.*, 2022), and antioxidant capacity (Kara *et al.*, 2021; Abdullah *et al.*, 2022; Ali, 2018). Despite the improvements observed, an excess of dietary Cu may potentially hinder the bioavailability of specific minerals like iron or calcium and could pose a hazard to the environment due to increased excretion in faeces and urine (Kara *et al.*, 2021; Ibrahim *et al.*, 2022). Some countries have already limited the dietary content of Cu in species such as pigs to levels of 75–100 mg/kg of feed (Kara *et al.*, 2021). Therefore, it is necessary to find safe sources and doses of Cu that improve animal production without compromising the environment (Lin *et al.*, 2020).

© The Author(s), 2024. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



Cu supplements are generally added to the avian diet such as ionic forms. Ionic sources such as Cu sulphate (CuSO₄), are extremely sensitive in contact with water and can be degraded readily both in the feed and in the intestinal tract. This process decreases its absorption in the small intestine due to its interaction with other nutrients (including vitamins, enzymes, or fats), and consequently increases its excretion into the environment (Olukosi *et al.*, 2019; Deo *et al.*, 2023). Hence, the use of new mineral additives has attracted the attention of the poultry industry in recent years, as an alternative to traditional trace mineral sources (Olukosi *et al.*, 2018; Lin *et al.*, 2020). Cu hydroxychloride is an organic salt, which is chemically more stable and bioavailable than Cu sulphate, which leads to enhanced absorption and consequently a reduction in excretion into the environment (Olukosi *et al.*, 2019).

Quail production has increased in recent decades, mainly due to their ease of management and high production rates (Sarmiento-García *et al.*, 2023). Nonetheless, unlike conventional avian species, fewer investigations have been carried out concerning the nutritional aspects of laying quails and are generally based on the nutritional values reported by the National Research Council (1994) 30 years ago (Olgun *et al.*, 2022). To the best of our knowledge, the number of studies on the Cu requirements of laying quails is quite limited, and even more so when alternative sources such as hydroxychloride are involved. Therefore, the objective of this study is to investigate how different levels of hydroxychloride copper in laying quail diets affect various parameters, such as performance, eggshell quality, yolk antioxidant capacity, tibia characteristics, and mineral excretion.

Materials and methods

Ethical approval

The following study was performed using farm animals; for this reason, there were no special requirements for keeping experimental animals. However, it is noteworthy that the criteria outlined in the European Animal Protection Policy (EPCEU, 2010) were strictly adhered to during the entire trial period.

Experimental design

The study was conducted according to a completely randomized design and took place on a farm in Selçuklu (Konya, Türkiye) at coordinates 38°1'36"N, 32°30'45"E. On arrival, one hundred-twenty-five female Japanese quails (*Coturnix coturnix Japonica*), were weighed (244 ± 12.3 g) and randomly allocated into five treatment groups with five replicates. All pens were standardized to identical size (30 × 45 cm) and upheld in well-ventilated, hygienic, and sanitized conditions. The ambient room temperature was controlled at 22°C (± 2.0), with a 16-h lighting schedule programme imposed. Additionally, each enclosure was outfitted with separate feeders and drinkers, allowing unrestricted access to both food and water.

Throughout the 12-week duration, all quails were provided with the identical basal diet comprising corn and soybean meal, characterized by a crude protein content of 200 g/kg, metabolizable energy of 12.14 MJ/kg, and copper content of 7.20 mg/kg, as detailed in Table 1. The basal diet containing 7.20 mg/kg of copper was employed for the diet devoid of supplemental hydroxychloride copper (54% Cu). According to the literature reviewed, the hydroxychloride Cu was incorporated into the basal diet at the

Table 1. Basal diet and its nutrient content (as fed)

Ingredients	g/kg	Nutrient composition	g/kg
Corn	544.0	Metabolizable energy (MJ/kg)	12.14
Soybean meal	344.0	Crude protein	200.1
Soybean oil	36.5	Crude fibre	28.3
Limestone	56.0	Crude fat	58.4
Dicalcium phosphate	11.4	Moisture	128.3
Salt	3.5	Lysine	10.9
Premix ^a	2.5	Methionine	4.5
DL-methionine	2.1	Cystine	3.7
Total	1000.0	Calcium	25.0
		Total phosphorus	6.4
		Available phosphorus	3.5
		Copper (mg/kg)	7.20

^aPremix (vitamin-mineral mixture) as contained per kg feed: vitamin A, 20000 IU; vitamin D3, 10000 IU; vitamin E, 125 mg; vitamin K3, 5 mg; vitamin B12, 0.0275 mg; biotin, 0.30 mg; folic acid, 2.5 mg; nicotinic acid, 112.5 mg; pantothenic acid, 37.5 mg; pyridoxine, 3.75 mg; riboflavin, 10 mg; thiamin, 5 mg; iodine, 3 mg; iron, 50 mg; manganese, 60 mg; zinc, 50 mg; selenium, 0.75 mg.

expense of corn to meet the desired concentration of Cu for the corresponding treatment at inclusion rates of 15, 30, 45 and 60 mg/kg Cu. The mash-formulated basal diet was formulated with the nutrient requirements recommended for laying quails according to the National Research Council (1994), except for Cu. The chemical composition of the basal diet was analysed using the methods specified by AOAC (2006). The basal diet's chemical composition was determined by subjecting samples to incineration and drying to ascertain ash content (method 942.05 analysing protein and fat levels using Kjeldahl (method 990.03) and Soxhlet techniques (method 2003.06), and analysing moisture content (method 2001.12) through drying at 105°C. Table 1 outlines the chemical composition and ingredients of the basal diet.

Evaluation of performance indicators and egg production

To establish performance parameters, all experimental quails ($n = 125$) underwent individual weighing at the beginning and conclusion of the study (± 0.01 g). Subsequently, changes in body weight (g) were determined based on the weight difference. Feed intake (g/quail/day) was calculated following the methodology outlined by Olgun *et al.* (2022) which involved monitoring the total feed dispensed and the remaining amount in the feed boxes. Egg production was evaluated by dividing the daily egg count by the total laying quails and multiplying the result by 100, expressing the result as per egg/100 quails. Additionally, every egg collected during the final three days of the study underwent weighing using a high-precision balance (±0.01 g) to ascertain egg weight. Egg mass (g/quail/day) and feed conversion ratio were determined according to the procedures delineated by Sarmiento-García *et al.* (2023).

Assessment of eggshell quality parameters

A total of 300 eggs were collected and sent to the Egg Quality Laboratory at the Faculty of Agriculture, Selcuk University,

Konya, Turkey, for evaluation of eggshell quality. Eggs gathered during the final three days of the trial were assessed for both internal and external quality parameters under ambient temperature conditions. Throughout the experiment, any instances of broken, damaged, or cracked eggs were noted and expressed as a percentage of the total egg count ($n = 300$). Shell strength was measured using the Egg Force Reader (Orka Food Technology Ltd., Ramat Hasharon, Israel) applied to the blunt part of the egg. Eggshell thickness was determined utilizing a micrometre (Mitutoyo, 0.01 mm, Japanese), with measurements taken at three different locations on the shell (equator, blunt, and pointed), and subsequently averaged. Relative eggshell weight was computed by weighing the cleaned and dried shells and dividing them by their respective egg weights.

Determination of yolk MDA and DPPH values

Malondialdehyde (MDA) and 1-diphenyl-2-picrylhydrazyl (DPPH) concentrations were determined on 100 fresh eggs to establish the lipid peroxidation of the yolk. The thiobarbituric acid reactive substances (TBARS) test was carried out to establish MDA value using a modified method reported by Kilic and Richards (2003) and Sarmiento-García *et al.* (2021), where each sample was measured in triplicate. MDA concentration was quantified by measuring absorbance at 530 nm wavelength using a spectrophotometer (Perkin Elmer, USA), based on a blank curve of MDA containing 1 ml of trichloroacetic acid (TCA) solution (7.5% TCA, 0.1% EDTA, 0.1% Propyl gallate) and 1 ml of thiobarbituric acid (TBA) solution (0.02 M). The results were reported as $\mu\text{mol MDA/kg}$ yolk.

The antioxidant capacity of the hydrolysates obtained was checked using a modified technique described by Sacchetti *et al.* (2005) and Olgun *et al.* (2022), which relies on the scavenging effect of DPPH radicals. Each sample was measured in triplicate to establish the average value. Absorbance at 517 nm was measured using a spectrophotometer (Perkin Elmer precisely UV/VIS Spectrometer) and compared against a blank curve substituted with 95% ethanol. Equation (1) was utilized to assess the scavenging effect:

$$\text{DPPH values} = \frac{[(\text{control absorbance} - \text{sample absorbance})]}{\text{control absorbance}} \times 100$$

Assessment of bone biomechanical properties

As the experiment ended, one female quail from each replicate ($n = 25$) was randomly euthanized via cervical dislocation. The right tibia was then removed, cleaned of tissue and cartilage, and stored at -20°C until the determination of bone biomechanical properties and mineral concentration. Before the examination, the samples were allowed to reach room temperature over 6 h in a controlled air environment. Cortical bone thickness, cortical bone cross-sectional area, shear force, and shear stress of bone were assessed according to protocols outlined by Gül *et al.* (2022), Armstrong *et al.* (2002), and Wilson and Ruszler (1996).

Analysis of mineral content in faeces and tibia

The mineral levels in faecal ($n = 25$) and tibia tissue ($n = 25$) samples were analysed using a wet digestion method following the

protocol by Olgun and Yıldız (2014). Samples from faeces and tibia, previously dried at 105°C for 24 h, were weighed (0.3 ± 0.01 g) and placed on Teflon-coated digestion plates (Milestone, Sorisole, Bergamo, Italy). A mixture of nitric acid (5 ml, 63.01 M) and perchloric acid (3 ml, 70%) was added to each plate. The samples were then heated in a microwave oven at 190°C for 40 min (CEM Corp, 3100 Smith Farm Road, Matthews, NC, USA). After cooling, the resulting supernatant solution was diluted with distilled water to 50 ml. Subsequently, 0.1 g of each ash sample was weighed to determine the mineral content, including concentrations of copper, calcium, phosphorus, manganese and zinc, using inductively coupled plasma atomic emission spectrometry (ICP-OES) with a Thermo Scientific 7200 analyser (ThermoFisher Scientific, Waltham, United States).

Statistical analysis

The data were analysed using analysis of variance (ANOVA) in SPSS 22.0 software (SPSS Inc., Chicago, IL, USA). Each cage replicate was considered as the experimental unit, and values presented in the Tables are displayed as means with Standard error means (S.E.M). Statistical significance was set at $P < 0.05$. Linear, quadratic, and cubic regressions were employed to examine the relationship between the dependent variable and increasing copper levels across all parameters.

Results

Performance indicators and egg production

Table 2 illustrates the effects of dietary Cu levels on the performance traits of laying quails. No statistically significant differences in performance traits were found among the experimental groups ($P > 0.05$). The range of values for these parameters was as follows: final body weight (265–277 g), body weight gain (22–32 g), egg production (90.4–92.1 per egg/100 quails), egg weight (12.6–13.3 g), egg mass (11.4–12.1 g/quail/day), feed intake (31.4–33.7 g/quail/day), and feed conversion ratio (2.70–2.85).

Eggshell quality

The eggshell quality parameters are expressed as damaged egg, egg-breaking strength, shell ratio, and shell thickness, as shown in Table 3. The effect of Cu supplementation in the diet of laying quails appeared to have no impact on eggshell parameters, as all values were similar ($P > 0.05$). Those values ranged from 0.2 to 1.6 per egg/100 eggs for damaged eggs, 12.0 to 12.9 N for egg-breaking strength, 8.18 to 8.72 g shell/ 100 g egg for shell ratio and 222 to 230 μm for egg-shell thickness.

Yolk DPPH and MDA values

The oxidative stability of the yolk measured as MDA and DPPH value is shown in Table 4. The MDA value was not impaired ($P > 0.05$) by Cu supplementation. Contrarily, a significant effect was described for the DPPH value, which significantly increased ($P < 0.01$) at 60 mg/kg of Cu compared to the rest of the experimental groups.

Tibia biomechanical traits

Data on tibia biomechanical properties are shown in Table 5. Cortical bone thickness (0.323–0.378 mm), cortical bone cross-

Table 2. Effect of dietary Cu levels on performance in laying quails ($n = 125$)

Parameters	Dietary Cu levels (mg/kg)					S.E.M.	P-value	Regression		
	7.20	15	30	45	60			L	Q	C
Initial body weight (g)	243	249	246	242	243.63	2.5	0.924	0.732	0.846	0.421
Final body weight (g)	269	271	277	265	274	3.5	0.887	0.910	0.903	0.451
Body weight gain (g)	26	22	32	24	30	1.9	0.512	0.490	0.972	0.741
Egg production (per egg/100 quails)	91.2	92.1	91.7	90.9	90.4	0.69	0.960	0.570	0.706	0.733
Egg weight (g)	13.0	12.9	13.0	12.6	13.3	0.14	0.587	0.799	0.276	0.352
Egg mass (g/quail/day)	11.9	11.9	11.9	11.4	12.1	0.16	0.786	0.939	0.514	0.350
Feed intake (g/quail/day)	32.4	33.7	32.1	31.4	33.0	0.43	0.537	0.698	0.430	0.156
Feed conversion ratio	2.73	2.85	2.70	2.74	2.74	0.031	0.683	0.667	0.879	0.507

S.E.M., Standard error means; L, Linear; Q, Quadratic; C, Cubic.

sectional area (1.22–1.30 mm²), shear force (131–154 N), and shear stress (100–119 N/mm²) were not affected by the Cu level ($P > 0.05$). Nevertheless, a linear regression ($P < 0.05$) was set for cortical bone thickness, which was linearly reduced from 0.378 to 0.323 mm, with increasing Cu content in the diet.

Tibia mineralization

Table 6 summarizes the tibia mineral concentration of laying quails in response to rising levels of Cu supplementation. Tibia Cu content decreased linearly ($P < 0.01$) with increasing Cu levels in the diet, resulting in higher values in the non-supplemented quails than in the quails receiving 30, 45 and 60 mg/kg of Cu. Contrarily, as the dietary Cu level increased ($P < 0.01$) the tibia manganese content linearly rose, recording the highest value at 60 mg/kg Cu than those reported for 15 mg/kg Cu diet and non-supplemented diets. Similarly, zinc concentration increased with enhanced Cu supplementation levels, and this increase was significant ($P < 0.01$) at 60 mg/kg Cu compared to other levels. The effect of Cu supplementation on tibia calcium content appeared to be not significant ($P > 0.05$). There was a significant effect ($P < 0.01$) of Cu supplementation on tibia phosphorus content, quails receiving 15 and 30 mg/kg diets had considerably higher P levels than the quails receiving 45 and 60 mg/kg Cu.

Mineral excretion

Faecal mineral content (Cu, manganese, zinc, calcium and phosphorus concentration) in response to the Cu diets are presented in

Table 7. Increasing the dietary Cu level from 7.20 to 60 mg/kg resulted in a linear increase in faecal Cu level from 18 to 56 mg/kg. Contrarily, the study indicated that the groups supplemented with Cu had considerably lower ($P < 0.01$) zinc and manganese (280–281 mg/kg) faecal content than the non-supplemented group. Faecal calcium content was not affected by dietary Cu level ($P > 0.05$), whereas a quadratic regression ($P < 0.05$) was observed with minimum values at 15 mg/kg of Cu. Faecal phosphorus was found to be significantly higher in the non-supplemented group compared to the laying quails supplemented with 15 and 30 mg/kg Cu.

Discussion

Trace elements, including Cu, are crucial in the diet of poultry to ensure optimum performance, since these elements due to these elements are involved in several biochemical processes (Olukosi *et al.*, 2019). However, the optimal dose of Cu supplementation to ensure optimal performance without compromising the environment is still controversial (Cufadar *et al.*, 2022). In this study, the addition of hydroxychloride copper did not yield any discernible impact on performance parameters. Comparable results have been reported by previous researchers in studies involving laying quails (Kara *et al.*, 2021; Cufadar *et al.*, 2022) and broilers (Nguyen *et al.*, 2020). On egg production in laying hens, similar to the current findings, Olukosi *et al.* (2019) observed no effect of Cu hydroxychloride supplementation, which is in line with that described by Cufadar *et al.* (2022) and Kara *et al.* (2021) when organic Cu was added to the diet of quails. Olukosi *et al.*

Table 3. Effect of dietary Cu levels on eggshell quality ($n = 300$) in laying quails

Parameters	Dietary Cu levels (mg/kg)					S.E.M.	P-value	Regression		
	7.20	15	30	45	60			L	Q	C
Damaged egg (per egg/100 egg)	1.7	0.5	0.3	1.5	0.2	0.32	0.452	0.440	0.834	0.097
Eggshell-breaking strength (N)	12.4	12.9	12.0	12.2	12.8	0.14	0.155	0.885	0.101	0.139
Relative eggshell weight (g shell/100 g egg)	8.72	8.29	8.18	8.48	8.41	0.073	0.172	0.539	0.095	0.070
Eggshell thickness (µm)	222	227	225	230	224	1.2	0.186	0.366	0.100	0.650

S.E.M., Standard error means; L, Linear; Q, Quadratic; C, Cubic.

Table 4. Effect of dietary Cu levels on DPPH and MDA values of yolk ($n = 100$) in laying quails

Parameters	Dietary Cu levels (mg/kg)*					S.E.M.	P-value	Regression		
	7.20	15	30	45	60			L	Q	C
MDA value ($\mu\text{mol MDA/kg}$)	1.7	2.4	1.4	2.4	1.9	0.16	0.192	0.776	0.912	0.691
DPPH (% reducing)	7.3	7.2	6.6	7.5	9.1	0.19	<0.001	<0.001	<0.001	0.707

S.E.M., Standard error of means; L, Linear; Q, Quadratic; C, Cubic; DPPH, 2,2-diphenyl-1-picrylhydrazyl; TBARS, thiobarbituric acid reactive substances.

(2019) proposed that the absence of differences could be attributed to the hens being at peak production, a circumstance consistent with the age of the quails in this study. In this research, the concentration of Cu in the basal diets approached the minimal requirement (6 mg/kg) reported for quails by the National Research Council (1994). Therefore, the comparative performance and egg production traits of quails fed a diet without copper supplementation imply that the basal diet, primarily composed of corn, did not exhibit a significant Cu deficiency. Therefore, based on these results and as pointed out by previous authors (Nguyen *et al.*, 2020; Cufadar *et al.*, 2022), it could be suggested that additional dietary Cu supplementation is not necessary to ensure optimal growth or egg production in laying quails.

Egg-shell quality remains a critical problem for avian production which accounts for large economic losses for the avian industry. Therefore, it is important to ensure that shell quality parameters are preserved (Gül *et al.*, 2022). The current research found that the hydroxychloride Cu supplementation did not affect the eggshell quality (in terms of damaged egg, eggshell breaking strength, relative eggshell weight, and eggshell thickness). Similarly, Olukosi *et al.* (2019) described an absence of differences in eggshell thickness and relative eggshell weight of the eggs from hydroxychloride Cu-fed hens, which follows those described by El Sherif *et al.* (2019). These findings align with those reported by Kara *et al.* (2021) who stated that the addition of propionate Cu (organic form) at the level of 80 mg/kg did not affect eggshell quality. However, Olgun *et al.* (2020) reported that administration of 20 mg/kg organic Cu to the basal diet (35.95 mg/kg Cu) decreased eggshell breaking strength and shell index, but conversely increased eggshell thickness. Similarly, Cufadar *et al.* (2022) demonstrated that incorporating high levels of organic copper (150 or 225 mg/kg) into the diets of laying quails enhanced eggshell thickness and eggshell ratio without affecting eggshell breaking strength. As can be perceived, there are several discrepancies between previous studies. Factors such as laying age, basal Cu doses, or the supplementation with organic, inorganic, or chelated Cu sources, among others, can impact the perceived effects. Nonetheless, the consensus is that using organic or hydroxychloride Cu sources rather than inorganic sources could result in improved eggshell quality, which may or may not, be projected in eggshell thickness (Olukosi *et al.*, 2019). In any case, the lack of differences in eggshell traits suggests that the Cu provided in the basal diet would be enough to ensure the eggshell quality of the laying quails.

Cu serves as a vital component of the antioxidant system, enhancing the activity of antioxidant enzymes (Kara *et al.*, 2021), but, excess of dietary Cu could act as a prooxidant due to the formation of hydroxyl radicals leading to lipid peroxidation (Ali, 2018; Lin *et al.*, 2020; Nguyen *et al.*, 2022). In the present investigation, the addition of Cu hydroxychloride at the maximum level (60 mg/kg) resulted in an improvement of the

DPPH value in comparison to the rest of the experimental diets, without promoting oxidative stress in any case. Those results are partially following those reported by Nguyen *et al.* (2022). These results suggest that Cu hydroxychloride, even at high doses, would have a reduced release of Cu ions, resulting in a significantly reduced reactivity potential and less production of reactive oxygen species leading to less oxidation (Lin *et al.*, 2020). Scarce data about the effect of Cu hydroxychloride dietary supplementation on the antioxidant capacity of the yolk have been reported. Most previous studies have focused on the effect of other dietary sources of Cu (inorganic or organic) or the combination of hydroxychlorinated minerals on the antioxidant capacity of different tissues. For example, Kara *et al.* (2021) reported that the administration of Cu proteinate at the level of 80 mg/kg reduced yolk MDA. Abdullah *et al.* (2022) noted an increase in serum SOD activity and a decrease in MDA values in broilers that had received Cu nanoparticles in the diet. Similar outcomes have been described by Wu *et al.* (2019), who found increased antioxidant protection (measured by SOD and GSH-x activities) in the serum of broiler chickens supplemented with organic Cu, which agrees with those described by Ali (2018). These authors found a reduction of serum MDA value in quails fed with dietary-containing Cu (cupric sulphate pentahydrate) regardless of the doses. It would appear that, overall, dietary Cu would have some antioxidant effect on different tissues. On the other hand, El Sherif *et al.* (2019) proposed that the total antioxidant capacity in the serum of laying hens was enhanced as the dietary Cu sulphate was reduced, while Nguyen *et al.* (2022) revealed that including 200 mg Cu/kg to the broiler diet did not affect liver MDA values. Nonetheless, it is important to note that the animals in the El Sherif *et al.* (2019) experiment were reared under stressful environmental conditions. The differences among studies are probably due to the dosage, the source of Cu involved, the age of the animals, and the environmental conditions.

The absorption of trace elements is crucial for the development of the skeletal system, potentially affecting overall bone health. In this sense, Cu is an important mineral to preserve the skeletal system as it serves as a cofactor for lysyl oxidase. This enzyme binds collagen and elastin proteins and is necessary to ensure structural integrity (Nguyen *et al.*, 2020). Furthermore, the antioxidant capacity of Cu has a major role in neutralizing the radicals which stimulate the osteoclasts. Hence, a deterioration of the biomechanical parameters of the tibia suggests a deficit in Cu requirements. As previously stated, no alteration of tibial biomechanical traits linked to dietary Cu level was described, which agrees with the results of previous reports (Nguyen *et al.*, 2020; dos Santos *et al.*, 2023). This observation supports the hypothesis that the Cu content of the basal diet would be enough to satisfy the requirements of laying quails.

Tissue mineral contents are commonly determined to assess mineral availability in livestock (Lin *et al.*, 2020) and are a good

Table 5. Effect of dietary Cu levels on biomechanical traits of tibia ($n = 25$) in laying quails

Parameters	Dietary Cu levels (mg/kg)					S.E.M.	P-value	Regression		
	7.20	15	30	45	60			L	Q	C
Cortical bone thickness (mm)	0.378	0.340	0.333	0.324	0.323	0.0074	0.102	0.022	0.209	0.449
Cortical bone cross-sectional area (mm ²)	1.42	1.22	1.29	1.23	1.30	0.027	0.116	0.314	0.094	0.368
Shear force (N)	144	131	132	131	154	5.5	0.616	0.510	0.167	0.931
Shear stress (N/mm ²)	102	110	100	108	119	4.4	0.690	0.317	0.477	0.651

S.E.M., Standard error means; L, Linear; Q, Quadratic; C, Cubic.

Table 6. Effect of dietary Cu levels on mineral contents of tibia ($n = 25$) in laying quails

Parameters	Dietary Cu levels (mg/kg)					S.E.M.	P-value	Regression		
	7.20	15	30	45	60			L	Q	C
Copper (mg/kg)	3.1	2.6	2.3	1.9	1.7	0.12	<0.001	<0.001	0.272	0.734
Manganese (mg/kg)	9.0	10.7	12.0	12.0	14.7	0.55	0.008	0.001	0.941	0.202
Zinc (mg/kg)	290	303	297	290	338	4.3	<0.001	<0.001	0.002	0.001
Calcium (g/kg)	210.9	237.6	222.9	222.3	229.4	0.63	0.771	0.721	0.827	0.322
Phosphorus (g/kg)	157.7	173.9	176.5	138.4	137.8	0.46	0.004	0.004	0.073	0.016

S.E.M., Standard error means; L, Linear; Q, Quadratic; C, Cubic.

marker of bone mineralization (Nguyen *et al.*, 2020). Interestingly, a reduction in the Cu content of the tibia was reported as the dietary Cu level was increased. This fact could be related to its role in phytase inhibition, vitamin degradation, and increased oxidation which could indirectly result in reduced bioavailability and decreased absorption of Cu in the tibia, as reported by Lin *et al.* (2020). The increase of Cu levels in faeces confirms these results. An increase in tibial zinc and manganese contents was only recorded in quails that had received the highest dietary Cu levels. At the same time, phosphorus showed the lowest values when dietary Cu levels were increased (45 and 60 mg/kg) which is by those described by Nguyen *et al.* (2020). It has been described that high doses of Cu could impair the bioavailability of phosphorus by the formation of an insoluble phytate-C complex as occurs in this research. Moreover, dietary Cu did not alter the deposition of calcium which is following those reported by Nguyen *et al.* (2020). However, according to the previous literature, the effects of dietary Cu on the tibia mineral content are controversial. Dos Santos *et al.* (2023), described that dietary

Cu in hydroxychloride form (at a level of 150 mg/kg) did not affect the Cu and phosphorus contents of the tibia in broilers, but reduced the calcium and zinc, while Nguyen *et al.* (2020) found that the majority of tibia minerals assessed remained unaffected by Cu dietary supplementation. Lin *et al.* (2020) suggested that the variations observed in previous research could stem from differences in factors such as the ages of the animals, forms and levels of copper supplementation, duration of the experiments, as well as the composition of the diets, among other factors. Nevertheless, our findings suggest that non-supplemented Cu diets are enough to ensure normal bone mineralization and absorption processes.

In the current study, dietary supplementation with Cu resulted in increased faecal excretion of Cu but decreased excretion of manganese, zinc, and phosphorus. As far as we know, there have been no studies investigating the effect of dietary copper on faecal mineral excretion in laying quails or hens. Somewhat congruent with these observations, Nguyen *et al.* (2022) observed that adding hydroxychloride Cu at high doses (200 mg/kg) to the

Table 7. Effect of dietary Cu levels on mineral contents of faecal ($n = 25$) in laying quails

Parameters	Dietary Cu levels (mg/kg)					S.E.M.	P-value	Regression		
	7.20	15	30	45	60			L	Q	C
Copper (mg/kg)	18	24	39	49	56	3.0	<0.001	<0.001	0.001	0.468
Manganese (mg/kg)	343	281	280	281	281	6.3	<0.001	0.001	0.003	0.010
Zinc (mg/kg)	304	242	241	247	247	5.7	<0.001	0.001	<0.001	0.001
Calcium (g/kg)	49.6	42.8	44.4	42.9	50.5	0.13	0.172	0.675	0.032	0.880
Phosphorus (g/kg)	21.4	17.8	17.9	20.0	19.6	0.04	0.014	0.843	0.026	0.005

S.E.M., Standard error means; L, Linear; Q, Quadratic; C, Cubic.

broiler diet' resulted in an increased Cu in distal ileum content, but did not influence the excretion of other minerals (calcium, phosphorus, zinc, and manganese). Mineral absorption is a complex process, and microminerals can use the same transport medium to be assimilated by the cell (dos Santos *et al.*, 2023). Consequently, even if the mineral content is equal, variations in the dietary components or the presence of chelating agents may lead to different results in the absorption process (Lin *et al.*, 2020). Hence, additional study is warranted to validate the mechanism underlying mineral excretion.

Conclusions

Copper plays a crucial role in maintaining, developing, and promoting the growth and overall health of animals. However, there seems to be no consensus on the appropriate dose to improve the development of laying quails without compromising the environment. This study revealed that while dietary intake of hydroxychloride copper at a level of 60 mg/kg had a beneficial impact on egg yolk antioxidant status, the basal diet composed of corn and containing 7.20 mg/kg copper could be sufficient to meet the requirements for animal development, egg production traits, egg quality, and bone biomechanical properties of laying quails, without compromising any of these parameters. It should be noted that reducing the dietary inclusion of Cu to basal levels would result in less Cu excretion to the environment, which would contribute to its protection.

Acknowledgements. We acknowledge the contributions of Dr Nicholas Collins (School of Agriculture Food and Wine, The University of Adelaide, Adelaide, SA, Australia) who supervised these projects, and thank all the technical staff involved in the field experiments.

Author contributions. Esra T. Gül, Osman Olgun, Alpönder Yıldız, Behlül Sevim and Ainhoa Sarmiento-García contributed to the conceptualization, investigation, methodology, writing – original draft and writing – review and editing. Gözde Kılınç contributed antioxidant analysis. Veli Uygun and Fatih Gökmen contributed mineral analysis. Esra T. Gül, Osman Olgun and Ainhoa Sarmiento-García contributed to supervision, review and editing. All authors have read and approved the final version of the manuscript.

Funding statement. This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Competing interests. None.

Ethical standards. The authors confirm the ethical policies of the journal, as noted on the journal's author guidelines page. The European National Research Council's guidelines for the Care and Use of Laboratory Animals were followed.

References

- Abdullah SS, Masood S, Zaneb H, Rabban I, Akba J and Kuthu ZH (2022) Effects of copper nanoparticle supplementation on growth, intestinal morphology, and antioxidant status in broilers exposed to cyclic cold stress. *Eminates Journal of Food and Agriculture* **34**, 436–445.
- Ali RAM (2018) Physiological and antioxidant responses of Japanese quail to dietary copper supplementation. *Journal of Animal and Poultry Production* **9**, 489–495.
- AOAC (2006) *Official Methods of Analysis Association*, 18th Edn. Washington, DC, USA: Association of Official Analytical Chemist.
- Armstrong TA, Flowers WL, Spears JW and Nielsent FH (2002) Long-term effects of boron supplementation on reproductive characteristics and bone mechanical properties in gilts. *Journal of Animal Science* **80**, 154–161.
- Cufadar Y, Curabay B, Gökmen SA, Bahtiyarca Y and Sevim B (2022) The effect of organic copper levels in breeding Japanese quail (*Coturnix coturnix Japonica*) diets on performance, egg quality, incubation parameters and blood parameters. *Journal of the Hellenic Veterinary Medical Society* **73**, 4619–4626.
- Deo C, Biswas A, Sharma D and Tiwari AK (2023) Effects of different concentrations of copper on performance, immunity and carcass traits in broiler Japanese quails. *Biological Trace Element Research* **201**, 4530–4537.
- dos Santos TS, Augusto KVZ, Han Y, Sartori MMP, Batistioli JS, Contin Neto AC, Ferreira Netto RG, Zanetti LH, Pasquali GAM, Muro EM, Araujo RGAC, Basso RM, Guimarães VY, Takahira RK, Kim WK and Sartori JR (2023) Effects of dietary copper and zinc hydroxychloride supplementation on bone development, skin quality and hematological parameters of broilers chickens. *Journal Animal Physiology and Animal Nutrition* **107**, 1241–1250.
- El Sabry MI, Stino FKR and El-Ghany WAA (2021) Copper: benefits and risks for poultry, livestock, and fish production. *Tropical Animal Health and Production* **53**, 487.
- El Sherif H, Fouad A, Nassar S, Wahba F, Elsabagh M and El-Iraqi K (2019) Effect of dietary copper sulphate on laying hen performance, egg quality, and oxidative stress in hot climate conditions. *European Poultry Science* **83**, 275.
- EPCEU-The European Parliament and the Council of the European Union (2010) Directive 2010/63/EU of the European Parliament and of the Council of 22 September 2010 on the Protection of Animals Used for Scientific Purposes; European Parliament: Strasbourg, France, pp. 1–47.
- Gül ET, Olgun O, Yıldız A, Tüzün AE and Sarmiento-García A (2022) Use of maca powder (*Lepidium meyenii*) as feed additive in diets of laying quails at different ages: its effect on performance, eggshell quality, serum, ileum, and bone properties. *Veterinary Sciences* **9**, 418.
- Hill GM and Shannon MC (2019) Copper and zinc nutritional issues for agricultural animal production. *Biological Trace Element Research* **188**, 148–159.
- Ibrahim IA, EL-Gendi GM, Nihad AA, Okasha HM and EL-Attrouny MM (2022) Effect of different dietary copper forms and levels on carcass characteristics and meat quality traits of broilers chickens. *Annals of Agricultural Science Moshtohor* **60**, 1091–1102.
- Kara K, Kocaoğlu Güçlü B, Şentürk M, Eren M and Baytok E (2021) Effects of catechin and copper or their combination in diet on productive performance, egg quality, egg shelf-life, plasma 8-OHdG concentrations and oxidative status in laying quail (*Coturnix coturnix japonica*). *Journal of Applied Animal Research* **49**, 97–103.
- Kaya H, Kaya A, Macit M and Kaynar Ö (2018) Effects of dietary copper supplementation on performance, egg quality parameters, yolk cholesterol and fatty acid profiles in laying hens. *Indian Journal of Animal Research* **52**, 1623–1627.
- Kilic B and Richards MP (2003) Lipid oxidation in poultry döner kebab: pro-oxidative and anti-oxidative factors. *Journal of Food Science* **68**, 686–689.
- Lin G, Guo Y, Liu B, Wang R, Su X, Yu D and He P (2020) Optimal dietary copper requirements and relative bioavailability for weanling pigs fed either copper proteinate or tribasic copper chloride. *Journal of Animal Science Biotechnology* **11**, 54.
- Mroczek-Sosnowska M, Tukaszewicz M, Wunk A and Sawosa E (2016) In ovo administration of copper nanoparticles and copper sulfate positively influences chicken performance. *Journal of the Science of Food and Agriculture* **96**, 3058–3062.
- National Research Council (1994) *Nutrient Requirement of Poultry*, 9 rev.th ed. Washington, DC: National Academy.
- Nguyen HTT, Morgan N, Roberts JR, Swick RA and Toghyani M (2020) Copper hydroxychloride is more efficacious than copper sulfate in improving broiler chicken's growth performance, both at nutritional and growth-promoting levels. *Poultry Science* **99**, 6964–6973.
- Nguyen HTT, Kheravii SK, Wu SB, Roberts JR, Swick RA and Toghyani M (2022) Sources and levels of copper affect liver copper profile, intestinal morphology and cecal microbiota population of broiler chickens fed wheat-soybean meal diets. *Scientific Reports* **12**, 2249.

- Olgun O and Aygun A** (2017) Effect of copper supplementation on performance, eggshell quality and heterophil: lymphocyte ratio in aged laying hens housed at different stocking densities. *Animal Nutrition and Feed Technology* **17**, 25–33.
- Olgun O and Yıldız AÖ** (2014) Effect of dietary supplementation of essential oils mixture on performance, eggshell quality, hatchability, and mineral excretion in quail breeders. *Environmental Science and Pollution Research* **21**, 13434–13439.
- Olgun O, Yıldız A and Şentürk ET** (2020) The effect of supplementation of organic copper to commercial quail diets on performance, egg quality and haematological parameters. *Turkish Journal of Agriculture-Food Science and Technology* **8**, 1517–1521.
- Olgun O, Gül ET, Kılınc G, Yıldız A, Çolak A and Sarmiento-García A** (2022) Performance, egg quality, and yolk antioxidant capacity of the laying quail in response to dietary choline levels. *Animals* **12**, 3361.
- Olukosi OA, Kujik SV and Han Y** (2018) Copper and zinc sources and levels of zinc inclusion influence growth performance, tissue trace mineral content, and carcass yield of broiler chickens. *Poultry Science* **97**, 3891–3898.
- Olukosi OA, Van Kuijk SJ and Han Y** (2019) Sulfate and hydroxychloride trace minerals in poultry diets—comparative effects on egg production and quality in laying hens, and growth performance and oxidative stress response in broilers. *Poultry Science* **98**, 4961–4971.
- Sacchetti G, Maietti S, Muzzoli M, Scaglianti M, Manfredini S, Radice M and Bruni R** (2005) Comparative evaluation of 11 essential oils of different origin as functional antioxidants, antiradicals and antimicrobials in foods. *Food Chemistry* **91**, 621–632.
- Sarmiento-García A, Palacios C, González-Martín I and Revilla I** (2021) Evaluation of the production performance and the meat quality of chickens reared in organic system. As affected by the inclusion of *Calliphora sp.* in the diet. *Animals* **11**, 324.
- Sarmiento-García A, Olgun O, Kılınc G, Sevim B and Gökmen SA** (2023) The use of purple carrot powder in the diet of laying quails improved some egg quality characteristics, including antioxidant capacity. *Tropical Animal Health and Production* **55**, 220.
- Scott A, Vadalasetty KP, Chwalibog A and Sawosz E** (2018) Copper nanoparticles as an alternative feed additive in poultry diet: a review. *Nanotechnology Reviews* **7**, 69–93.
- Wilson JH and Ruszler PL** (1996) Effects of dietary boron supplementation on laying hens. *British Poultry Science* **37**, 723–729.
- Wu X, Dai S, Hua J, Hu H, Wang S and Wen A** (2019) Influence of dietary copper methionine concentrations on growth performance, digestibility of nutrients, serum lipid profiles, and immune defenses in broilers. *Biological Trace Element Research* **191**, 199–206.