#### VIENNA RADIUM INSTITUTE RADIOCARBON DATES XI

#### HEINZ FELBER

Institut für Radiumforschung und Kernphysik der Österr Akademie der Wissenschaften, Vienna, Austria

Measurements have continued with the same proportional counter system, pretreatment procedure, methane preparation and measurement, and calculation, as described previously (R, 1970, v 12, p 298-318). Uncertainties quoted are single standard deviations originating from standard, sample, background counting rates, and half-life. No  $^{13}\mathrm{C}/^{12}\mathrm{C}$  ratios were measured. Sample descriptions have been prepared in cooperation with submitters.

#### **ACKNOWLEDGMENTS**

I express my thanks to Ing L Stein for excellent work in sample preparation, and to Konrad Flandorfer for careful operation of the dating equipment.

#### SAMPLE DESCRIPTIONS

I. GEOLOGIC, GLACIOLOGIC, LIMNOLOGIC, AND FOREST SAMPLES

#### Austria

### VRI-606. Scharfling, O Ö

 $2480 \pm 110$ 

Wood fragments, detritus of semi-terrestrial series, at depth 160 to 180cm in lake marl of Egelsee near Scharfling (47° 48′ N, 13° 23′ E), alt 493m, Upper Austria. Coll 1977 and subm by Roland Schmidt, Limnolog Inst, Österr Akad Wiss, Vienna.

### Gmunden, series, O Ö

Wood from boring Gschliefgraben, Gmunden, E shore of Lake Traunsee, between Traunstein and Grünberg Mts (47° 53′ N, 13° 50′ E), O Ö. Coll 1977 and subm by Jedlitschka, Forsttechn Dienst f Wildbachu Lawinenverbauung, Gmunden.

General Comment (J): dating of material that slipped down from above.

Boring 1, alt 460m:

VRI-660.	Depth 4.4m	<350
VRI-661.	Depth 29m	$2250\pm80$
VRI-662.	Depth 39m	$9500 \pm 160$
VRI-663.	Depth 53m	$10,090 \pm 150$
VRI-664.	Depth 43.5m	$9690 \pm 180$

Boring 2:

VRI-665. Depth 0.6 to 1.7m <220

House Point 1:

# VRI-675. Depth 2.5m

 $2420 \pm 80$ 

### Velden series, Kärnten

Boring core samples from lake Jeserzer See near Velden (46° 38′ 02″ N, 14° 01′ 45″ E), Carinthia. Coll 1978 and subm by Ekkehard Schultze, Limnolog Inst, Österr Akad Wiss, Wien.

General Comment (ES): absolute chronology of pollen spectrum. No NaOH pretreatment.

## VRI-639. 405 to 395cm

 $8100 \pm 140$ 

Fine detritus gyttja, depth 405 to 395cm. Comment (ES): beginning of Fagus domination; Postglacial.

## VRI-640. 485 to 475cm

 $9560 \pm 70$ 

Fine detritus gyttja, depth 485 to 475cm. Comment (ES): increasing Quercetum mixtum curve, crossing with Pinus curve; boundary Late glacial/Postglacial.

### VRI-641. 515 to 505cm

 $10,520 \pm 150$ 

Fine detritus gyttja, depth 515 to 505cm. Comment (ES): increasing NAP values; younger Late glacial.

## VRI-642. 565 to 555cm

 $13,850 \pm 310$ 

Clayey gyttja, depth 565 to 555cm. Comment (ES): stagnation in vegetational development, crossing of Pinus and Juniperus curves; older Late glacial.

## VRI-643. 585 to 575cm

 $15,200 \pm 400$ 

Clayey gyttja, depth 585 to 575cm. Comment (ES): beginning of Juniperus bush phase; older Late glacial.

## VRI-666. Pasterze, Kärnten

 $340 \pm 80$ 

Stem wood of *Pinus cembra*, in deposits of artificially dammed lake at Margaritzenboden below tongue of Pasterze glacier (47° 04′ N, 12° 46′ E), Mt Glockner Group, Carinthia. Coll 1952 by W Hosnedl and subm by Friedrich Ucik, Landesmus f Kärnten, Klagenfurt. *Comment* (FU): answers question whether stem is medieval or deposited after glacial advance ca AD 1600 (Patzelt, 1969).

## VRI-667. Linsendorf, Kärnten

 $2530 \pm 100$ 

Wood (Castanea sativa) -3.5m in basal terrace of recent R Drau, Linsendorf (46° 34′ N, 14° 28′ E), ca 2km from Anna Bridge, Carinthia. Coll 1979 and subm by Friedrich Ucik. Comment (FU): chronologically fixed point in youngest history of Drau Valley. Edible chestnut in Carinthia known for 100 yr from cultivation only.

# VRI-671. Badgastein, Salzburg

 $2670 \pm 100$ 

Cyperaceae peat, bog Zitterauer Moor, depth 85 to 95cm. Mt Stubnerkogel, 1920m asl, Badgastein (47° 06′ 10″ N, 13° 06′ 15″ E), Salzburg.

Coll 1979 and subm by Friedrich Kral, Inst Waldbau, Univ Bodenkultur, Vienna. *Comment* (FK): above this layer, palynologic evidence for beginning of human influence exists for the Gastein Valley.

### Pürgschachen series, Steiermark

Peat from different depths below soil surface of bog near Pürgschachen (47° 35′ N, 14° 20′ E), Enns Valley, Styria. Coll 1977 by Birkner, subm by Franz Wolkinger, Inst f Umweltwiss u Naturschutz, Österr Akad Wiss, Graz. No NaOH pretreatment.

VRI-610.	Depth 1m	$790 \pm 80$
VRI-611.	Depth 2m	$1790 \pm 90$
VRI-612.	Depth 3m	$1950 \pm 90$
VRI-613.	Depth 4m	$3440 \pm 100$
VRI-614.	Depth 5m	$3890 \pm 100$
VRI-615.	Depth 6m	$5760 \pm 100$
VRI-616.	Depth 7m	$7590 \pm 140$
VRI-617.	Depth 8m	$8520 \pm 140$
VRI-654. Mt	Dachstein, Stmk	$8240 \pm 260$

Pinus wood from boring core in detritus gyttja of lake Schwarzsee, Mt Dachstein E plateau (47° 30′ 30″ N, 13° 50′ E), alt 1420m, Styria. Coll 1979 and subm by Roland Schmidt. Comment (RS): dates spreading of Abies and Fagus at Mt Dachstein and palynologically recognized climatic oscillation.

#### II. ARCHAEOLOGIC AND HISTORIC SAMPLES

#### Austria

### Falkenstein series, N Ö

Charred wood with roots and inorganic detritus in hearth pit at depth 120cm in hut cut in Quads 6 to 9, excavation at Schanzboden (ring rampart) near Falkenstein (48° 43′ 22″ N, 16° 35′ 05″ E), Lower Austria. Coll 1976 and subm by Johannes-Wolfgang Neugebauer, Bundesdenkmalamt, Vienna.

General Comment (JN): repeat of VRI-577:  $5140 \pm 90$  (R, v 22, p 113) found too young by ca 600 yr.

### VRI-577a. Charcoal

 $5430 \pm 260$ 

Charcoal, 4% sample, practically free of roots and humic acids.

#### VRI-577b. Humic acids

 $5250 \pm 110$ 

Sodium humates, 26% sample, precipitated from NaOH pretreatment fraction.

# Stillfried series, N Ö

Prehistoric finds in loess, Stillfried a d March (48° 25′ 00″ N, 16° 50′ 20″ E), Lower Austria. Subm by Fritz Felgenhauer, Inst Ur- u Frühgeschichte, Univ Vienna.

General Comment (FF): absolute dating.

# VRI-635. W Rampart 1

 $2880 \pm 90$ 

Charred grass from W rampart of prehistoric bulwark. Coll 1978 by Walpurga Weiser.

## VRI-636. W Rampart 2

 $2880 \pm 90$ 

Charcoal from wooden construction of W rampart of prehistoric bulwark. Coll 1976 by Walpurga Weiser.

### VRI-637. Küssleracker

 $450 \pm 90$ 

Charcoal from cultural layer Küssleracker. Coll 1976 by Sigrid von Osten. Comment (FF): date too young.

+2000

## VRI-649. St Margarethen, Bgld

23,500

-1600

Bone find in cave fissure of quarry St Margarethen (47° 48′ N, 16° 48′ E), Burgenland. Coll by Christine Gossy, Eisenstadt, Burgenland.

## Rabensburg series, N Ö

Wood at depth -1.5m below sand layer. Lot 1762, Rabensburg (48° 41′ N, 16° 54′ E), near Mistelbach, Lower Austria. Coll 1979 and subm by Horst Adler, Bundesdenkmalamt, Vienna.

General Comment (HA): probably remnants of mill at former arm of R Thaya. Age between late high middle ages and modern times was expected.

VRI-658.  $690 \pm 90$ 

Comment (HF): de Vries correction (Suess, 1970) gives AD 1280  $\pm$  70.

VRI-659.  $520 \pm 80$ 

Comment (HF): de Vries correction (Suess, 1970) gives AD  $1400 \pm 50$ .

# VRI-673. Eggenburg, N Ö

 $2940 \pm 100$ 

Small animal bones of young Pleistocene hyena cave "Teufelslucke" near Eggenburg (48° 39′ N, 15° 49′ E), Lower Austria. Coll 1976 and subm by G Rabeder, Inst Paläont, Univ Vienna. Comment (GR): dated for comparison with other dating methods. Collagen extracted after Longin (1971).

# VRI-676. Horn, N Ö

 $23,210 \pm 510$ 

Bones in three layers below loess, -2 to -3m, at Horn-Raabs St near Horn (48° 40′ N, 15° 40′ E), Lower Austria. Coll 1931 in un-

stratified sample by Josef Bayer; subm 1980 by Wolfgang Heinrich, Inst Humanbiol, Univ Vienna. *Comment* (WH): palaeolithic hunting site with few artifacts. Collagen extracted from 1000g bone according to Longin (1971). Result is average age.

### VRI-651. Mattsee, Salzburg

 $1080 \pm 80$ 

Wood of tree-coffin -1.5m below collegiate church in Mattsee (47° 58′ 30″ N, 13° 06′ 30″ E), Salzburg. Coll 1978 and subm by Gertrud Mossler, Bundesdenkmalamt, Vienna. *Comment* (GM): coffin and remnants of oldest church was expected to be 1st millennium AD.

### VRI-674. Lamprechtshausen, Salzburg

>Modern

Bones of domestic pig. Site covered by 1 to 1.5m peat which was removed by peat-cutting, bog Waidmoos near Lamprechtshausen (48° 01' N, 12° 57' E), Salzsburg. Coll 1977 by E Stüber, Haus d Natur, Salzburg. Comment (HF): wood from same site (VRI-603, R, v 22, p 113) is  $7000 \pm 130$ . Obviously bone is atomic bomb age. Collagen extracted according to Longin (1971).

### VRI-657. Schwarzenbach, Stmk

 $3010 \pm 100$ 

Charcoal, find No. 14, -80cm in prehistoric slag deposit "Sunk Church" at Schwarzenbach (47° 29′ 30″ N, 14° 27′ 35″ E) near Trieben, Styria. Coll 1979 and subm by Clemens Eibner, Inst Ur- u Frühgesch, Univ Vienna. *Comment* (CE): ceramics point to Urnfield culture. de Vries corrected age (Suess, 1970) not unique: 1320 or 1450 BC.

#### REFERENCES

- Longin, R, 1971, New method of collagen extraction for radiocarbon dating: Nature, v 230, p 241-242.
- Patzelt, G, 1969, Zur Geschichte der Pasterzenschwankungen: Wiss Alpenvereinshefte, v 21, p 171-179.
- Suess, H E, 1970, Bristlecone pine calibration of the radiocarbon time scale 5300 BC to the present, in: Olsson, I U, Radiocarbon variatons and absolute chronology, Nobel symposium 12th, Proc: Stockholm, Almqvist & Wiksell, p 303-313.