

Testing the Wind-shock Paradigm for B-Type Star X-Ray Production with θ Car

T. F. Doyle¹, V. Petit¹, D. Cohen² and M. Leutenegger^{3,4}

¹Dept. of Physics & Space Sciences, Florida Institute of Technology
Melbourne, FL 32901

email: trisha.mizusawa@gmail.com

²Dept. of Physics & Astronomy, Swarthmore College
Swarthmore, PA 19081

³NASA/Goddard Space Flight Center
Baltimore, MD 20771

⁴University of Maryland, Baltimore County/CRESST
Baltimore, MD 21250

Abstract. We present Chandra X-ray grating spectroscopy of the B0.2V star, θ Carina. θ Car is in a critical transition region between the latest O-type and earliest B-type stars, where some stars are observed to have UV-determined wind densities much lower than theoretically expected (e.g., Marcolino *et al.* 2009). In general, X-ray emission in this low-density wind regime should be less prominent than for O-stars (e.g., Martins *et al.* 2005), but observations suggest a higher than expected X-ray emission filling factor (Lucy 2012; Huenemoerder *et al.* 2012); if a larger fraction of the wind is shock-heated, it could explain the weak UV wind signature seen in weak wind stars, but this might severely challenge predictions of radiatively-driven wind theory.

We measured the line widths of several He-, H-like and Fe ions and the f/i ratio of He-like ions in the X-ray spectrum, which improves upon the results from Nazé *et al.* (2008) (*XMM-Newton* RGS) with additional measurements (*Chandra* HETG) of MgXI and SiXIII by further constraining the X-ray emission location. The f/i ratio is modified by the proximity to the UV-emitting stellar photosphere, and is therefore a diagnostic of the radial location of the X-ray emitting plasma. The measured widths of X-ray lines are narrow, <300 km s⁻¹ and the f/i ratios place the X-rays relatively close to the surface, both implying θ Car is a weak wind star. The measured widths are also consistent with other later-type stars in the weak wind regime, β Cru (Cohen *et al.* 2008), for example, and are smaller on average than earlier weak wind stars such as μ Col (Huenemoerder *et al.* 2012). This could point to a spectral type divide, where one hypothesis, low density, works for early-B type stars and the other hypothesis, a larger fraction of shock-heated gas, explains weak winds in late-O type stars. Archival IUE data still needs to be analyzed to determine the mass loss rate and hydrodynamical simulations will be compared with observations to determine which hypothesis works for θ Car.

Keywords. line: profiles, techniques: spectroscopic, stars: winds, outflows, X-rays: stars

References

- Cohen, D. *et al.* 2008, *MNRAS*, 386, 1855
Huenemoerder, D. *et al.* 2012, *ApJL*, 756, L34
Lucy, L. B. 2012, *A&A*, 544, A120
Marcolino, W. L. F. *et al.* 2009, *A&A*, 498, 837
Martins, F. *et al.* 2005, *A&A*, 441, 735
Nazé, Y. *et al.* 2008, *A&A*, 490, 801