

# PHOTOMETRIC AND H $\alpha$ OBSERVATIONS OF LSI+61°303

J. M. PAREDES, J. MARTÍ, F. FIGUERAS, C. JORDI, G. ROSSELLÓ,  
and J. TORRA

*Dep. d'Astronomia i Meteorologia, Universitat de Barcelona, Av. Diagonal 647, E-08028  
Barcelona, Spain*

P. MARZIANI

*Dep. of Physics and Astronomy, University of Alabama, Tuscaloosa AL 35487-0324,  
U.S.A.*

J. FABREGAT, and V. REGLERO

*Dep. de Matemàtica Aplicada y Astronomia, Universidad de Valencia, 46100 Burjassot,  
Valencia, Spain*

M. J. COE, C. EVERALL, P. ROCHE, and S. J. UNGER

*Physics Department, University of Southampton, Southampton SO9 5NH, U.K.*

J. M. GRUNSFELD

*Caltech, Pasadena, CA 91125, U.S.A.*

A. J. NORTON

*Dep. of Physics, The Open University, Walton Hall, Milton Keynes MK7 6AA, U.K.*

and

R. ZAMANOV

*National Astronomical Observatory Rozhen, POB 136, 4700 Smoljan, Bulgaria*

## 1. Introduction

The Be massive X-ray binary LSI+61°303 is a 26.5 days periodic radio source (Taylor & Gregory, 1984), exhibiting radio outbursts maxima between phases 0.6-0.8. Evidence of a photometric period of similar value has also been reported (Paredes & Figueras, 1986; Mendelson & Mazeh, 1989). The previous spectroscopic radial velocity observations of Hutchings & Crampton (1981) are in agreement with the radio period, and give support to the presence of a companion. We present new optical and infrared photometric observations and high resolution H $\alpha$  spectra of LSI+61°303.

## 2. Observations and results

The photometric and spectroscopic observations are presented in Fig. 1. The JHK data show a  $\sim 0.2$  mag modulation with a deep minimum, which is reminiscent of eclipsing binaries. A detailed model involving attenuation and eclipsing of an emitting source associated to the secondary will be presented in Martí & Paredes (1993). A periodicity analysis applied to our V-




Fig. 1. *Left*) VIJHK photometric observations of LSI+61°303 folded on the 26.496 radio period. Phase zero has been set at JD 2443366.775. *Right*) Normalized  $H_{\alpha}$  line profiles.

band data indicates  $25.8 \pm 0.3$  d as most significant period. A similar analysis merging our JHK data, after subtracting their respective mean and dividing by the rms, gives  $27.0 \pm 0.3$ . Both values are close to the radio period. On the other hand, our  $H_{\alpha}$  spectroscopic observations show evidence of line parameter variability with radio phase. In particular, the FWHM of the  $H_{\alpha}$  red hump increase significantly during the phases of radio maximum, while the minimum value of the  $H_{\alpha}$  EW and the maximum value of the B/R peak ratio are observed between radio phases 0.7-0.8. Further details are reported in Paredes et al. (1993).

## References

- Hutchings J.B., Crampton D. 1981, PASP, 93, 486  
 Martí J., Paredes J.M., 1993 (in preparation)  
 Mendelson H., Mazeh T. 1989, MNRAS, 239, 733  
 Paredes J.M., Figueras, F. 1986, A&A, 154, L30  
 Paredes et al., 1993 (in preparation)  
 Taylor A.R., Gregory P.C. 1984, ApJ, 283, 273