

Diphtheria outbreak in Maranhão, Brazil: microbiological, clinical and epidemiological aspects

L. S. SANTOS¹, L. O. SANT'ANNA¹, J. N. RAMOS^{1,2}, E. M. LADEIRA²,
R. STAVRACAKIS-PEIXOTO^{1,3}, L. L. G. BORGES⁴, C. S. SANTOS¹,
F. NAPOLEÃO¹, T. C. F. CAMELLO¹, G. A. PEREIRA^{1,5}, R. HIRATA JR.¹,
V. V. VIEIRA^{2,6}, L. M. S. S. COSME⁷, P. S. SABBADINI^{1,4}
AND A. L. MATTOS-GUARALDI^{1*}

¹ *Laboratory of Diphtheria and Corynebacteria of Clinical Relevance, Faculty of Medical Sciences, Rio de Janeiro State University, UERJ, Rio de Janeiro, RJ – LDCIC/FCM/UERJ; The Collaborating Centre for Reference and Research on Diphtherial/National Health Foundation/Ministry of Health – FNS/MS, Brazil*

² *National Institute of Quality Control in Health, Oswaldo Cruz Foundation, Rio de Janeiro, RJ, Brazil*

³ *Institute of Microbiology Professor Paulo de Góes, Federal University of Rio de Janeiro - IMPPG/UFRJ, Rio de Janeiro, RJ, Brazil*

⁴ *Laboratory of Bacterial Respiratory and Systemic Diseases, Ceuma University – UNICEUMA, São Luís, MA – Brazil*

⁵ *General Coordination of Public Health Laboratories, Secretariat of Health Surveillance, Ministry of Health – CGLAB/SVS/MS, Brazil*

⁶ *Laboratory of Genetics of Microorganisms, Oswaldo Cruz Foundation, Rio de Janeiro, RJ, Brazil*

⁷ *Central Laboratory of Public Health, Oswaldo Cruz Institute – LACEN, São Luís, MA, Brazil*

Received 24 September 2013; Final revision 23 April 2014; Accepted 24 April 2014;
first published online 22 May 2014

SUMMARY

We describe microbiological, clinical and epidemiological aspects of a diphtheria outbreak that occurred in Maranhão, Brazil. The majority of the 27 confirmed cases occurred in partially ($n=16$) or completely ($n=10$) immunized children ($n=26$). Clinical signs and characteristic symptoms of diphtheria such as cervical lymphadenopathy and pseudomembrane formation were absent in 48% and 7% of the cases, respectively. Complications such as paralysis of lower limbs were observed. Three cases resulted in death, two of them in completely immunized children. Microbiological analysis identified the isolates as *Corynebacterium diphtheriae* biovar *intermedius* with a predominant PFGE type. Most of them were toxigenic and some showed a decrease in penicillin G susceptibility. In conclusion, diphtheria remains endemic in Brazil. Health professionals need to be aware of the possibility of atypical cases of *C. diphtheriae* infection, including pharyngitis without pseudomembrane formation.

Key words: Brazil, *Corynebacterium diphtheriae* biovar *intermedius*, diphtheria outbreak.

INTRODUCTION

Corynebacterium diphtheriae is the main causative agent of diphtheria, a toxemic disease whose

prevention depends on the implementation of effective immunization programmes by using diphtheria toxoid (dT) vaccines [1]. Toxigenic *C. diphtheriae*

* Author for correspondence: Prof. Dr. A. L. Mattos-Guaraldi, Laboratório de Difteria e Corinebactérias de Importância Clínica-LDCIC. Disciplina de Microbiologia e Imunologia, Faculdade de Ciências Médicas, Universidade do Estado do Rio de Janeiro, Av. 28 de Setembro, 87 – Fundos, 3º andar, Vila Isabel, RJ, Brazil, CEP 20 551-030. (Email: guaraldi@uerj.br)

may circulate in a community for 20 years after a reported case of diphtheria, even in countries where immunization programmes are followed with great efficiency [2, 3]. The introduction of toxigenic strains in a susceptible population may result in diphtheria outbreaks. All these aspects emphasize the need of vaccination strategies directed at persons of all ages and different ethnic groups and continuous surveillance of a population's immunity and new diphtheria cases [4]. Despite all relevant knowledge acquired in different areas (microbiology, pathology, internal and preventive medicine) during the years, eradication of *C. diphtheriae* has not yet been achieved [5].

Diphtheria causes significant illness and death in developing countries where vaccination coverage is low. Higher risk of acquiring the disease and potentially life-threatening complications are possible in inadequately immunized or unimmunized travellers to countries with endemic diphtheria [6, 7].

More recently in the Americas, diphtheria outbreaks have occurred in Haiti and the Dominican Republic. Diphtheria is rare in the USA; the last case occurred in an elderly traveller returning from Haiti in 2003. Although diphtheria is uncommon in industrialized countries because of longstanding routine use of vaccines containing dT [8], a changing epidemiology has been observed: *Corynebacterium ulcerans* has emerged as an important diphtheria toxin-producing pathogen and in some countries the number of diphtheria cases due to this species exceed the number reported with the classical aetiological agent, *C. diphtheriae*. Despite this, its capacity to cause disease in humans, including the inhabitants of urban centres, is still often neglected [9].

In Brazil, a developing country presenting a very large territory (8 547 403.5 km²) with varied geographical, social and economic conditions unfavourable for prevention of infectious diseases, during the last decades cases of diphtheria have been notified in many states, including in Maranhão, a northern state [6, 9–11]. Recently, one case of co-infection by *C. diphtheriae* and the infectious mononucleosis virus was also reported in a Brazilian 11-year-old child whose vaccination against diphtheria was incomplete [12]. It is important to mention that accurate data have not been available, particularly from the north, northern and central-west states, because reporting is infrequent, laboratory confirmation is not available, and the extent of carriers is not clearly known [6].

In view of these facts and considering that the vaccine does not affect the infection/colonization by

C. diphtheriae, since it is only directed against the toxin, not the whole bacterium, the endemicity of the disease in Brazil motivates diagnostic procedures appropriate to the environment, i.e. epidemiological and molecular investigations of the microorganism. The aim of the present investigation was to describe microbiological, clinical and epidemiological aspects of a diphtheria outbreak that recently occurred in Maranhão, Brazil.

METHODS

Clinical and epidemiological features

Data provided by the Public Health Secretary of the state of Maranhão showed that from January to June 2010, suspect diphtheria cases ($n=57$) and contacts of diphtheria patients ($n=117$) were notified in different villages of three municipalities of Maranhão: Jatobá ($n=95$), Colinas ($n=46$) and São Domingos ($n=33$).

Diphtheria cases were confirmed by laboratory, clinical or clinical-epidemiological criteria, according to the recommendations laid out in the Epidemiological Surveillance Guide of the Brazilian Health Ministry [13].

Origin of diphtheria bacilli strains, culture conditions and phenotypic analysis

Microorganisms were isolated from clinical samples collected with swabs from the nasopharynx and throat of the individuals at the Central Laboratory of Public Health from the state of Maranhão (LACEN-MA) by methods described previously [14]. Gram-positive *Corynebacterium*-like colonies obtained from cultures of clinical specimens collected from six patients (case nos. 10–15) were sent to the Collaborating Centre for Reference and Research on Diphtheria/National Health Foundation/Ministry of Health – FNS/MS, Brazil (LDCIC/FCM/UERJ) for further phenotypic and genotypic analysis.

Positive bacterial cultures for irregular Gram-positive rods (IGPR) were preliminarily characterized by colonial morphology, pigmentation, haemolysis, and DNase activity. Phenotypic characterization of *Corynebacterium*-like colonies was performed by conventional biochemical assays and the semi-automatized API-Coryne System v. 3.0 (bioMérieux, France) with the API web decoding system (www.apweb.biomerieux.com) [14–16]. Toxicity

was evaluated by Vero cell cytotoxicity assays as described previously [17].

C. diphtheriae biovar mitis non-toxigenic ATCC 27010 [C7s(-) tox⁻ (NCTC 11397)] type strain and the homologous toxigenic ATCC 27012 (tox⁺) strain from the American Type Culture Collection (USA), TR241 (sucrose fermenting strain) and the strain Park–Williams no. 8 (PW8), used for toxoid vaccine preparation were used as controls of experiments in addition to the *C. diphtheriae* biovar gravis VA01 strain and *C. ulcerans* 809 human isolate.

Stock cultures in 10% skim milk with 25% added glycerol were maintained at -70 °C and recovered as required by cultivation in trypticase soy broth (TSB; Difco Laboratories, USA) [18].

Antimicrobial susceptibility testing

The sensitivity to antimicrobial agents (Oxoid, UK), penicillin G (10 U), erythromycin (15 µg), clindamycin (2 µg), rifampicin (5 µg), tetracycline (30 µg), linezolid (30 µg) and vancomycin (30 µg) was determined by the disk diffusion method using inoculum equivalent to a 0.5 McFarland standard, according to Clinical Laboratory Standards Institute (CLSI) guidelines [19]. Plates were incubated at 37 °C for 24 h and reconfirmed at 48 h using a cation-adjusted Mueller–Hinton agar with 5% sheep's blood. Breakpoints for the susceptible strains were used as suggested by CLSI. The breakpoints for *Staphylococcus aureus* established by CLSI were considered in cases of penicillin.

Genotypic identification and toxigenicity evaluation using the multiplex polymerase chain reaction (mPCR) technique

A mPCR using three different primer pairs developed for detection of *C. diphtheriae* and differentiation between toxigenic and non-toxigenic strains was performed based on protocols described elsewhere [20, 21]: two primer pairs targeted to domains A and B of the *tox* gene (Dipht 2F and Dipht 4R, 719 bp) and a third primer pair targeted to a region of the *dtxR* gene (DtxR1F and DtxR1R, 258 bp) [22, 23].

DNA fingerprinting by pulsed-field gel electrophoresis (PFGE)

Genomic DNA was prepared following a method described previously [24]. The DNA was cleaved with *Sfi*I (New England BioLabs, USA) according to the

manufacturer's instructions. PFGE was performed in 0.5 × Tris-borate-EDTA 1.2% agarose gels at 13 °C with a CHEF DRII system (Bio-Rad, USA). The pulse times were 3–18 s over 20 h. A concatenated lambda DNA (New England BioLabs) was used as a molecular size marker. PFGE banding profiles were analysed using visual comparison among the strains and with the aid of automated analysis using the BioNumerics Fingerprinting software v. 4.0 (Applied Math, Belgium). PFGE types were identified by roman numerals and subtypes were identified by roman numerals followed by a letter. The similarity index of the strains was calculated using the Dice correlation coefficient with a band position tolerance of 1% and the unweighted pair-group method using average linkages (UPGMA) was used to construct a dendrogram. Strains were considered to belong to the same PFGE group if the similarity index was ≥85% band-based similarity coefficients as the cut-off values.

RESULTS

Epidemiological features

Data provided by the Public Health Secretary of the state of Maranhão showed that from January to June of 2010, suspect diphtheria cases ($n=57$) and contacts of diphtheria patients ($n=117$) were notified in different villages of three municipalities of Maranhão: Jatobá ($n=95$), Colinas ($n=46$) and São Domingos ($n=33$). A total of 27 diphtheria cases [(females ($n=18$), males ($n=9$))] was confirmed by laboratory ($n=9$), clinical ($n=7$) or clinical-epidemiological ($n=11$) criteria (Table 1). Data displayed in Figure 1 showed that the majority of the confirmed cases occurred in Jatobá ($n=20$). The highest number of diphtheria cases was observed from January to February. Most ($n=26$) of the confirmed cases occurred in children and adolescents of varied ages [<7 years ($n=13$), 7–15 years ($n=13$)] that were partially ($n=16$) or completely ($n=10$) immunized. Only one case occurred in a 31-year-old and partially immunized female patient. Three cases ended up in death: one child was partially immunized, i.e. did not receive all doses of DTP vaccine provided in the Brazilian immunization schedule [13], and the other two were completely immunized.

Clinical signs and symptoms

Clinical signs and symptoms included fever, weakness, cervical lymphadenopathy ($n=14$) and

Table 1. Clinical and epidemiological aspects of 27 confirmed cases of diphtheria notified during outbreak in Maranhão, Brazil (January–June, 2010)

Diagnostic criteria	Clinical signs and symptoms												
	Gender (Male/female)	Age group (<7, 7–20, ≥20 years)	Immunization (complete/partial)	Neck swelling	Pseudo-membrane	Tonsils	Palate	Pharynx	Larynx	Nasal cavity, lungs or skin	Paralysis of lower limbs	Serotherapy	Death
Clinical (n=7)	2/5	4/3/0	6/1	4	7	7	1	1	1	0	2	0	2
Clin/Epidemiol (n=11)	5/6	5/5/1	6/3	5	8	7	1	5	2	0	1	0	0
Laboratorial (n=9)	2/7	4/3/0	5/4	5	9	9	0	2	0	0	1	0	1

Clin/Epidemiol, Clinically compatible case epidemiologically linked to a laboratory-confirmed case; from 27 diphtheria cases, data of paralysis were not informed in nine patients: five and four cases with clinical and laboratory diagnosis, respectively. Data were provided by the Health Secretary of the state of Maranhão.

pseudomembrane formation in the tonsils (n=25), palate (n=2), pharynx (n=8), larynx (n=3) but not in the nasal cavity, lungs or skin; complications such as paralysis of lower limbs were also observed (n=4). Most of the patients exhibited pseudomembranes (92.59%). Several patients (51.85%) also exhibited enlarged lymph nodes of the neck giving a ‘bull neck’ appearance. The patients were treated with antibiotics but were not submitted to anti-diphtheria serotherapy (Table 1).

C. diphtheriae phenotypic properties

All six strains presented phenotypic characteristics of *C. diphtheriae* biovar intermedius including non-haemolytic and lipophilic colonies <1 mm. Microorganisms were positive for catalase, DNase, nitrate reductase and α-glucosidase. Pyrazinamidase, urease, gelatinase, alkaline phosphatase, esculin hydrolysis and CAMP tests gave negative results. Fermentation tests were positive for glucose, maltose, ribose and mannose; negative for glycogen, sucrose, xylose, mannitol, galactose, trehalose, arabinose and lactose. The API Coryne System confirmed the result, in which the samples showed a numerical profile 1010 324.

Genotypic identification and toxigenicity evaluation by mPCR

Results of analysis by mPCR confirmed that all six strains tested corresponded to the species *C. diphtheriae* (*dtxR* gene positive). Similar to the (tox⁺) control *C. diphtheriae* ATCC 27012 strain, five strains exhibited the presence of the *tox* gene. An MA136 strain isolated from the throat with pseudomembrane of a child living in Colinas, gave negative results by mPCR for the *tox* gene analogous to the (tox⁻) control *C. diphtheriae* ATCC 27010 strain. Complete agreement between the results of mPCR and the gold standard Vero cell cytotoxicity assays was observed for gene *tox*-positive *C. diphtheriae* strains tested.

Antimicrobial susceptibility profiles

All clinical isolates tested showed resistance to clindamycin and susceptibility to erythromycin, rifampicin, linezolid and vancomycin. Resistance to tetracycline was observed in three (MA23, MA131, MA150)




Fig. 1. Diphtheria outbreak in the state of Maranhão, Brazil. From January to June 2010. Confirmed ($n=27$) and suspect ($n=57$) cases and contacts ($n=117$) of diphtheria cases were notified in municipalities, located 400 km from the metropolitan area of São Luís (●), the capital of Maranhão State: (i) Jatobá with a surface area of 406 km² and a population of 8526 inhabitants (95 diphtheria cases); (ii) Colinas with a surface area of 2034 km² and a population of 39 167 inhabitants (46 diphtheria cases); and São Domingos with a surface area of 1303 km² and a population of 33 630 inhabitants (33 diphtheria cases). † Fatal cases ($n=3$).




Fig. 2. Pulsed-field gel electrophoresis (PFGE) types of *Corynebacterium diphtheriae* strains isolated from children with diphtheria living in the state of Maranhão, Brazil. Lane 1, λ DNA ladder PFGE marker; lanes 2–5, PFGE type Ia (MA19, MA23, MA52, MA131 strains, respectively); lane 6, PFGE type II (MA136 strain); lane 7, PFGE type Ib (MA150 strain); lane 8, profile III (TR241 strain); lane 9, profile IV (VA01 strain); lane 10, profile V (PW8 strain).

strains. Two strains (MA23 and MA52) showed intermediate susceptibility to penicillin G.

PFGE analysis

Three distinct PFGE types (Ia, Ib, II) were found in the *C. diphtheriae* isolates from Maranhão (Fig. 2). The PFGE subtypes Ia and Ib showed a similarity coefficient $\geq 95\%$ and were considered genetically

related. PFGE type Ia was the most frequently observed in the five strains evaluated in this study. Only the non-toxicogenic MA136/13 strain exhibited PFGE type II. Three other PFGE types (III, IV, V) different from those presented by the strains from Maranhão were exhibited by the PW8 strain, which is the only major strain used in toxoid vaccine production and by TR241 and VA01 strains isolated from patients with diphtheria in Rio de Janeiro city [10].

DISCUSSION

It is well known that the effectiveness of the vaccination using the dT against diphtheria caused by *C. diphtheriae* varies from 45% to 90% [25]. In the recent diphtheria outbreak in Maranhão the majority of Brazilian patients were children and many of them had been covered by a complete course of immunization. Cervical lymphadenopathy (70·37%) and neck oedema (51·85%) were observed in our study, indicating the occurrence of a severe form of the disease in Maranhão. The presence of pseudomembrane, a pathognomonic sign of the disease [1], was found in 92·59% of the patients despite immunization; mostly children (aged 1–12 years) and females; 11·12% cases were fatal, possibly due to the lack of diphtheria antitoxin and delayed treatment, as previously observed at the beginning of epidemic diphtheria in the states of the Former Union of Soviet Socialist Republics [4]. In Dhule, India, there was a shift in the median age of disease to adolescents (10–15 years) without gender differences; only 18% reported disease despite immunization. About 64% of the confirmed cases presented with a well-defined pseudomembrane [25]. In the newly independent states of the Former Soviet Union, a high proportion of cases of pharyngeal or tonsillar diphtheria without pseudomembrane formation in adults were observed [26].

A case of diphtheria with pseudomembrane formation in a Brazilian 32-year-old woman has previously been reported. Her history included complete paediatric immunization (DTP) and three doses of adult formulation tetanus and dT 2 years earlier. Clinical diagnosis of diphtheria was not made until microbiological examination of specimens confirmed *C. diphtheriae* biovar *gravis* (VA01 strain) [10]. Multilocus sequence typing (MLST) to study genetic relationships in *C. diphtheriae* strains isolated in the urban area of Rio de Janeiro showed that the VA01 strain was assigned to an already known sequence type (ST), indicating that it was part of a clonal complex that comprises strains isolated in Canada (ST80) [27]. These cases reinforced the potential susceptibility of Brazilian children and adults to diphtheria that may be caused by both endemic and imported clones.

Although in Brazil and Eastern Europe outbreaks mainly occur due to the dissemination of *C. diphtheriae* biovars *mitis* and *gravis*, respectively [4, 5], in Maranhão the diphtheria outbreak was caused by *C. diphtheriae* biovar *intermedius*. Interestingly,

diphtheria cases and deaths also caused by *C. diphtheriae* biovar *intermedius* have been documented in previously immunized individuals in India [25, 28]. By using PFGE it was possible to conclude that five toxigenic strains isolated in the state of Maranhão in 2010 were classified within the same clone or PFGE group. However, MA136 strain, the only one without the *tox* gene, was classified in a different PFGE type unrelated to toxigenic strains from Maranhão, indicating the circulation of more than one PFGE type of *C. diphtheriae* biovar *intermedius* during the outbreak which had taken place in this northern state of Brazil.

The presence of diphtheria bacilli resistant to drugs, frequently used in the treatment of infections from different sources (e.g. penicillin and erythromycin) has been noticed in some countries, including Brazil [21]. Penicillin tolerance has been hypothesized to be a cause of treatment failure of *C. diphtheriae* infections [29, 30]. Presently, even considering the breakpoints for *S. aureus*, we observed a decreased susceptibility to penicillin in *C. diphtheriae* strains isolated from the patients of municipalities of the state of Maranhão. Microorganisms also showed resistance to tetracycline, as previously observed in Rio de Janeiro [21]. *C. diphtheriae* strains of PFGE types Ia and Ib and II showed 100% susceptibility to erythromycin, linezolid and vancomycin. The representative strain of PFGE type II (MA136 strain) showed susceptibility to most of the antimicrobial agents tested, except clindamycin. Continuous surveys of antibiotic susceptibility of *C. diphtheriae*, especially in developing countries where diphtheria is endemic and invasive infections may occur remain necessary.

In conclusion, diphtheria remains endemic in Brazil, including the northern state of Maranhão. Unvaccinated or inadequately vaccinated people who travel to countries where diphtheria is endemic may be at increased risk. Our review of diphtheria patients hospitalized in Maranhão suggests that the clinical features of the disease in partially vaccinated patients may still be similar to those that were observed in the pre-vaccine era. However, we cannot exclude the possibility of some cases of diphtheria being misdiagnosed or not notified to the public health authorities due to the lack of knowledge by physicians regarding the clinical and laboratory diagnosis of diphtheria, particularly when patients are not fully protected against diphtheria toxin. The scarcity of epidemiological data concerning diphtheria in developing countries may be due to the fact that doctors

and microbiologists are not aware of the possibility of atypical cases of *C. diphtheriae* infection, including pharyngitis without pseudomembrane [25] as well as invasive infections such as pneumonia [31, 32], arthritis and endocarditis [21, 33–36] and catheter-related infections [16].

ACKNOWLEDGEMENTS

This work was supported by FAPEMA, FAPERJ, CNPq, CAPES, SR-2/UERJ, Programa de Núcleo de Excelência (PRONEX) and Programa Nacional de Pós-Doutorado – PNPd (CAPES/MEC).

DECLARATION OF INTEREST

None

REFERENCES

1. **Hadfield TL, et al.** The pathology of diphtheria. *Journal of Infectious Disease* 2000; **181** (Suppl. 1): S116–120.
2. **Golaz A, et al.** Epidemiology of diphtheria in South Dakota. *South Dakota Journal of Medicine* 2000; **53**: 281–285.
3. **Wagner KS, et al.** Screening for *Corynebacterium diphtheriae* and *Corynebacterium ulcerans* in patients with upper respiratory tract infections 2007–2008: a multicentre European study. *European Society of Clinical Microbiology and Infectious Disease* 2011; **17**: 519–525.
4. **Dittman S, et al.** Successful control of epidemic diphtheria in the State of former Union of Soviet Socialist Republics: lessons learned. *Journal of Infectious Disease* 2000; **181** (Suppl. 1): S10–20.
5. **Mattos Guaraldi AL, Hirata Jr. R, Damasco PV.** Diphtheria in Brazil and around the world: aspects of the current scenario. *Revista Imunizações* 2011; Suppl. 1: S2–20.
6. **Mattos-Guaraldi AL, et al.** Diphtheria remains a threat to health in the developing world – an overview. *Memórias do Instituto Oswaldo Cruz* 2003; **98**: 987–993.
7. **Bitragunta S, et al.** Persistence of diphtheria, Hyderabad, India, 2003–2006. *Emerging Infectious Disease* 2008; **14**: 1144–1146.
8. **Center for Disease Control and Prevention (CDC).** The Yellow Book: CDC Health Information for International Travel. *Diphtheria* 2012; **1**: 162–165.
9. **Dias AA, et al.** *Corynebacterium ulcerans* diphtheria: an emerging zoonosis in Brazil and worldwide. *Revista de Saúde Pública* 2011; **45**: 1176–1191.
10. **Mattos-Guaraldi AL, et al.** Diphtheria in a vaccinated adult in Rio de Janeiro, Brazil. *Brazilian Journal of Microbiology* 2001; **32**: 236–239.
11. **Information System for Notifiable Diseases (SINAN) of the Brazilian Ministry of Health** (http://portal.saude.gov.br/portal/arquivos/pdf/casos_difteria.pdf). Accessed 10 April 2010.
12. **Mattos-Guaraldi AL, et al.** Concurrent diphtheria and infectious mononucleosis: difficulties for management, investigation and control of diphtheria in developing countries. *Journal of Medical Microbiology* 2011; **60**: 1685–1688.
13. **Brazilian Secretariat of Health Surveillance.** Guide to Epidemiological Surveillance – 7th edn. Brasília, Brazil: Brazilian Ministry of Health, 2009; **3**: 21–43. (Series A. Standards and Technical Manuals).
14. **Funke G, Bernard KA.** Coryneform gram-positive rods. In: Murray PR, Baron EJ, Jorgensen JH, Landry MA, Pfaller MA, eds. *Manual of Clinical Microbiology*. Washington (DC): American Society for Microbiology Press, 2007, pp. 485–514.
15. **Pimenta FP, et al.** DNase test as a novel approach for the routine screening of *Corynebacterium diphtheriae*. *Letters in Applied Microbiology* 2008; **46**: 307–311.
16. **Gomes DL, et al.** *Corynebacterium diphtheriae* as an emerging pathogen in nephrostomy catheter-related infection: evaluation of traits associated with bacterial virulence. *Journal of Medical Microbiology* 2009; **58**: 1419–1422.
17. **Efstratiou A, et al.** Comparison of phenotypic and genotypic methods for detection of diphtheria toxin among isolates of pathogenic corynebacteria. *Journal of Clinical Microbiology* 1998; **36**: 3173–3177.
18. **Hirata R Jr., et al.** Intracellular viability of toxigenic *Corynebacterium diphtheriae* strains in HEp-2 cells. *FEMS Microbiology Letters* 2002; **215**: 115–119.
19. **CLSI.** Performance standards for antimicrobial susceptibility testing; twentieth informational supplement. CLSI document M100-S20. Wayne, PA: Clinical and Laboratory Standards Institute, 2010.
20. **Pimenta FP, et al.** A multiplex PCR assay for simultaneous detection of *Corynebacterium diphtheriae* and differentiation between non-toxigenic and toxigenic isolates. *Journal of Medical Microbiology* 2008; **57**: 1438–1439.
21. **Pimenta FP, et al.** A PCR for dtxR gene: application to diagnosis of non-toxigenic and toxigenic *Corynebacterium diphtheriae*. *Molecular and Cellular Probes* 2008; **22**: 189–192.
22. **Pallen M J, et al.** Polymerase chain reaction for screening clinical isolates of corynebacteria for the production of diphtheria toxin. *Journal of Clinical Pathology* 1994; **47**: 353–356.
23. **Nakao H, et al.** Heterogeneity of diphtheria toxin gene, *tox*, and its regulatory element, *dtxR*, in *Corynebacterium diphtheriae* strains causing epidemic diphtheria in Russia and the Ukraine. *Journal of Clinical Microbiology* 1996; **34**: 1711–1716.
24. **Baio PV, et al.** Clonal multidrug-resistant *Corynebacterium striatum* within a nosocomial environment, Rio de Janeiro, Brazil. *Memórias do Instituto Oswaldo Cruz* 2013; **108**: 23–29.
25. **Phalkey RK, et al.** Preventing the preventable through effective surveillance: the case of diphtheria in a rural

- district of Maharashtra, India. *BMC Public Health* 2013; **13**: 317.
26. **Kadirova R, Kartoglu Ü, Strebel PM.** Clinical characteristics and management of 676 hospitalized diphtheria cases, Kyrgyz Republic, 1995. *Journal of Infectious Disease* 2000; **181** (Suppl. 1): S110–115.
 27. **Viguetti SZ, et al.** Multilocus sequence types of invasive *Corynebacterium diphtheriae* isolated in the Rio de Janeiro urban area, Brazil. *Epidemiology and Infection* 2012; **140**: 617–620.
 28. **Dravid MN, Joshi SA.** Resurgence of diphtheria in Malegaon and Dhule regions of north Maharashtra. *Indian Journal of Medical Research* 2008; **127**: 616–617.
 29. **Von Hunolstein C, et al.** Penicillin tolerance amongst non-toxigenic *Corynebacterium diphtheriae* isolated from cases of pharyngitis. *Journal of Antimicrobial Chemotherapy* 2002; **50**: 125–128.
 30. **Gladin DP, et al.** Sensitivity of *Corynebacterium diphtheriae* isolated in Saint Petersburg to antibacterial drugs. *Antibiotics and Chemoterapy* 1999; **44**: 17–21.
 31. **Honma Y, et al.** A case of afebrile pneumonia caused by non-toxigenic *Corynebacterium diphtheriae*. *Japanese Journal of Infectious Disease* 2009; **62**: 327–329.
 32. **Trost E, et al.** Pangenomic study of *Corynebacterium diphtheriae* that provides insights into the genomic diversity of pathogenic isolates from cases of classical diphtheria, endocarditis, and pneumonia. *Journal of Bacteriology* 2012; **194**: 3199–3215.
 33. **Guran P, et al.** Septic arthritis due to a nontoxigenic diphtheria bacillus. *Archives Française de Pédiatrie* 1979; **36**: 926–929.
 34. **Lin RVTP, et al.** *Corynebacterium diphtheriae* endocarditis in an adult with congenital heart disease: a case report. *Journal of Tropical Medicine and Hygiene* 1994; **97**: 189–191.
 35. **Mattos-Guaraldi AL, Formiga LCD.** Bacteriological properties of a sucrose fermenting *Corynebacterium diphtheriae* strain isolated from a case of endocarditis. *Current Microbiology* 1998; **37**: 156–158.
 36. **Hirata R, et al.** Potential pathogenic role of aggregative adhering *Corynebacterium diphtheriae* of different clonal groups in endocarditis. *Brazilian Journal of Medical Biological Research* 2008; **41**: 986–991.