

ARTICLE

Psychogenic amnesia: when memory complaints are medically unexplained

Gavin C. M. McKay & Michael D. Kopelman

Gavin C. M. McKay was a Specialist Registrar in Neuropsychiatry at the Memory Disorders Clinic at St Thomas' Hospital. He has obtained his certificate of completion of training and is now a locum consultant in general adult psychiatry, maintaining a special interest in neuropsychiatry. **Michael D. Kopelman** is Professor of Neuropsychiatry at the Institute of Psychiatry, King's College London, and Consultant Neuropsychiatrist with the South London and Maudsley NHS Foundation Trust, based at the Neuropsychiatry and Memory Disorders Clinic at St Thomas' Hospital, London. Professor Kopelman has published widely on many aspects of memory disorders (including the amnesic syndrome, Alzheimer's and semantic dementia, confabulation, psychogenic amnesia, amnesia for offences, and false confessions), as well as calculation disorders, sleep disorder, post-traumatic stress disorder and neuroimaging. **Correspondence** Dr Gavin McKay, Mascalls Park, Mascalls Lane, Brentwood, Essex CM14 5HQ, UK. Email: Gavin@doctors.org.uk

SUMMARY

The focus of this article is the assessment and management of medically unexplained ('psychogenic') amnesia, which we classify here as global or situation specific. Other psychiatric causes for memory disorder and neurological conditions that could cause diagnostic confusion are briefly reviewed, as are forensic aspects of memory complaints. Finally, brain and physiological mechanisms potentially associated with psychogenic amnesia are discussed.

DECLARATION OF INTEREST

None.

Amnesia (Fig. 1) has been defined as 'an abnormal mental state in which memory and learning are affected out of all proportion to other cognitive functions in an otherwise alert and responsive patient' (Victor 1971). Memory impairment can affect the learning of new material (anterograde amnesia), owing to impairments in the encoding, storage or retrieval stages. It can also affect the recall of previously acquired memories (retrograde amnesia), which might involve personal experiences (episodic memory), general information (semantic memory) or perceptuomotor skills (procedural memory). Psychological factors can cause both anterograde and retrograde memory loss (or a combination of the two).

Types of psychogenic amnesia

Psychologically based amnesia includes the persistent anterograde memory impairment present in mental disorders such as depression, which in an extreme form can lead to depressive pseudodementia. Alternatively, it can cause transient or discrete episodes of retrograde and/or anterograde memory

loss, which we refer to here as 'psychogenic' amnesia (i.e. loss not due to identifiable brain disease).

Why 'psychogenic'?

A number of terms have been used to describe medically unexplained amnesia, including 'hysterical', 'psychogenic', 'dissociative' and 'functional'. Each requires the exclusion of an underlying neurological cause and the identification of a precipitating stress that has resulted in amnesia. Unfortunately, the presence of amnesia may make it difficult to identify the stress until either informants have come forward or the amnesia itself has resolved.

Both DSM-IV (American Psychiatric Association, 2000) and ICD-10 (World Health Organization, 1992) favour the term 'dissociative' amnesia. Some have argued strongly for 'functional' amnesia as a description more acceptable to patients (Stone 2005). Others prefer 'medically unexplained amnesia'. We favour 'psychogenic' amnesia, because it points to underlying psychological processes without assuming that any particular psychological mechanism is involved (a difficulty with 'dissociative' amnesia). Also, it does not specify whether the memory loss is produced (partly or entirely) consciously ('factitious' or 'exaggerated' amnesia) or purely unconsciously ('hysterical' amnesia). The term 'functional' amnesia has the problem that the amnesia could in many respects be considered dysfunctional. However, regardless of the term used, the question remains, at what point, in cases of doubt, should the clinician assume that a psychological stressor or a marker of neuropathology or brain pathology is the primary cause of the symptoms? A subsidiary to this is, does the amnesia have both psychological and neurological contributions to its aetiology?




FIG 1 A partial classification of amnesia.

Types of global psychogenic amnesia

Psychogenic fugue

This is a syndrome of sudden onset, involving loss of all autobiographical memories, including personal identity. It is usually associated with a period of wandering, for which there is a gap in continuous memories on recovery. Fugue states usually last a matter of hours or days. There is anecdotal evidence that they are more common in wartime, particularly in soldiers about to return to the front. Psychogenic fugue is classified in DSM-IV as dissociative fugue.

Case 1 Psychogenic fugue

A 40-year-old woman 'came round' on the London Underground. She had approached a staff member, telling them that she had no recollection of who she was or what she was doing. The only clues to her identity were a bag with a few clothes and a letter that turned out to be addressed to someone else. Staff contacted the police, who took her to an accident and emergency department. After being medically cleared she was admitted to a psychiatric unit. Initially, she appeared depressed but this resolved with no medication. In an effort to trace her identity a picture was shown on national television, with no responses. An amyltal interview was tried, again with no success. She was eventually discharged from hospital and the retrograde memory loss persisted. One year later, relatives in the USA sent a missing person poster to the UK police and she was traced. It emerged that, following a marital crisis in the context of an unstable marriage, she had disappeared from her home and taken a flight to the UK. She had a history of depression.

After obtaining a clear history from the family a second amyltal interview was attempted and she did recover most of her retrograde memory function (Kopelman 1994).

Three main factors may predispose to psychogenic fugue:

- precipitating stress such as emotional, relationship, marital or financial problems
- depressed mood, sometimes with suicidal thoughts
- a history of a transient organic amnesia: some studies report that more than half of affected individuals have experienced severe head injury (Berrington 1956).

Psychogenic focal retrograde amnesia

This is a persistent state in which the individual loses all retrograde memories. The amnesia is described as 'focal' because anterograde memory is relatively or completely spared. There is considerable controversy concerning whether a brain lesion in isolation can ever cause this dissociation between anterograde and retrograde amnesia (Kopelman, 2002a). Often, spouses or close family members are not initially recognised and later the patient claims to have relearned who they are. Clinically, there may be two aetiological routes to the appearance

of this syndrome. The first is following a typical fugue with wandering and loss of personal identity, when disproportionate retrograde amnesia persists. The other is after a minor brain insult such as mild concussion in a predisposed person. These cases are often difficult to manage, both because of the coexistence of minor organic, psychological factors and the possibility of financial compensation for injuries, and because the underlying stressors are often not obvious on initial inquiry. In both types of aetiology abnormal illness behaviour may arise and be reinforced by the gains derived from the emotional and practical support of family, friends, helping professions and the welfare system.

Case 2 Psychogenic focal retrograde amnesia

'I put things in boxes, I choose to put them in the back of my mind. I've always done that'. These are the words, during treatment, of a 62-year-old man who collapsed at work with a transient right-sided weakness and a complete loss of autobiographical memory. On admission, he was disoriented in time and place as well as person, and there was a mild loss of power in the right arm and leg with an equivocally up-going right plantar response. A computed tomography (CT) scan was normal but a magnetic resonance imaging (MRI) brain scan showed evidence of a few pinpoint regions of altered signal bilaterally, consistent with a history of previously diagnosed hyperlipidaemia and diabetes. However, the physicians attending this man felt confident that his memory loss was entirely disproportionate to his neurological signs, which rapidly resolved. The patient did not recognise his wife, and he could not remember the names and ages of his children. He claimed to have relearned this (personal semantic) knowledge about his life and that 'each day I remember more of the day before'. On formal tests, he showed severe and extensive autobiographical memory loss, with intact anterograde memory. When first seen, he and his family were angry at any suggestion that there might be a psychological component to his memory loss. However, during the succeeding weeks, his wife provided information about his difficult childhood and subsequent emotional problems. The initial onset had occurred after the patient had been confronted about 'moonlighting' in two employments and his dismissal from both. After being seen on a regular basis for several weeks, he was more willing to accept a psychological contribution to his amnesia and, following an amyltal interview, virtually all of his memories were recovered.

(This patient is now out of contact. A previous brief report was published (Kopelman 2000) with his permission. Identifying details have been changed.)

Multiple personality disorder (DSM-IV dissociative identity disorder)

A key symptom in this disorder is some degree of between-personality amnesia. How much access each personality has to the others' memories is variable (for a review see Kihlstrom 2000). A recent study (Huntjens 2006) showed evidence that items claimed not to have been remembered in one personality did influence the choice made later in a recognition task

BOX 1 Differential diagnosis of psychogenic fugue

- Alcoholic ‘blackout’
- Transient global amnesia
- Transient epileptic amnesia
- Epileptic fugue (rare)
- Post-traumatic amnesia following head injury
- Simulation/malingering

after a personality switch. In contrast to ‘simulators’, who tended to choose implausible alternative answers, the choice made was often a ‘near miss’ or plausible alternative to the correct answer. This led the study’s authors to conclude that there is no evidence of a memory retrieval problem but instead that individuals with dissociative identity disorder hold incorrect beliefs about their memory functioning. How much this disorder is the result of iatrogenic influences on patients already vulnerable to psychogenic memory disorders is still a matter of debate (Merskey, 1995).

Differentiation of transient psychogenic and organic amnesia

The differential diagnosis of psychogenic fugue is described in Box 1 (Kopelman 2002b). The features common to, and differentiating between, transient organic and psychogenic amnesias are shown in Table 1. The retrograde memory impairment in a transient neurological amnesia such as transient global amnesia, or in a persistent one such as dementia or Korsakoff syndrome, usually follows Ribot’s law: recently learned material is more affected than earlier memories. This leads to a characteristic ‘temporal gradient’ when retrograde memory test performance is plotted against time. The memory impairment in psychogenic amnesia, in contrast, characteristically shows a reversed gradient, with earlier memories preferentially affected (see Fig. 2).




FIG 2 Example of retrograde memory performance plotted against age of memory for a psychogenic amnesia (—■—) and a neurological amnesia such as transient global amnesia (---◆---). Note the characteristic ‘reversed’ temporal gradient for psychogenic amnesia (see Kritchevsky 1997).

Transient global amnesia

Transient global amnesia is a syndrome of sudden-onset anterograde and variable retrograde amnesia, with repetitive questioning not associated with other neurological deficits. It usually lasts for a matter of minutes to hours, with complete recovery.

Its aetiology is poorly understood. One theory is that an upsurge in venous pressure, which could explain the association with emotional or physical stress, causes hypoxo-ischaemia in the hippocampus, an area of key importance for memory. There is an association with migraine in 25% of cases (Hodges 1990). Hyperintensities in the hippocampus have been demonstrated on diffusion weighted imaging (Sander 2005). Hodges & Warlow (1990) found an epileptic basis for a small proportion of cases (7% of 153) and the term ‘transient epileptic amnesia’ was coined (Kapur 1990). Here the attacks tend to be briefer and more recurrent than in transient global amnesia, they are associated with other seizure types and they usually respond well to anticonvulsants.

TABLE 1 Differentiating transient organic and psychogenic amnesias

Characteristic	Transient organic amnesia	Transient psychogenic amnesia
Is preceded by precipitating stress/significant life-event	Possible	Usually
Normal results in standard investigations (routine EEG, CT, MRI)	Possible	Usually
Loss of personal identity	Never in transient global amnesia	Always in fugue
Repetitive questioning	Common in transient global amnesia/transient epileptic amnesia	Seldom in fugue/psychogenic amnesia where there may be <i>la belle indifference</i>
Other symptoms/signs	Sensorimotor symptoms in transient epileptic amnesia	Wandering in fugue
Time-memory performance curve	‘Temporal gradient’ of retrograde amnesia in transient global amnesia/transient epileptic amnesia	‘Reversed gradient’ in psychogenic amnesia
Anterograde memory	Impaired	Often spared

Assessment of psychogenic amnesia

Box 2 shows the salient features to be elicited in the history, mental state and physical examinations, and the investigations to be performed when assessing a likely psychogenic amnesia. Neuropsychological testing should be carried out whenever possible.

Management of global psychogenic amnesia

The patient and family should be engaged sympathetically, avoiding confrontation. The patient often has coexisting depression and this should be treated. The disadvantages of the amnesia should be gently drawn to attention and the advantages minimised, and the patient should be encouraged to resume normal activities, including employment.

Interview under sedation

It has long been noted, anecdotally, that an interview under intravenous sedation (abreaction) can help recover memories in psychogenic amnesia (Herman, 1938). This is a controversial technique and its use in the UK is in decline (Patrick 1990). There are concerns about the suggestibility of individuals during sedation and the reliability of memories obtained (Rogers 1988). Adding to the controversy, the technique has been used to ‘recover’ memories of childhood sexual abuse, leading several international bodies to caution against its use in this particular situation (Brandon 1997). Although problems remain, we would distinguish this use (hunting for specific, long distant abuse memories) from the use of amytal in the treatment of acute-onset non-neurological loss of recent (and remote) memories.

There are no randomised placebo-controlled trials of interview under sedation in psychogenic amnesia (Kavirajan, 1999). Lambert & Rees (1940) compared barbiturates, hypnosis and supportive treatment in a group of 247 servicemen with hysteria, 56 of whom had amnesia. The groups were non-randomly chosen and the barbiturate group contained patients who had not recovered spontaneously or by interview without sedation. The authors noted that, although there was no difference in remission rates between the groups (82%, 75% and 88% respectively) in a *post hoc* analysis, interview under sedation was found to be ‘by far the quickest and the easiest method’ of recovering memory. Despite the absence of high-quality evidence demonstrating its efficacy and the controversy regarding its use, interview under sedation can still have a place in the management of psychogenic amnesia if it is used cautiously and without any attempt at making suggestions.

Conducting an amytal interview

As much collateral history from the family and other informants should be elicited as possible before

BOX 2 The assessment of likely global psychogenic amnesia

History

- Precipitant of amnesia
Head injury? Stress?
- What memories are affected?
Retrograde only, or anterograde also?
Complete, or ‘islets’ of preserved memory?
Is it confined to autobiographical incidents?
Is knowledge of facts about oneself or facts in general (semantic memory) affected?
Personal identity?
Close family, spouse’s identity?
Repetitive questioning?
- How does it feel to the patient? Indifference?
An insistence that ‘recovered’ memories have been relearned?
- Psychiatric history
Somatisation, depression, other isolated and ‘unexplained’ physical symptoms?
- Medical history
Previous transient neurological amnesia?
Seizures?
Alcohol/substance intake?

Mental state examination

- *La belle indifférence*?
- Other hysterical symptoms or symptoms of somatising?
- Depression?

Physical examination

- Particular emphasis on the cardiovascular and neurological systems

Neuropsychological tests

- Anterograde memory may be relatively normal, with disproportionate retrograde amnesia
- Tests of retrograde memory (e.g. the Autobiographical Memory Interview) show a ‘reversed gradient’
- Tests of simulation can be performed: malingers will perform worse than by chance alone

Investigations

- Complete dementia blood screen
- MRI
- EEG if relevant

Essential collateral history

- GP and hospital records
- Witness/family accounts
- Undisclosed stressors from family

proceeding and the interview must not be rushed into. It is best to arrange an in-patient admission for the interview, so that any psychological issues arising can be discussed with staff. Clinical impression suggests that sodium amytal is the most effective drug for this purpose; we have found that lorazepam has too slow an onset and midazolam is more difficult to titrate. Amytal can be more accurately titrated to gain greater control of the degree of sedation. It must be noted that amytal is unlicensed for this purpose and it is advisable to obtain written informed consent before the procedure. A protocol is available (Perry 1982). The risk of respiratory arrest and the lack of a pharmacological antagonist are considerations and a crash trolley and oxygen should be readily available. Significant cardiopulmonary disease would be a contraindication. For guidelines on the safe use of ‘conscious sedation’ see American Society of Anesthesiologists Task Force (2002). Complications are rare; at a time when such interviews were more commonplace, a study reported 1 respiratory arrest in 500 amytal interviews, and this because the amytal was administered too quickly (Hart 1945).

The principle of interview under sedation is to reach a level of intoxication in which the patient is deeply relaxed but not asleep. This can be judged by getting the patient to count backwards from 50 until numbers are mixed up or skipped. At this point conversation about neutral topics can turn towards the amnesic gap, using clues gained from the history. Questions are as open as possible and leading questions should not be used (Rogers 1988). This level of sedation is maintained by slowly injecting more amytal until no further progress is made or the maximum dose reached. Information revealed is talked over again while the patient is recovering, to ensure that the memories are retained in awareness.

Prognosis

If the amnesia is long-standing and the family is enmeshed in a system that maintains the symptoms then it is difficult to change. The longer the amnesia has persisted, the less likely is complete recovery.

Memory complaints or a psychogenic amnesia may form part of a wider picture of somatisation disorder. If this is the case, efforts should be made to limit the number of investigations and doctors involved, and some form of regular psychological treatment should be instigated.

Situation-specific amnesia

Amnesia for offences

Offenders as well as victims of crimes commonly claim amnesia regarding the offence and psychiatrists can be called upon to comment on this. Cross-sectional studies have found that in 25–45% of homicides, 8% of other violent crimes and a small percentage of non-violent crimes, offenders claim amnesia for the offence (Kopelman, 2002a).

Differentiation from underlying neurological conditions such as an epileptic automatism, post-ictal confusional state, head injury, hypoglycaemia, sleepwalking or rapid eye movement (REM) sleep disorder is important (for references see Kopelman 2002a). Amnesia for an offence can also occur in alcohol intoxication, substance misuse and acute psychosis but purely psychological amnesia occurs most commonly in crimes of passion. In one series, 40% of offenders claiming amnesia on arrest for murder still had amnesia for the offence 3 years after conviction. They were less likely to deny the offence than a similarly convicted comparison group, thus providing some evidence that malingering was not the sole cause of their memory loss (Pyszora 2003).

Post-traumatic stress disorder

The relationship between trauma and memory is complex, with some memories apparently enhanced and others forgotten. This forgetting is sometimes

interpreted in terms of ‘peri-traumatic dissociation’, which is one of the diagnostic criteria for acute stress disorder in DSM–IV. However, it may also be the case that, when something extraordinary happens, we ask ourselves to recall far more detail than we would normally expect, and experience the shortfall as ‘gaps’ in memory (Kopelman 2002a). The presence of head injury can complicate the clinical picture. With regard to the memory disturbance seen, amnesia predominates in head injury, and intrusive memories predominate in post-traumatic stress disorder (PTSD), where memory lapses are less common. The two could be seen to lie at the opposite extremes of a continuum.

One particular theoretical model proposes a dual-component memory system to explain memory of personally experienced traumatic events (Brewin 1996). On the one hand are verbally accessible memories (autobiographical memories) that have received some conscious processing and have been transferred to the long-term store. These memories can be retrieved either automatically or deliberately. On the other are the situational accessible memories responsible for trauma-related flashbacks and nightmares. These memories have undergone little conscious processing and involve the autonomic and motor responses at the time of the trauma. Because these memories are not encoded verbally they are difficult to communicate to others and to control, often being triggered by trauma-related cues.

Anterograde memory dysfunction has been demonstrated in people with PTSD and there are claims that they have a loss of hippocampal volume on MRI (Bremner 1999), which has been attributed to effects of glucocorticoids (Markowitsch 1996).

Childhood sexual abuse

Amnesia for childhood sexual abuse has been an issue of debate and can have important medico-legal implications. Doubts over the veracity of memories of such abuse that come to light during psychotherapy are expressed in the proposition of ‘false memory syndrome’.[†] However, even studies by protagonists of false memory syndrome (Loftus 1994) show that some forgetting of abuse does occur.

As with PTSD, mechanisms have been proposed whereby such forgetting and subsequent (possibly erroneous) retrieval might occur (Schacter 1996), particularly in relation to certain cues or triggers (Andrews 2000).

Other common psychiatric disorders causing memory complaints

Cognitive symptoms are increasingly seen as part of the schizophrenic syndrome. A meta-analysis concluded that significant deficits were found in

[†]For a discussion of false memory syndrome, see Merskey H (1998) Prevention and management of false memory syndrome. *Advances in Psychiatric Treatment*; 4: 253–60. Ed.

verbal recall (Aleman 1999). Performance on delayed recall tests by people with schizophrenia was one whole standard deviation below that of people without. Both *post-mortem* and structural MRI studies have shown people with schizophrenia to have smaller medial temporal lobes compared with controls (Shenton 2001).

Depression is associated with both subjective complaints of forgetting and measurable deficits on anterograde memory tests (Dalglish 2002). In its extreme form, the picture may be of a pseudodementia, especially in the elderly. These deficits are at least partly corrected by treatment. Again, it is interesting that one meta-analysis concluded that there is evidence of reduced hippocampal volume in depression (Campbell 2004). Bipolar disorder, especially in older patients with multiple episodes, has been reported to be associated with measurable impairments on neuropsychological tests of memory, even when patients are in remission (Savitz 2005).

Another way of looking at these data is to suggest that psychopathology has a non-specific effect on memory. What is not clear is how much the cognitive deficits are directly related to the pathophysiology of each condition, how much is an epiphenomenon of the psychological condition, and whether and to what extent they are secondary to confounders such as medication or substance misuse. Well-designed prospective studies would clarify these issues.

What are the brain mechanisms of psychogenic amnesia?

There is evidence from 'lesion' studies that memories of emotional or traumatic events are processed differently from 'ordinary memories'. In particular, emotional memories, especially those involving fear, may use amygdaloid circuits, whereas 'normal' learning involves other brain regions (Fine 2000). The amygdala is a brain area richly connected to the emotional circuitry of the brain and the autonomic nervous system.

Lesions in the medial temporal and the diencephalic areas of the brain cause an amnesic syndrome with anterograde amnesia and also a variable degree of retrograde amnesia. Frontal areas are believed to be more involved in the effortful retrieval of autobiographical memories (Kopelman 2002a). A few years ago, one of us (M.D.K) proposed a model of how brain systems and psychosocial factors interact to produce a global psychogenic amnesia (as in fugue or focal retrograde amnesia) (Kopelman, 2002a). It hypothesises that stress affects a frontal/executive retrieval system, especially if there has been a past learning experience of an organic amnesia or the person is severely depressed. Stress sometimes has a direct effect on the medial

temporal–diencephalic system, but if this remains unaffected new learning can continue, so that only the retrieval of past autobiographical information is impaired. Severe stress can also affect retrieval from the 'personal semantic belief system', resulting in the transient loss of personal identity. In situation-specific amnesia, it has been suggested that some aspects of forgetting are due to a narrowing of consciousness, with attention focused on central perceptual details.

Functional neuroimaging is offering insights into the brain mechanisms involved in psychogenic amnesia. Several functional MRI (fMRI) studies have been carried out in psychogenic amnesia. Although there do appear to be abnormalities in areas important for memory, the location and even the direction of the activation or deactivation has differed from study to study, making interpretation difficult (references can be supplied on request). Anderson and colleagues (2004) reported evidence that executive mechanisms can be recruited to prevent unwanted memories from entering awareness, and that repeated use of this strategy inhibits the subsequent recall of the suppressed memories. This research group went on to use event-related fMRI to investigate the neural substrates of these phenomena. They found that during the suppression of unwanted memories, poorer recall correlated with activation of the lateral prefrontal cortex and deactivation of the medial temporal lobe. Further research using event-related fMRI during autobiographical memory retrieval paradigms in patients with psychogenic amnesia will clarify the brain mechanisms associated with memory suppression.

Conclusions

We hope that this article has given the clinician knowledge of the range of presentations of psychogenic amnesia and a practical framework on which to base an assessment. This should allow differentiation between neurologically based amnesias and comorbid psychiatric conditions. Management may involve an amytal interview, but this should not be rushed into: first, a full understanding of the patient's underlying issues should be sought and any depression treated.

Psychogenic amnesia may result from the effects of stress on the frontal memory retrieval systems and inhibitory mechanisms, rather than from an effect on the medial temporal lobes or diencephalic structures, which are the usual sites of pathology in neurologically based amnesias. Functional neuroimaging is now allowing the investigation of the interaction of these structures during normal retrieval and suppression in memory tasks, and may eventually provide insights into the brain mechanisms underlying psychogenic amnesia.

MCQ answers

1	2	3	4	5
a f	a f	a f	a f	a f
b f	b f	b t	b t	b f
c t	c f	c f	c f	c f
d f	d f	d f	d f	d t
e f	e t	e f	e f	e f

References

Aleman A, Hijman R, de Haan EHF, et al (1999) Memory impairment in schizophrenia. A meta-analysis. *American Journal of Psychiatry*; **156**: 1358–66.

American Psychiatric Association (2000) *Diagnostic and Statistical Manual of Mental Disorders (4th edn) (DSM-IV)*. APA.

American Society of Anesthesiologists Task Force on Sedation and Analgesia by Non-Anesthesiologists (2002) Practice guidelines for sedation and analgesia by non-anesthesiologists. *Anesthesiology*; **96**: 1004–17.

Anderson MC, Ochsner KN, Kuhl B, et al (2004) Neural systems underlying the suppression of unwanted memories. *Science*; **303**: 232–5.

Andrews B, Brewin CR, Ochera J, et al (2000) The timing, triggers and qualities of recovered memories in therapy. *British Journal of Clinical Psychology*; **39**: 11–26.

Berrington WP, Liddell DW, Foulds GA (1956) A re-evaluation of the fugue. *Journal of Mental Science*; **102**: 280–6.

Brandon S, Boakes J, Glaser D, et al (1997) Reported recovered memories of child sexual abuse. Recommendations for good practice and implications for training, continuing professional development and research. *Psychiatric Bulletin*; **21**: 663–5.

Bremner JD, Randall P, Scott TM, et al (1995) Memory-based measurement of hippocampal volume in patients with combat-related posttraumatic stress disorder. *American Journal of Psychiatry*; **152**: 973–81.

Brewin CR, Dalgleish T, Joseph S (1996) A dual representation of posttraumatic stress disorder. *Psychological Review*; **103**: 670–86.

Campbell S, Marriott M, Nahmias C, et al (2004) Lower hippocampal volume in patients suffering from depression: a meta-analysis. *American Journal of Psychiatry*; **161**: 598–607.

Dalgleish T, Cox SG (2002) Memory and emotional disorder. In *The Handbook of Memory Disorders* (eds MD Kopelman, AD Baddeley, BA Wilson): 437–49. John Wiley & Sons.

Fine C, Blair RJ (2000) The cognitive and emotional effects of amygdala damage. *Neurocase*; **6**: 435–50.

Hart WL, Ebaugh F, Morgan DC (1945) The amytal interview. *American Journal of Medical Science*; **210**: 125–31.

Herman M (1938) The use of intravenous sodium amytal in psychogenic amnesic states. *Psychiatric Quarterly*; **12**: 738–42.

Hodges JR, Warlow CP (1990) Syndromes of transient amnesia: towards a classification. A study of 153 cases. *Journal of Neurology, Neurosurgery and Psychiatry*; **53**: 834–43.

Huntjens RJ, Peters ML, Woertman L, et al (2006) Inter-identity amnesia in dissociative identity disorder: a simulated memory impairment? *Psychological Medicine*; **36**: 857–63.

Kapur N (1990) Transient epileptic amnesia: a clinically distinct form of neurological memory disorder. In *Transient Global Amnesia and Related Disorders* (ed HJ Markowitsch): 140–51. Hogrefe & Huber.

Kavirajan H (1999) The amobarbital interview revisited: a review of the literature since 1966. *Harvard Review of Psychiatry*; **7**: 153–65.

Kihlstrom JF, Schacter, DL (2000) Functional amnesia. In *Handbook of Neuropsychology* (ed LS Cermak): 409–27. Elsevier Science.

Kopelman MD, Christensen H, Parfitt A, et al (1994) The Great Escape. A neuropsychological study of psychogenic amnesia. *Neuropsychologia*; **32**: 675–91.

Kopelman MD (2000) Focal retrograde amnesia and the attribution of causality: An exceptionally critical review. *Cognitive Neuropsychology*; **17**: 585–621.

Kopelman MD (2002a) Disorders of memory. *Brain*; **125**: 2152–90.

Kopelman MD (2002b) Assessment of psychogenic amnesia. In *Handbook of Memory Disorders* (2nd edn) (eds AD Baddeley, MD Kopelman, BA Wilson): 427–48. John Wiley & Sons.

Kritchevsky M, Zouzounis J, Squire L (1997) Transient global amnesia and functional retrograde amnesia. Contrasting examples of episodic memory loss. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*; **352**: 1747–54.

Lambert C, Rees WL (1940) Intravenous barbiturate in the treatment of hysteria. *BMJ*; **2**: 70–3.

Loftus EF, Polonsky S, Fullilove MT (1994) Memories of childhood sexual abuse: remembering and repressing. *Psychology of Women*; **18**: 67–84.

Markowitsch HJ (1996) Organic and psychogenic retrograde amnesia. Two sides of the same coin? *Neurocase*; **2**: 357–71.

Merskey H (1995) Multiple personality disorder and false memory syndrome. *British Journal of Psychiatry*; **166**: 281–3.

Patrick M, Howells R (1990) Barbiturate-assisted interviews in modern clinical practice. *Psychological Medicine*; **20**: 763–5.

Perry JC, Jacobs D (1982) Overview: clinical applications of the amytal interview in psychiatric emergency settings. *American Journal of Psychiatry*; **139**: 552–9.

Pyszora NM, Barker AF, Kopelman MD (2003) Amnesia for criminal offences: a study of life sentence prisoners. *Journal of Forensic Psychiatry*; **14**: 475–90.

Rogers R, Wettstein RM (1988) Drug-assisted interviews to detect malingering and deception. In *Clinical Assessment of Malingering and Deception* (ed R Rogers): 197–204. Guilford Press.

Sander K, Sander D (2005) New insights into transient global amnesia: recent imaging and clinical findings. *Lancet Neurology*; **4**: 437–44.

Savitz J, Solms M, Ramesar R (2005) Neuropsychological dysfunction in bipolar affective disorder: a critical opinion. *Bipolar Disorder*; **7**: 216–35.

Schacter DL (1996) *Searching for Memory: The Brain, the Mind, and the Past*. Basic Books.

Shenton ME, Dickey CC, Frumin M, et al (2001) A review of MRI findings in schizophrenia. *Schizophrenia Research*; **49**: 1–52.

Stone J, Carson A, Sharpe M (2005) Functional symptoms in neurology: management. *Journal of Neurology, Neurosurgery and Psychiatry*; **76**: 13–21.

Victor M, Adams RD, Collins GH (1971) *The Wernicke-Korsakoff syndrome*. F.A. Davis.

World Health Organization (1992) *The ICD-10 Classification of Mental and Behavioural Disorders: Clinical Descriptions and Diagnostic Guidelines*. WHO.

MCQs

1 Psychogenic fugue:

- a usually lasts a few months
- b is increasing in incidence
- c is commonly preceded by head injury
- d is associated with repetitive questioning
- e is rarely associated with a loss of personal identity.

2 Psychogenic focal retrograde amnesia:

- a is never associated with head injury
- b has a classic temporal gradient
- c is associated with severe anterograde amnesia
- d often causes repetitive questioning
- e persists longer than fugue in the majority of cases.

3 The management of psychogenic amnesia:

- a should always involve an amytal interview
- b should involve the family
- c should involve an element of confrontation
- d will be made easier if the patient is encouraged to apply for social support benefits
- e is more likely to result in a good outcome the longer the amnesia persists.

4 Transient global amnesia:

- a shows a pattern of retrograde memory loss very similar to that of psychogenic amnesia
- b is often associated with stressful antecedents
- c is characteristically associated with abnormal EEG

- d is associated with a loss of personal identity
- e usually responds to anticonvulsants.

5 Amnesia for an offence:

- a is commonly reported in shoplifting
- b is caused by malingering in almost all cases
- c is rare in homicide
- d can result from an automatism
- e persists in only a tiny minority once the trial is over.