

alcohol, thiamine shortage, and alcohol-related cerebrovascular illness. Moreover, it is worth noting that alcohol use disorder (AUD) has the potential to worsen the advancement of neurodegenerative processes in individuals already diagnosed with dementia.

Conclusions: The association between AUD (Alcohol Use Disorder) and dementia is complex and involves multiple factors, presenting considerable difficulties in terms of clinical intervention and treatment. The use of early intervention strategies and public health initiatives focused on addressing alcohol use disorder (AUD) could have a significant impact on preventing or reducing the development of dementia.

Disclosure of Interest: None Declared

EPV0761

Under-attribution in self-agency on pre-reflexive task connected to positive schizotypal traits among healthy students

I. Szendi^{1,2*}, N. Domján³, H. Pásztor³, T. Jenei², O. Bóna², C. Kovács², A. Pejin³, P. Pajkossy⁴, Á. Szöllösi⁴ and M. Racsmány⁵

¹Department of Personality, Clinical and Health Psychology, University of Szeged, Szeged; ²Department of Psychiatry, Kiskunhalas Semmelweis Hospital, Kiskunhalas; ³Department of Psychiatry, University of Szeged, Szeged; ⁴Department of Cognitive Science, Budapest University of Technology and Economics, Budapest and ⁵Department of Cognitive and Neuropsychology, University of Szeged, Szeged, Hungary

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.1394

Introduction: The aim of this study was to identify low-risk traits of schizophrenia among healthy undergraduate student volunteers, and the investigation of these traits with regards to their specificity in contrast to individuals with a latent disposition towards bipolar disorder. Self-agency, as a phenomenon closely related to psychomotor functioning, provides a unique opportunity for the investigation of subjective self-perception.

Objectives: The implicit self-agency performances that are considered illness- (or risk state-) specific were compared between groups to find early markers of a specific schizotypic developmental path.

Methods: In a sample of 710 healthy university students, with the help of screening questionnaires, we were able to successfully form two risk groups, in one of them the emphasis on cyclothymia (CTF: Cyclothymia factor group, N=25), and in the other (PSF: Positive schizotypy factor group, N=26) the tendency to unusual experiences and paranoid thinking emphasis was typical. We assigned a properly matched control group (N=29) displaying both features on average. We focused on the implicit aspect of self-agency, using the well-known paradigm of intentional binding, as well as the self-developed device that exclusively tests the pre-reflexive feeling of movement initiation, the sense of self-agency.

Results: During the examination of intentional binding, although the specific predictive and retrospective component indicators did not show any significant difference for either group, the association of the sound alone could induce a binding effect in the control group. In the predictable frequency condition, there was a strong significant effect ($W = 65.00$, $p = .007$, $r_{rb} = -.60$), and in the non-predictable condition a trend-level effect. Remarkably, this binding

effect did not develop in either the CTF or PSF groups, indicating an implicit agency impairment in both risk groups. However, during the examination of sense of self-agency, we observed a disturbance specifically among healthy college students with positive schizotypal traits, in the form of falsely attributing their movement initiation to external influences. The percentage of this 'miss'-type answering differed between groups, $H(2) = 7.68$, $p = .021$, $\epsilon^2 = .10$. The Dwass-Steel-Critchlow-Fligner pairwise comparisons showed that this difference was due to the PSF Group showing a significant difference from the Control Group ($W = -3.83$, $p = .019$), but not from the CTF group, and the CTF Group also did not differ from the Control Group.

Conclusions: Thus, in premorbid conditions, in at-risk groups of non-help-seeking individuals, or in cases of early detection of prodromal abnormalities, objective confirmation of suspected susceptibility to schizophrenia may be aided by, among other things, instrumental assessment of self-agency.

Disclosure of Interest: None Declared

EPV0762

Chronotype and health related quality of life among undergraduate university students

E. L. Nikolaev¹, I. Poverinov² and M. Alhasan^{3*}

¹Department of Social and Clinical Psychology; ²Philosophy, Sociology and Pedagogy Department and ³Medical Faculty, Ulianov Chuvash State University, Cheboksary, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.1395

Introduction: Chronotype represents genetically determined behavioral characteristics of a person's twenty-four-hour activity. Research shows that a person's chronotype is interrelated with their mental health. Are there similar connections with general health and health related quality of life?

Objectives: To establish how various chronotypes are represented in university students and if there are any interrelations between chronotypes and health-related quality of life

Methods: We used SF-12 Health Survey и Morningness–Eveningness Questionnaire (MEQ) by Horne and Ostberg to survey 305 university students of both genders.

Results: The results showed that the majority of the students (71.2%) have an intermediate chronotype. The second goes a moderate morning chronotype (17.7%), the third – a moderate evening chronotype (9.8%). Definite morning and definite evening chronotypes were revealed in less than 1% of the students. SF-12 Health Survey scale indicators that assess quality of life corresponded to standard scores for the given group of the respondents. We have revealed valid relations in correlational interconnections of the achieved parameters. Thus, the morning chronotype is most consistently associated ($p < 0.01$) in undergraduate university students with higher indicators of health related quality of life including General Health ($r = .23$), Vitality ($r = .21$), Role Physical ($r = .18$), Role Emotional ($r = .17$), Physical Functioning ($r = .16$), Social Functioning ($r = .13$). Mental Health and Bodily Pain in university students are not connected with the chronotype ($p > 0.05$).

Conclusions: Therefore, this research establishes that the majority of the students are related to the intermediate chronotype, and the