

WSRT 21-cm continuum field in Cygnus OB2: in search of more Wolf-Rayet stars*

Diah Y.A. Setia Gunawan^{1,2}, Karel A. van der Hucht²,
A. Ger de Bruyn^{1,3}, and Peredur M. Williams⁴

¹*Kapteyn Astronomical Institute, RU Groningen, the Netherlands*

²*Space Research Organization Netherlands, Utrecht, the Netherlands*

³*Netherlands Foundation for Research in Astronomy,
Dwingeloo, the Netherlands*

⁴*Royal Observatory, Edinburgh, United Kingdom*

Abstract. We present a deep look into the Cygnus OB2 region using the WSRT. A $2^\circ \times 2^\circ$ map of the optically highly-obscured region was obtained at 1400 MHz, with a noise level down to 0.2 mJy and an angular resolution of $1''$. We will compare the resulting radio point source list with optical and near-IR catalogues, in order to identify stellar wind sources like WR and OB stars.

1. Introduction

The direction towards the Cygnus OB2 association is known to have strong and varying extinction ($E_{B-V} \simeq 1.4\text{--}2.3$). WR 146 (WC6) has a visual extinction of $A_v=9.5$, while WR 147 (WN8) has $A_v=12.6$. Placing them at the 1.7 kpc distance of the Cyg OB2 association would make these WR stars too luminous. Similarly, placed at the distance of Cyg OB2, Cyg OB2 #12 would have $M_v = -10$ and $M_{bol} = -11$ which would make it one of the most luminous objects in the Galaxy. However, this star may be a foreground object like WR 146 and WR 147: recently accepted distances for WR 146 and WR 147 (1.2 kpc and 0.63 kpc, respectively) bring their X-ray, optical and IR luminosities closer to those of other WR stars of their respective subclasses. Recent studies also show both WR 146 and WR 147 to be WR+OB binary systems with colliding stellar winds, generating excess non-thermal radio fluxes (Dougherty *et al.* 1996; Williams *et al.* 1997; Niemela *et al.* 1998), and making them bright radio stars.

In our 21-cm WSRT observations of the WR 146 field, we note a number of additional point sources which are not present in, *e.g.*, DSS and IRAS maps of the same area. Comparison with our 6-cm maps indicates inverted spectral indices, suggesting a galactic (thermal) origin of those possibly stellar wind sources like OB and WR stars.

Encouraged by those examples, we have started a project to search for more of such point sources in the line-of-sight to the Cyg OB2 association, which is veiled by the thick IS clouds. At radio wavelengths one can see through the huge

*Based on observations made with the Westerbork Synthesis Radio Telescope. The WSRT is operated by the Netherlands Foundation for Research in Astronomy (NFRA) which is financially supported by the Netherlands Organization for Scientific Research (NWO).

and varying extinction, and expose sources which are more difficult to discover at optical and infrared wavelengths. Therefore, we have mapped the Cyg OB2 region at 21-cm continuum with the *WSRT*.

2. Observations and data reduction

Our mosaic observation, a $2^\circ \times 2^\circ$ field centered at $\alpha(1950) = 20^h 31^m 17^s.30$ and $\delta(1950) = 40^\circ 39' 0''$, was divided into 6×6 pointings with a grid of about 0.4 degrees. The *WSRT* East-West-array of 14 antennas, with a maximum baseline of 2.8 km, observed each grid point cyclically with an integration time of 50 seconds. The total observing time was 6×12 hours spanning two observing periods: 2×12^h in 1996 and 4×12^h in 1997. The continuum back-end used had a frequency bandwidth of 1360-1420 MHz.

The reduction was carried out using the *WSRT-NEWSTAR* reduction package. The data were first calibrated for antenna corrections using standard calibrators observed directly before and after each 12^h run. Thereafter, they were SELF-CAL-ed. The resulting maps, with a $13'' \times 19''$ beam, were subsequently CLEAN-ed and the restored maps were combined into one mosaic. The noise level across the map is 0.2–0.6 mJy per beam, allowing us to detect 2 mJy sources at a 5σ level.

3. Preliminary result and future work

We have made use of the SIMBAD data base to identify the point sources in the field. Several point sources remain unidentified. Comparison with the *WSRT* 92-cm survey including the same field (Vasisht & de Bruyn, unpublished) and 6-cm observations would yield the spectral indices α ($S_\nu \propto \nu^\alpha$) of these sources. WR stars are expected to have thermal spectral indices, $\alpha \simeq 0.6$ – 0.8 . At the distance of Cyg OB2, new WR stars would be rather weak sources, but less weak if they are foreground objects. In addition, if they are non-thermal colliding wind sources like WR 146 ($\alpha_{6-21} \simeq -0.7$) and WR 147 ($\alpha_{6-21} \simeq -1$), they could be relatively bright radio stars.

Follow-up deep optical and infrared photometry and spectroscopy has to determine interstellar extinction and distance towards the individual point sources and thus to assess their classification and luminosity properly.

Acknowledgments. DYASG thanks the Leids Kerkhoven-Bosscha Fonds for a grant allowing her to attend the symposium. The support from the Computer Group at the Kapteyn Laboratory is gratefully acknowledged.

References

- Dougherty, S.M., Williams, P.M., van der Hucht, K.A., Bode, M.F., Davis, R.J. 1996, *MNRAS* 280, 963
- Niemela, V.S., Shara, M.M., Wallace, D.J., Zurek, D.R., Moffat, A.F.J. 1998, *AJ* 115, 2047
- Williams, P.M., Dougherty, S.M., Davis, R.J., van der Hucht, K.A., Bode, M.F., Setia Gunawan, D.Y.A. 1997, *MNRAS* 289, 10