December 1966

Vol. LXXX No. 12

The Journal of Laryngology and Otology

EDITED BY G. H. BATEMAN

ASSISTANT EDITOR LIONEL TAYLOR

Contents

THE TREATMENT OF POST-CRICOID CARCINOMA BY SINGLE STAGE PHARYNGO-LARYNGECTOMY AND COLON TRANSPLANT

VOICE PRODUCTION FOLLOWING LARYNGO-PHARYNGO-OESOPHAGECTOMY WITH COLON TRANSPLANT

THE USE OF IMPULSIVE MECHANICAL STIMULI OF VARIABLE DURATION IN THE EVALUATION OF THE INDICATION TIME OF THE OTOLITH ORGANS

THE USE OF ELECTRONYSTAGMOGRAPHY IN CLINICAL PRACTICE

GLOTTIC CANCER-THE IMPORTANCE OF A BALANCED VIEW OF TREATMENT

ON SUBARACHNOID-TYMPANIC FISTULAE. A REPORT OF Two Cases of the Rare Indirect Type •

Consanguinity among Parents of Deaf Children in the Jewish Population in Israel

C. REMINGTON HOBBS and KENNETH MULLARD

> MOTI LALL and GORDON EVISON

> > E. G. WALSH

A. G. D. MARAN

H. J. SHAW

R. J. BENNETT

SRAEL M. FEINMESSER, L. BAUBERGER-TELL and F. MARKUS

CLINICAL RECORDS-

ANGIOMA OF MAXILLARY SINUS · · · L. H. HIRANANDANI, D. K. GOSAVI, R. D. MELGIRI and G. K. HIRANANDANI PERIPHERAL GANGLIONEUROMA OF NECK · PREM. K. KAKAR, K. L. SAWHNEY and P. S. SAHARIA A SIMPLE AID FOR AURAL MICROSURGERY · H. J. SHAW FIBRO-ANGIOMA OF THE FACIAL NERVE · ABDEL SALAM EL BARBARY

Societies' Proceedings

BOOK REVIEW

GENERAL NOTES

London Headley Brothers

109 Kingsway WC2

Annual Subscription for Spublished on line by Cambridge University Press

Monthly, 12/6 net post free

ADVERTISEMENTS

The Journal of Laryngology and Otology

(FOUNDED in 1887 by MORELL MACKENZIE and NORRIS WOLFENDEN)

EDITED BY

G. H. BATEMAN

ASSISTANT EDITOR LIONEL TAYLOR

1. Original articles which have not been published elsewhere are invited and should be sent to the Editor. They are considered for publication on the understanding that they are contributed to this *Journal* solely. Reproduction elsewhere, in whole or in part, is not permitted without the previous written consent of the Author and Editor and the customary acknowledgment must be made.

2. Manuscripts should be typewritten, on one side only of the paper, and well spaced. Captions to illustrations should be typed on a separate sheet and sent at the same time as original photographs, etc.

The Harvard system of recording references should be used, e.g. GREEN, C., and BROWN, D. (1951) J. Laryng., 65, 33. Abbreviations of Journals should follow the style recommended in *World Medical Periodicals*, published by World Health Organization, 1952.

Health Organization, 1952. It is most important that authors should verify *personally* the accuracy of every reference before submitting a paper for publication.

3. Galley proofs and engraver's proofs of illustrations are sent to the author. Corrections, which should be kept to a minimum, must be clearly marked, and no extra matter added. Proofs should be returned within ς days.

4. Illustration blocks will normally be held by the Printers for three years, after which they will be destroyed. Any author is entitled to have these returned to him, but a request for this must be sent within three years of the appearance of the article, to HEADLEY BROTHERS, 109 Kingsway, London, WC2. Coloured illustrations will be charged to authors, unless a special grant is authorized by the Editor.

5. Orders for reprints must be sent when returning galley proofs, and for this purpose special forms are supplied.

6. Authors of original communications on Oto-Laryngology in other journals are invited to send a copy, or two reprints, to the *Journal of Laryngology*. If they are willing, at the same time, to submit their own abstract (in English, French, Italian, or German) it will be welcomed.

7. Editorial communications may be addressed to THE EDITOR, Journal of Laryngology, c/o HEADLEY BROTHERS, 109 Kingsway, London, WC2.

8. The annual subscription is five guineas sterling (U.S.A. \$15) post free, and payable in advance.

9. Single copies will be on sale at 125. 6d. each; copies of parts up to Vol. LXIII which are available may be purchased at 75. 6d. each.

10. All subscriptions, advertising and business communications should be sent to the publishers, HEADLEY BROTHERS, 109 KINGSWAY, LONDON, WC2.

United States of America

Orders for this Journal may be sent through local booksellers, or to STECHERT-HAFNER, INC., 31-33 East 10th Street, New York, or direct to the publishers, HEADLEY BROTHERS, 109 KINGSWAY, LONDON, WC2, England.

© Journal of Laryngology and Otology, 1966

CONTENTS

PAGE

THE TREATMENT OF POST-CRICOID CARCINOMA BY SINGLE STAGE PHARYNGO-LARYNGECTOMY AND COLON TRANSPLANT. C. Remington Hobbs (Winchester) and Kenneth Mullard (South- ampton)	1193
Voice Production following Laryngo-Pharyngo-Oesophagectomy with Colon Transplant. Moti Lall and Gordon Evison (Bristol)	
The Use of Impulsive Mechanical Stimuli of Variable Duration in the Evaluation of the Indication Time of the Otolith Organs. E. G. Walsh (Edinburgh)	1218
THE USE OF ELECTRONYSTAGMOGRAPHY IN CLINICAL PRACTICE. A. G. D. Maran (Edinburgh)	1224
GLOTTIC CANCER—THE IMPORTANCE OF A BALANCED VIEW OF TREATMENT. H. J. Shaw (London)	
On Subarachnoid-Tympanic Fistulae. A Report of Two Cases of the Rare Indirect Type. R. J. Bennett (Newcastle, Staffs.)	1242
CONSANGUINITY AMONG PARENTS OF DEAF CHILDREN IN THE JEWISH POPULATION IN ISRAEL. M. Feinmesser, L. Bauberger-Tell and F. Markus (Jerusalem)	1253
Clinical Records	
ANGIOMA OF MAXILLARY SINUS. L. H. Hiranandani, D. K. Gosavi, R. D. Melgiri and G. K. Hiranandani (Bombay)	
PERIPHERAL GANGLIONEUROMA OF NECK. Prem. K. Kakar, K. L. Sawhney and P. S. Saharia (New Delhi, India)	1260
A SIMPLE AID FOR AURAL MICROSURGERY. H. J. Shaw (London)	
FIBRO-ANGIOMA OF THE FACIAL NERVE. Abdel Salam El Barbary (Cairo, U.A.R.)	1265
Societies' Proceedings	1268
Book Review · · · · · · · · · · · ·	1271
General Notes · · · · · · · · · · · · · · · · · · ·	1272

BÉKÉSY Accessory

For the very first time a fully automatic unit has been made available for a Clinic Audiometer, making the combined units into a true BÉKÉSY Audiometer.

This automatic accessory is being made for the PETERS SPD/5 CLINIC AUDIOMETER, which has a continuous frequency range and continuously variable attenuation. The result is therefore a true Békésy audiometer.

It can be attached to or detached from the audiometer in under one minute so that the two units together offer the facilities of an advanced clinic audiometer or Békésy audiometry at will.

It can be fitted to any existing SPD/5 Audiometer.

Please write for full details and demonstration to:

ALFRED PETERS & SONS LIMITED 51 GELL STREET, SHEFFIELD 3



'Otosporin' rapidly destroys virtually all bacteria commonly found in ear infections, reduces irritation, inflammation and pain. 'Otosporin' brand Drops contain polymyxin B sulphate, neomycin sulphate and hydrocortisone in a bland suspension designed to facilitate penetration. Issued in bottles of 5 ml.





Burroughs Wellcome & Co. London (The Wellcome Foundation Ltd.)



the antistaphylococcal antibiotic that penetrates tissues at these levels...

bears the FUCIDIN label



The clinical significance of these levels has been confirmed in many publications during the past five years. Over 100 authors have endorsed the value of Fucidin in the treatment of staphylococcal Hayes Middlesex

infection. Fucidin (Sodium Fusidate B.P.) 250 mgm capsules.

Leo Laboratories Limited



Please mention The Journal of Laryngology and Otology when replying to advertisements

iv

WHEN A HEARING AID IS RECOMMENDED ONLY AMPLIVOX OFFERS THIS HEARING AID SERVICE

The design and manufacture of a complete range with individual fitting facilities.
Best of other makes in addition to Amplivox aids—Amplivox hearing aids are ONLY available from Amplivox.
Highest degree of training of any hearing aid organisation. Master Hearing Aid evaluations.
Expert follow-up and rehabilitation advice.
15 branch offices with while-you-wait service.
FOR YOUR PATIENTS' GREATEST SATISFACTION SPECIFY AMPLIVOX



AMPLIVOX HEARING ADVISORY SERVICE 80 NEW BOND STREET LONDON, W.1



Specifically designed for you and your patients. Precision-engineered, functionally-built equipment for the Ear, Nose and Throat and other Specialists.

S.M.R. keeps you and your patients' comfort and satisfaction in mind. Whatever your needs . . . whether they be for one unit or a completely equipped treatment room, you are assured of the finest . . . a room that reflects professional good taste.



Picture shows what can be done with small treatment room areas. The rooms are complete from the compact treatment unit to the physician's stool. Yet, this equipment fits easily and functionally into a small area. This is just one of many room arrangements available from S.M.R.



Please send for our new catalog Surgical Mechanical Research,

INC.

883 West 16th Street, P.O. Box 1185 Newport Beach, Calif. 92663

Direct factory sales, F.O.B. Newport Beach, Calif.

ONLY **AMPLIVOX** provides a <u>complete</u> audiometric service!



Transistor Audiometers

with air conduction or air and bone conduction with masking, for hospitals, schools, consulting rooms, industrial medical departments, screening programmes, etc.



Clinical Audiometers

with narrow band masking for the highest accuracy and extended resolution.


Audiometer Booths

for ideal testing conditions in hospitals, clinics, factories. Standard and special sizes available.



PLUS An efficient and speedy calibration service please write for technical Ilterature to meet your needs 80 New Bond Street, London, W.1. Tel: HYDe Park 9888

viii



It is only natural to manage without an aid or with an old aid for far longer than one should.

New aids give facilities undreamt of even five years ago. Today a tiny Multitone behind the ear aid gives nearly everything body-aids gave five years ago. Even good aids, like people, get older. Where should one go?... To Multitone.


They lead technically, having pioneered such developments as telesonic coils and automatic volume control. Even more important, Multitone the leading British manufacturers, are scrupulously correct. In the recent criticism of hearing aid firms expressed in the House of Commons, Multitone were exempted by name, as a firm above reproach. Every Multitone Centre will give fair *free advice* without obligation.



ELECTRIC COMPANY LIMITED

Underwood St., London N.1

CLE 8022



--that is the triumph of the **MACKIE** hearing aid

The new Mackie post-aural hearing aid is made in several different circuit versions so that the gain and frequency response can be selected to match the patient's needs; and it is available where required with a unique form of automatic volume control which has proved highly successful in cases of recruitment. The forward pick-up microphone gives maximum intelligibility of speech without background interference.

It is exported to many countries including the United States of America–a tribute to its high quality and advanced design.

In Great Britain it is available for trial by your patients from *independent* hearing aid dealers in all parts of the country.

MANUFACTURED BY A. G. R. MACKIE LTD. A member of the Bonochord Group of Companies FOR FURTHER DETAILS WRITE TO 48 WELBECK STREET, LONDON WI

THE LARYNGOSCOPE

A Monthly Journal devoted to the disease of EAR, NOSE AND THROAT


Official organ for the American Laryngological Rhinological and Otological Society


Price \$18.00 per year Foreign \$19.00 per year

Established 1896

THEODORE E. WALSH, M.D. EDITOR

517 SOUTH EUCLID AVENUE SAINT LOUIS MO. 63110.





Please mention The Journal of Laryngology and Otology when replying to advertisements

ORBENIN and its success in Gram-Positive Infections

Otitis media Otitis externa

Boils Carbuncles Infected dermatoses

Pneumonia Lung abscess Acute bronchitis

Osteomyelitis Osteitis

Tonsillitis Pharyngitis

Post-operative wound infections Burns Skin graft protection

Septicaemia Acute endocarditis

Staphylococcal enterocolitis

Staphylococcal urinary tract infections

Staphylococcal meningitis

ORBENIN (cloxacillin sodium B.P.) is a penicillin active against virtually all Gram-positive organisms, including staphylococci, β -haemolytic streptococci and pneumococci, irrespective of their resistance to other antibiotics.

It is encouraging that the widespread use of Orbenin in hospitals has not resulted in the emergence of resistant strains.

More than 300 references in the World Literature testify to the excellent clinical results obtained with Orbenin in a wide variety of Gram-positive infections.

Detailed literature is available on request.

Dosage:

Two capsules (500mg.) or one vial (250mg.) q.i.d.

Availability:






Orbenin (cloxacillin sodium B.P.)(regd.) is a product of British research at **Beecham Research Laboratories** Brentford, England. Telephone : ISLeworth 4111



21510006655X Published online by Cambridge University Press

What do you do about a case like this?





Now, for the profoundly deaf child and the adult with a severe fragmentary loss-a completely new hearing aid. The Zenith 'Vocalizer' enables hearing correction to be maintained down to frequencies previously beyond the scope of all but desk appliances. The 'Vocalizer' actually amplifies down to one hundred cycles. It can prove helpful for children and adults with some residual hearing for low frequencies but a total loss for the medium and high frequencies. It incorporates new facilities to provide for Loop Listening simultaneously with the child hearing his own voice for speech training. Because the 'Vocalizer' is light yet robust, he can continue to use the same aid at home as he uses at school.

REFER YOUR PATIENTS TO INGRAMS who will gladly provide this unique hearing aid on trial—or any other aid from their comprehensive range of twenty-six different makes. Patients have the widest choice with Ingrams. If you would like details, please write.

ngrams

THE DIFFERENT MAKES **TO CHOOSE FROM**

2 Shepherd St, Shepherd Market, London W1. HYD 9041 Branches throughout Great Britain




for the girl who has everything...(even a stuffy nose)

Otrivine provides gentle, long-lasting relief from the misery of nasal congestion. One application maintains a clear nasal airway for up to eight hours, without subsequent rebound hyperaemia. Suitable for adults and children.

Nasal spray (0.1%) $\frac{1}{2}$ fl. oz, drops (0.1%) $\frac{1}{2}$ fl. oz, and paediatric drops (0.05%) 10 ml. (in special safety dropper-bottle)





In chronic bronchitis **PENBRITIN** helps him put up a better fight

By swiftly reducing both the quantity and purulence of the sputum, routine dosage Penbritin brings a rapid clinical response in the majority of patients: moreover clinical trials have shown these benefits to be more marked than those observed in patients on tetracycline therapy. The success of this regimen (250mg. four times daily for one week or more) can be attributed to Penbritin's higher activity against the causative organisms, H. influenzae and pneumococci.

In addition, severely disabled patients, including those with extensive bronchiectasis and cases whose sputum is winter-long purulent despite other antibiotic therapy, may gain unprecedented relief from a short course of high dosage Penbritin. In published trials on this regimen (1 gram four times daily for one week or more) infection was cleared in the majority of patients, and subsequent freedom from exacerbation for up to 16 weeks was reported.

A full list of supporting references is available on request.



PENBRITIN* (ampicillin B.P.) is a product of British research at **Beecham Research Laboratories**



"Regd.

10. 017/\$002221510006655X Published online by Cambridge University Press




'Triominic' timed-release tablets and orangeflavoured syrup contain antihistamines and a decongestant in balanced proportions to ensure relief from nasal congestion for between 6 and 8 hours. The active constituents of 'Triominic'—the original oral decongestant—reach the nasal mucosa systemically. This obviates the rebound congestion which may follow the topical use of decongestants. 'Triominic' is also available as an orange-flavoured syrup. Basic N.H.S. prices: 50 tablets 5/9½; 4 fl. oz. 2/2 (ex dispensing packs).

Literature and sample on request
A. WANDER LIMITED · 42 UPPER GROSVENOR STREET · LONDON W1

Please mention The Journal of Laryngology and Otology when replying to advertisements

KAMPLEX AUDIOMETRIC EQUIPMENT

AUDIOMETERS



CLINIC AUDIOMETER DA1


ACOUSTIC IMPEDANCE APPARATUS ZA2

PNEUMOMETER ZP1

SKIN RESISTANCE APPARATUS RA4 SCREENING PORTABLE TA16 DIAGNOSTIC PORTABLE TA15

CLINIC AUDIOMETER DA1 DOUBLE CHANNEL DA2

AUTOMATIC AUDIOMETER AB1



SOUND LEVEL METER GA6

VISIBLE SPEECH APPARATUS

AUDITORY TRAINERS

Write for full catalogue and detailed descriptions to:

ALFRED PETERS & SONS LTD 51 GELL STREET, SHEFFIELD 3



Ceporin (cephaloridine) What makes this antibiotic so valuable in the E.N.T. unit?

Spectrum?

Exceptionally broad. It covers most gram-positive organisms—including the penicillin-resistant staphylococci—and many gram-negative organisms. Action?

Intensely bactericidal. It gives a rapid kill of bacteria at or near the minimum inhibitory concentration, especially in the case of gram-negative organisms. Safety?

Wide clinical usage has confirmed the laboratory evidence that Ceporin has a remarkably low toxicity—and it is usually well tolerated by patients allergic to penicillin.

Clinical record?

An analysis of results from over 1500 cases has recently shown high success rates in a wide variety of infections. Ear, nose and throat infections were included in a group of 87 soft tissue infections which showed an 85% satisfactory response. Many reports mention the rapid clinical response obtained with Ceporin after other antibiotics had failed. Conditions which respond well to Ceporin include: mastoiditis, otitis media, sinusitis, quinsy, pharyngeal abscess, and intracranial infections secondary to infected sinuses and mastoiditis. Because of its three main attributesbroad spectrum, bactericidal action and low toxicity—Ceporin has proved valuable for use in place of more toxic drugs or combinations of antibiotics. It also has special value when a quick clinical response is essential in infections caused by penicillin-resistant staphylococci and in the penicillinhypersensitive patient.

Ceporin is issued in vials containing 250 mg, 500 mg and 1 gram of cephaloridine, packed singly and in boxes of five.

Detailed literature is available from Glaxo Laboratories Limited, Greenford, Middlesex.

Ceporin is a Glaxo trade mark