

Editorial

Cite this article: Fisher EW, Sandler D. Virtual issue: ENT care of older adults. *J Laryngol Otol* 2022;**136**:785–787. <https://doi.org/10.1017/S0022215122001207>

Received: 10 May 2022

Accepted: 10 May 2022

First published online: 26 May 2022

Virtual issue: ENT care of older adults

Edward W Fisher^{1,2} and David Sandler³

¹Editor of The Journal of Laryngology & Otology, Birmingham, UK, ²Consultant ENT Surgeon, University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK and ³Consultant Physician, University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK

This virtual issue aims to provide a resource for ENT surgeons and physicians with responsibility for the care of older adults, and draws together relevant articles from the last decade or so of issues from *The Journal of Laryngology & Otology* to give an up-to-date survey of the field.

Demographics

The demographics of the developed world have changed during the late twentieth and early twenty-first centuries such that the care of older adults has assumed ever more importance. In the UK in 2016, there were 11.8 million residents aged 65 years and over, representing 18 per cent of the total population; 25 years before, there were 9.1 million, accounting for 15.8 per cent of the population. The projection is that by 2066, this will have risen to 26 per cent of the population.

Not only are adults living longer, but they have longer ‘active’ lives and are often also economically productive.¹ The reasons for this are numerous and well documented: better nutrition, medical care and fewer wars; a rise in retirement ages well beyond 60 years for both men and women in most Western countries; young families with two working parents; a reduced proportion of young adults in the population compared to older adults; economic barriers to the provision of adequate pensions; and both older adult and child care provision, with the potential this brings for intergenerational conflict. The result is that the later period in a person’s life is unlikely to be spent principally in gardening (‘growing cabbages’, as a retired Roman Emperor was described).

An ‘elderly’ person will spend some of their time in the world of work, whether paid or voluntary, and have an important role in caring for spouses, children or grandchildren, and friends, often in combination with work.² These activities require a fully functioning physiology. Therefore, specialties such as ENT and the healthcare of older adults must strive to limit or ideally abolish disability and handicap in older patients, to allow them to function in these important roles. Both specialties are as much involved in improving quality of life as in extending the lifespan of their patients.

Hearing loss (and olfaction)

Hearing loss in older adults is multifactorial, with the legacy of occupational hearing loss receding as a consequence of the decline of heavy industry, and with current aetiologies dominated by mainly recessive polygenic factors and the numerous lifestyle risk factors for degenerative vascular disorders.³

Most older adults will require diagnosis and treatment for hearing loss. Even in developed countries, there are deficiencies in the provision of an early diagnosis and treatment,⁴ although this provision is much starker in low- and middle-income countries,^{5,6} some of which have little or no audiology or otology service.

Rehabilitative devices such as bone-anchored hearing aids and cochlear implants,⁷ with indications for the latter widening, are increasingly used in older adult patients, and this can be achieved safely. Active research is under way in regenerative medicine, but this is still a long way off from offering practical therapies to restore damaged inner-ear cells.⁸

With the rise in dementia and cognitive disorders in the population, attention has focused on the relationships between hearing impairment, the auditory brain and cognitive decline.⁹ Improved afferent inputs from the use of hearing aids, for example, may have a role in limiting cognitive decline, so there is every reason to seek help for hearing loss. Screening for acoustic neuroma (vestibular schwannoma) is often age-limited, but this needs to be considered on an individual basis.^{10,11} A relationship between cognitive decline and olfaction has been established, in parallel with the relationship with hearing loss, and hyposmia is very common in dementia patients.¹²

Tinnitus

Tinnitus and its inter-dependent relationship with psychological ill health is well recognised;¹³ both conditions are common in the general population and especially in older

adults. Rehabilitative treatments are effective for tinnitus in many cases. Mental health services are under great strain, and audiological rehabilitation cannot entirely compensate for this deficiency.

Necrotising otitis externa

An important, and often overlooked and misdiagnosed condition of the older age group (usually diabetic), is necrotising ('malignant') otitis externa. This condition usually presents with unpleasant and severe ear pain. The condition is on the rise because of increasingly virulent organisms and an increasing prevalence of diabetes mellitus, and is really best described as a 'skull base osteomyelitis'.

'Otitis externa' as a term is misleading, as it is all too easily dismissed as a minor condition. Necrotising otitis externa is often misdiagnosed as a stroke or a 'Bell's palsy' if cranial nerves are involved. Awareness of this treacherous condition among physicians is important as it is associated with significant morbidity and mortality, with the ability to produce lower cranial nerve palsies if it remains untreated. The financial burden of necrotising otitis externa on health services is considerable.^{14–16}

Dizziness and imbalance

Dizziness and imbalance result from disorders and degeneration of peripheral inputs (inner ears, joint and muscle position sense, vision), as well as central nervous system disorders, and any vascular or metabolic upsets that can affect central function.

Managing dizziness and imbalance in older patients often involves many specialties working together, and requires skilled diagnosis and rehabilitation. Studies from many countries show the size of the problem.^{17,18} Simple conditions with a quick and effective treatment such as benign positional vertigo are easily overlooked,¹⁹ but many cases are multifactorial. There is also a link between presbycusis and vestibular deficiency (sometimes referred to as presbyastasis).²⁰ The importance in managing this effectively is amplified if an older person needs to be able to drive.²¹

Swallowing problems

Swallowing problems in older adults are common for neurological or malignant reasons, and there is the interesting condition of pharyngeal pouch.²² Pharyngeal pouches are eminently treatable, but carry high mortality due to aspiration and chest infection if overlooked. Isolated cricopharyngeal dysfunction is another niche condition that is amenable to therapy.²³ In addition, patients have considerable swallowing difficulties after partial laryngectomy, and following treatment for laryngeal and oropharyngeal cancer.^{24,25} The recent coronavirus disease 2019 (Covid-19) pandemic has introduced a whole new range of swallowing problems in older patients, either due to the infection or the treatment.²⁶

The assessment and treatment of voice and swallowing problems are labour-intensive with respect to speech and language therapy services, and such services are usually under-resourced for the wide variety of conditions of all ages that they treat. This includes a wide variety of malignant and non-malignant causes of swallowing dysfunction.^{27,28}

Voice problems

Voice problems in older adults, although common, are somewhat amplified if the patient has a role that involves voice use, which encompasses many occupations, as well as family responsibilities and leisure activities. Most choirs have an older demographic in their membership, and charity shops, and National Trust properties and their guided tours, are staffed largely by retired volunteers. There is a close relationship between voice and hearing function; hence, it is particularly important that hearing rehabilitation takes place in a person who relies on their voice, so that both voice and hearing function are optimised.²⁹

Salivary gland obstruction and sialadenitis

Salivary gland obstruction and sialadenitis are common problems in older adults, with hospital in-patients and care home residents being at particular risk, often due to dehydration and poor oral intake. A recent article in *The Journal* reviews the management of this distressing condition.³⁰

Nose bleeds

Nose bleeds (epistaxis) in older adults have always been associated with significant mortality, often related to associated co-morbidities and the deleterious effects on the cardio-respiratory system of nasal packing and bed rest.^{31,32} This difficult problem has been rendered even more difficult by the rise in prescriptions for anticoagulant agents,³³ particularly the newer oral anticoagulants ('NOACs').

Surgical treatment options

The days when 'lifestyle' surgery, or other ENT, head and neck or thyroid surgical procedure, performed in older adults is dismissed as an option simply because of age are long gone, if they ever existed. The recognition that outcomes are good, and that morbidity and mortality can be reduced to acceptable levels by techniques such as local anaesthesia, has encouraged older adult patients to be offered a wide variety of surgical options, from endoscopic nasal surgery³⁴ to major head and neck surgery,³⁵ although there are times when conservatism is best encouraged, such as for some thyroid cancers.³⁶ One problem when optimising head and neck cancer care in older adults is that such individuals are often left out of cancer trials.³⁷ The topic of head and neck cancer care in the 'elderly' is covered in a very recent article in *The Journal*.³⁸

Head and neck cancer diagnosis

The diagnosis of head and neck cancer aims to be rapid in order to facilitate early treatment, but administrative systems such as the 'two-week-wait' clinics have had limited success, and delays in diagnosis are all too common.^{39,40} These clinics have a disappointing detection rate, and as many cancer patients are diagnosed in the 'routine' system – so getting routine waiting times down to acceptable levels is just as important if timely cancer treatment is the real aim. The Covid-19 pandemic has introduced further barriers to early cancer diagnosis.⁴¹

Liam Flood has reviewed a useful book by Sataloff and Johns which would make good reading for any physician

with an interest in the care of older adults,⁴² as would perusing some of the articles cited in this virtual issue.

References

- 1 Storey A. *Living Longer: How Our Population Is Changing and Why It Matters*. Office for National Statistics: London, 2018
- 2 Storey A, Coombs N, Leib S. *Living Longer: Caring in Later Working Life*. Office for National Statistics. London, 2019
- 3 Bentur N, Valinsky L, Lemberger J, Ben Moshe Y, Heymann AD. Primary care intervention programme to improve early detection of hearing loss in the elderly. *J Laryngol Otol* 2012;**126**:574–9
- 4 Chew HS, Yeak S. Quality of life in patients with untreated age-related hearing loss. *J Laryngol Otol* 2010;**124**:835–41
- 5 Nieman CL, McMahon CM. The World Health Organization's World Report on Hearing: a call to action for hearing care providers. *J Laryngol Otol* 2020;**134**:377–8
- 6 Graydon K, Waterworth C, Miller H, Gunasekera H. Global burden of hearing impairment and ear disease. *J Laryngol Otol* 2019;**133**:18–25
- 7 Hilly O, Hwang E, Smith L, Shipp D, Nedzelski JM, Chen JM *et al*. Cochlear implantation in elderly patients: stability of outcome over time. *J Laryngol Otol* 2016;**130**:706–11
- 8 Jackler RK, Jan TA. The future of otology. *J Laryngol Otol* 2019;**133**:747–58
- 9 Lau K, Dimitriadis PA, Mitchell C, Martyn-St-James M, Hind D, Ray J. Age-related hearing loss and mild cognitive impairment: a meta-analysis and systematic review of population-based studies. *J Laryngol Otol* 2022;**136**:103–18
- 10 Basu S, Youngs R, Mitchell-Innes A. Screening for vestibular schwannoma in the context of an ageing population. *J Laryngol Otol* 2019;**133**:640–9
- 11 Gair J, Borsetto D, Donnelly N, Axon P, Bance M, Tysome JR. Comments on 'Screening for vestibular schwannoma in the context of an ageing population'. *J Laryngol Otol* 2020;**134**:275–6
- 12 Gros A, Manera V, De March CA, Guevara N, König A, Friedman L *et al*. Olfactory disturbances in ageing with and without dementia: towards new diagnostic tools. *J Laryngol Otol* 2017;**131**:572–9
- 13 Pinto PCL, Marcelos CM, Mezzasalma MA, Osterne FJV, de Melo Tavares de Lima MA, Nardi AE. Tinnitus and its association with psychiatric disorders: systematic review. *J Laryngol Otol* 2014;**128**:660–4
- 14 Stapleton E, Watson G. Emerging themes in necrotising otitis externa: a scoping review of the literature 2011–2020 and recommendations for future research. *J Laryngol Otol* 2021. Epub 2021 Oct 20
- 15 Hutson KH, Watson GJ. Malignant otitis externa, an increasing burden in the twenty-first century: review of cases in a UK teaching hospital, with a proposed algorithm for diagnosis and management. *J Laryngol Otol* 2019;**133**:356–62
- 16 Godbehere J, Hutson KH, Watson GJ. Necrotising otitis externa: the increasing financial burden on the National Health Service. *J Laryngol Otol* 2021. Epub 2021 Nov 10
- 17 Soto-Varela A, Faraldo-García A, Del-Río-Valeiras M, Rossi-Izquierdo M, Vaamonde-Sánchez-Andrade I, Gayoso-Diz P *et al*. Adherence of older people with instability in vestibular rehabilitation programmes: prediction criteria. *J Laryngol Otol* 2017;**131**:232–8
- 18 Lasisi AO, Gureje O. Disability and quality of life among community elderly with dizziness: report from the Ibadan Study of Ageing. *J Laryngol Otol* 2010;**124**:957–62
- 19 Lindell E, Karlsson T, Johansson M, Magnusson M, Finizia C. Investigation of self-reported dizziness in the elderly when lying down or turning over in bed, and possible benign paroxysmal positional vertigo. *J Laryngol Otol* 2019;**133**:275–80
- 20 Tan JL, Tang J, Lo S, Yeak S. Investigating the risk factors of vestibular dysfunction and the relationship with presbycusis in Singapore. *J Laryngol Otol* 2016;**130**:816–21
- 21 Sinnott JD, Mahoney H, Wilkinson AS, Broomfield SJ. Dizziness, driving, and the Driver and Vehicle Licensing Agency: audit of advice given to patients, and design of a patient information leaflet. *J Laryngol Otol* 2019;**133**:822–4
- 22 Stewart K, Sen P. Pharyngeal pouch management: an historical review. *J Laryngol Otol* 2016;**130**:116–20
- 23 Ashman A, Dale OT, Baldwin DL. Management of isolated cricopharyngeal dysfunction: systematic review. *J Laryngol Otol* 2016;**130**:611–15
- 24 Seino Y, Miyamoto S, Nakayama M, Yamashita T, Miles A, Allen J. Characteristics that predict penetration and aspiration in elderly patients following supracricoid laryngectomy with cricothyroidopiglotomy: a videofluoroscopic study. *J Laryngol Otol* 2022;**136**:439–46
- 25 Burnip E, Owen SJ, Barker S, Patterson JM. Swallowing outcomes following surgical and non-surgical treatment for advanced laryngeal cancer. *J Laryngol Otol* 2013;**127**:1116–21
- 26 Dawson C, Capewell R, Ellis S, Matthews S, Adamson S, Wood M *et al*. Dysphagia presentation and management following coronavirus disease 2019: an acute care tertiary centre experience. *J Laryngol Otol* 2020;**134**:981–6
- 27 Hanna R, Randall DR. Progression of swallowing dysfunction and associated complications of dysphagia in a cohort of patients with serial videofluoroscopic swallow examinations. *J Laryngol Otol* 2021;**135**:593–8
- 28 Kwon DI, Villegas BC, Ouyoung LM, Sinha UK. Improved swallow outcomes with early intervention using combined swallow therapy in advanced oropharyngeal carcinoma. *J Laryngol Otol* 2022;**136**:433–8
- 29 Rodrigues Dias D, Santos M, Sousa F, Azevedo S, Sousa e Castro S, Freitas S *et al*. How do presbylarynx and presbycusis affect the Voice Handicap Index and the emotional status of the elderly? A prospective case-control study. *J Laryngol Otol* 2021;**135**:1051–6
- 30 Familiari M, Ferrario F, Giordano L, Di Santo D, Indelicato P, Battista RA *et al*. Management of obstructive pathology of the salivary glands in elderly patients: a preliminary study. *J Laryngol Otol* 2022;**136**:60–3
- 31 Corr MJ, Tikka T, Douglas CM, Marshall J. One-year all-cause mortality for 338 patients admitted with epistaxis in a large tertiary ENT centre. *J Laryngol Otol* 2019;**133**:487–93
- 32 Integrate (The National ENT Trainee Research Network). Epistaxis and mortality. *J Laryngol Otol* 2018;**132**:1061–6
- 33 Ho JPK, Bari N, Riffat F. Management of epistaxis in patients on novel oral anticoagulation therapy. *J Laryngol Otol* 2020;**134**:316–22
- 34 Casale M, Costantino A, Sabatino L, Luchena A, Moffa A, Cassano M *et al*. Minimally invasive surgery under local anaesthesia for chronic rhinosinusitis with nasal polyps: our experience in older adults. *J Laryngol Otol* 2020;**134**:145–9
- 35 Subramaniam N, Balasubramanian D, Rka P, Rathod P, Murthy S, Vidhyadharan S *et al*. Peri-operative outcomes following major surgery for head and neck cancer in the elderly: institutional audit and case-control study. *J Laryngol Otol* 2018;**132**:742–7
- 36 Megwalu UC. Observation versus thyroidectomy for papillary thyroid microcarcinoma in the elderly. *J Laryngol Otol* 2017;**131**:173–6
- 37 Abbasi J. Older patients (still) left out of cancer clinical trials. *JAMA* 2019;**322**:1751–3
- 38 Iqbal MS, Navarro-Rodriguez C, Munro S, Ozalp B, Kelly C. Patterns of care for octogenarian patients with head and neck cancer. *J Laryngol Otol* 2022;**136**:(April or May issue)
- 39 Mettias B, Charlton A, Ashokkumar S. Outcome of two-week head and neck cancer pathway for the otolaryngology department in a tertiary centre. *J Laryngol Otol* 2021;**135**:869–73
- 40 Nash R, Hughes J, Sandison A, Stewart S, Clarke P, Mace A. Factors associated with delays in head and neck cancer treatment: case-control study. *J Laryngol Otol* 2015;**129**:383–5
- 41 Taylor R, Omakobia E, Sood S, Glore RJ. The impact of coronavirus disease 2019 on head and neck cancer services: a UK tertiary centre study. *J Laryngol Otol* 2020;**134**:684–7
- 42 Flood LM. Geriatric Otolaryngology. *J Laryngol Otol* 2015;**129**:1042