

Main-sequence and sub-giant stars in the globular cluster NGC 6397: The complex evolution of the lithium abundance

J. I. González Hernández^{1,2,†}, P. Bonifacio^{1,2,3}, E. Caffau¹, M. Steffen⁴,
H.-G. Ludwig^{1,2}, N. Behara^{1,2}, L. Sbordone^{1,2}, R. Cayrel¹, and
S. Zaggia⁵

¹GEPI, Observatoire de Paris, CNRS, Université Paris Diderot;
Place Jules Janssen 92190 Meudon, France
email: Jonay.Gonzalez-Hernandez@obspm.fr

²Cosmological Impact of the First STars (CIFIST) Marie Curie Excellence Team

³Istituto Nazionale di Astrofisica - Osservatorio
Astronomico di Trieste, Italy

⁴Astrophysikalisches Institut Potsdam, An der Sternwarte 16,
D-14482 Potsdam, Germany

⁵INAF - Osservatorio Astronomico di Padova,
Vicolo dell'Osservatorio 5, Padua 35122, Italy

Abstract. Thanks to the high multiplex and efficiency of Giraffe at the VLT we have been able for the first time to observe the Li I doublet in the Main Sequence stars of a globular cluster. At the same time we observed Li in a sample of Sub-Giant stars of the same B-V colour.

Our final sample is composed of 84 SG stars and 79 MS stars. In spite of the fact that SG and MS span the same temperature range we find that the equivalent widths of the Li I doublet in SG stars are systematically larger than those in MS stars, suggesting a higher Li content among SG stars. This is confirmed by our quantitative analysis carried out making use of 1D hydrostatic plane-parallel models and 3D hydrodynamical simulations of the stellar atmospheres.

We derived the effective temperatures of stars in our the sample from H α fitting. Theoretical profiles were computed using 3D hydrodynamical simulations and 1D ATLAS models. Therefore, we are able to determined 1D and 3D-based effective temperatures. We then infer Li abundances taking into account non-local thermodynamical equilibrium effects when using both 1D and 3D models.

We find that SG stars have a mean Li abundance higher by 0.1 dex than MS stars. This result is obtained using both 1D and 3D models. We also detect a positive slope of Li abundance with effective temperature, the higher the temperature the higher the Li abundance, both for SG and MS stars, although the slope is slightly steeper for MS stars. These results provide an unambiguous evidence that the Li abundance changes with evolutionary status.

The physical mechanisms responsible for this behaviour are not yet clear, and none of the existing models seems to describe accurately these observations. Based on these conclusions, we believe that the cosmological lithium problem still remains an open question.

Keywords. Stars: abundances, fundamental parameters, Population II – Galaxy: globular clusters: individual: NGC 6397

† Present address: Dpto. de Astrofísica y Ciencias de la Atmósfera, Facultad de Física, Universidad Complutense de Madrid, E-28040 Madrid, Spain. Email: jonay@astrax.fis.ucm.es

1. Introduction

The determination of the baryonic density from the fluctuations of the cosmic microwave background (CMB) by the WMAP satellite (Spergel *et al.* 2007, Cyburt *et al.* 2008) implies a primordial Li abundance which is $\log(\text{Li}/\text{H}) + 12 = 2.72 \pm 0.06$, at least 0.3–0.5 dex higher than the Li abundance determined in metal-poor stars of the Galactic halo (Spite & Spite 1982).

Many different models of Li depletion have been proposed to explain discrepancy: (a) Piau *et al.* (2006) proposed that the first generation of stars, Population III stars, could have processed some fraction of the halo gas, lowering the lithium abundance; (b) other authors suggest that the primordial Li abundance has been uniformly depleted in the atmospheres of metal-poor dwarfs by some physical mechanism (e.g. turbulent diffusion as in Richard *et al.* (2005), Korn *et al.* (2006); gravitational waves as in Charbonnel & Talon (2005), etc.); and (c) finally, it has been also suggested that the standard Big Bang nucleosynthesis (SBBN) calculations should be revised, possibly with the introduction of new physics as in e.g. Jedamzik (2004), Jedamzik (2006), Jittoh *et al.* (2008), Hisano *et al.* (2009).

Here we present the determination of Li abundances of subgiant (SG) and main-sequence (MS) stars of the cluster NGC 6397. This work provides the first observations of the Li doublet in MS stars of a globular cluster.

2. Observations

We performed spectroscopic observations of the globular cluster NGC 6397 with the multi-object spectrograph FLAMES-GIRAFFE at the VLT on 2007 April, May, June and July, covering the spectral range $\lambda\lambda 6400\text{--}6800 \text{ \AA}$ at resolving power $\lambda/\delta\lambda \sim 17,000$.

We selected subgiant and dwarf stars in the colour range $B - V = 0.60 \pm 0.03$, which ensures that both set of stars fall in a similar and narrow effective temperature range (see Fig. 3 online in González Hernández *et al.* 2009a).

The spectra were reduced with the ESO pipeline and later on treated within MIDAS. We correct the spectra for sky lines and, barycentric and radial velocity. We typically combine 17 spectra of dwarfs and 4 spectra of subgiants to achieve a similar S/N ratio between 80 and 130 in both sets of stars. The mean radial velocity of the cluster stars is $v_r = 18.5 \text{ km s}^{-1}$.

3. Stellar parameters

We derived the effective temperature by fitting the observed $\text{H}\alpha$ line profiles with synthetic profiles, using 3D hydrodynamical model atmospheres computed with the CO⁵BOLD code. The details of this code are provided in Freytag *et al.* (2002) and Wedemeyer *et al.* (2004). The ability of 3D models to reproduce Balmer line profiles has been shown in Behara *et al.* (2009). In that work, the $\text{H}\alpha$ profiles of the Sun as well as the metal-poor stars HD 84937, HD 74000 and HD 140283 are studied. Ludwig *et al.* (2009) also quantified, from a purely theoretical point of view, the difference between the effective temperatures derived from $\text{H}\alpha$ fitting using 1D and 3D models.

We also derived the effective temperatures of MS and SG stars using 1D ATLAS 9 model atmospheres (see Kurucz 2005) and the same fitting procedure. In Fig. 1 we display the histograms of the effective temperatures derived for MS and SG stars using both 1D and 3D models. In the 1D case we got similar effective temperatures for both sets of stars. However, using 3D hydrodynamical models we obtained hotter temperatures by




Figure 1. Histograms of 1D (left panel) and 3D (right panel) effective temperatures derived by fitting the observed $H\alpha$ profiles with theoretical profiles computed using 1D (left panel) and 3D (right panel) model atmospheres, in bins of 100 K, in MS (solid line) and SG (dashed-dotted line) stars in the globular cluster NGC 6397.

approximately 250 K in MS stars and 150 K for SG stars. In the 3D case the histogram of SG stars is slightly shifted with respect to the histogram of MS stars to cooler temperatures, as expected from the difference in surface gravity between MS and SG stars and the sensitivity of the $B - V$ colour to the surface gravity.

Fixed values for the surface gravity were adopted for both subgiant and dwarf stars in the sample, according to the values that best match the position of the stars on a 12 Gyr isochrone (Straniero *et al.* 1997). The adopted values were $\log(g/\text{cm s}^2) = 4.40$ and 3.85 for MS and SG stars, respectively.

4. Li abundances

We measure the equivalent width (EW) of the $\text{Li I } 6708 \text{ \AA}$ line in SG and MS stars by fitting synthetic spectra of known EW to the observed Li profiles. González Hernández *et al.* (2009a) showed the histograms of the EWs measured in SG and MS stars of this cluster (see their Fig. 1). In that figure it is clearly seen that the EWs of SG stars are larger than those of MS stars. They also estimate the weighted mean EW of the SG stars, being ~ 1.1 pm larger than the weighted mean EW of MS stars. Although the colour $B - V$ is sensitive to surface gravity, a priori, this result was not expected, and clearly suggests that subgiants in this cluster have actually higher Li abundances than dwarfs.

We derived Li abundances using 3D model atmospheres. The line formation of Li was treated in non-local thermodynamical equilibrium (NLTE) using the same code and model atom used in Cayrel *et al.* (2007). The analysis was also done using 1D model atmospheres providing essentially the same picture, even when T_{eff} in 1D are lower (see also Fig. 6 online in González Hernández *et al.* 2009a). In the 1D case we used the Carlsson *et al.* (1994) NLTE corrections.

5. Discussion and conclusions

In Fig. 2 we display the histograms of the derived 1D-NLTE and 3D-NLTE Li abundances of dwarf and subgiant stars of the globular cluster NGC 6397. In both 1D and 3D cases, the SG stars have on average larger amounts of Li content in their atmospheres than MS stars. The difference in the mean Li abundance of dwarfs and subgiants is ~ 0.14 dex in the 1D case and ~ 0.07 dex in the 3D case. This difference between the 1D and the 3D cases is due to the effective temperatures derived using 1D and 3D models,




Figure 2. Histograms of 1D (left panel) and 3D (right panel) non-LTE Li abundances, in bins of 0.07 dex, in MS (solid line) and SG (dashed-dotted line) stars in the globular cluster NGC 6397.

and is not related to the NLTE corrections which were independently computed in the 1D and 3D cases. In fact, due to the cooler temperatures derived in 1D with respect to the 3D case, the mean 1D Li abundance lower by ~ 0.13 dex for the MS stars ~ 0.06 dex for the SG stars than the mean 3D abundance.

Lind *et al.* (2009) also find different mean Li abundances, using 1D models, in MSs and SGs, but only by 0.03 dex although still significant at 1σ . However, their result is partially affected by the very narrow range of T_{eff} for MSs deduced by Lind *et al.* (2009) (~ 80 K) compared to the wide range (~ 450 K) for the SGs (see Fig. 7 online in González Hernández *et al.* 2009a).

González Hernández *et al.* (2009a) showed, in their Fig. 2, the 3D-NLTE abundances of SG and MS stars versus 3D effective temperatures (see Fig. 1 in González Hernández *et al.* (2009b) for a similar picture but with T_{eff} and Li abundances computed using 1D models). The points in that figure display a decreasing trend of Li abundance with decreasing temperature. This lithium abundance pattern is different from what is found among field stars (see e.g. Meléndez & Ramírez 2004, Bonifacio *et al.* 2007, González Hernández *et al.* 2008).

Our results imply that the Li surface abundance depends on the evolutionary status of the star. In Fig. 2 of González Hernández *et al.* (2009a), the Li isochrones are shown for different turbulent diffusion models (Richard *et al.* 2005). These models were shifted up by 0.14 dex in Li abundance to make the initial abundance of the models, $\log(\text{Li}/\text{H}) = 2.58$, coincide with the primordial Li abundance predicted from fluctuations of the microwave background measured by the WMAP satellite (Cyburt *et al.* 2008). The models assuming pure atomic diffusion, and, among those including low levels of turbulent mixing, in particular the model T6.0, preferred model in Korn *et al.* (2006) and Lind *et al.* (2009) are ruled out by our observations, since they predict the Li content in MS stars to be higher than in SG stars. The only model that predicts a Li pattern which is qualitatively similar to that observed, is the T6.25 model. This model shows a decreasing trend of Li abundances with decreasing temperatures and also predicts higher abundances for SG stars than for MS stars although at temperatures lower than 6000 K, which is not consistent with the observed trend even in the 1D case (see Fig. 1 in González Hernández *et al.* 2009b).

Models including atomic diffusion and tachocline mixing (Piau 2008) do not seem to reproduce our observations either, since they provide a constant Li abundance up to 5500 K. More sophisticated models are required, for instance, those models that besides including diffusion and rotation also take into account the effect of internal gravity waves

(Talon & Charbonnel 2004), seem to predict accurately the Li abundance pattern in solar-type stars, at solar metallicity (Charbonnel & Talon 2005), but models at low metallicity are still needed.

The cosmological lithium discrepancy still needs to be solved. Given that none of the existing models of Li evolution in stellar atmospheres matches our observations in the globular cluster NGC 6397, we hope our results will prompt further new theoretical investigations.

References

- Behara, N. T., Ludwig, H.-G., Steffen, M., & Bonifacio, P. 2009, *American Institute of Physics Conference Series*, 1094, 784
- Bonifacio, P., Molaro, P., Sivarani, T., Cayrel, R., & Spite, M. 2007, *A&A*, 462, 851
- Carlsson, M., Rutten, R. J., Bruls, J. H. M. J., & Shchukina, N. G. 1994, *A&A*, 288, 860
- Cayrel, R., Steffen, M., Chand, H., Bonifacio, P., Spite, M., Spite, F., Petitjean, P., Ludwig, H. G., & Caffau, E. 2007, *A&A (Letters)*, 473, 37
- Charbonnel, C. & Talon, S. 2005, *Science*, 309, 2189
- Cyburt, R. H., Fields, B. D., & Olive, K. A. 2008, *Journal of Cosmology and Astro-Particle Physics*, 11, 12
- Freytag, B., Steffen, M., & Dorch, B. 2002, *Astronomische Nachrichten*, 323, 213
- González Hernández, J. I. *et al.* 2008, *A&A*, 480, 233
- González Hernández, J. I., Bonifacio, P., Caffau, E., Steffen, M., Ludwig, H.-G., Behara, N. T., Sbordone, L., Cayrel, R., & Zaggia, S. 2009a, *A&A (Letters)*, 505, L13
- González Hernández, J. I. *et al.* 2009b, *Star Clusters: Basic Galactic Building Blocks Throughout Time And Space*. Edited by R. de Grijs and J. Lepine. Proceedings of IAU Symposium #266, held 10-14 August, 2009 in Brazil, Rio de Janeiro. Cambridge, UK: *Cambridge University Press*, in press
- Hisano, J., Kawasaki, M., Kohri, K., & Nakayama, K. 2009, *Phys. Rev. D.*, 79, 063514
- Jedamzik, K. 2004, *Phys. Rev. D.*, 70, 083510
- Jedamzik, K. 2006, *Phys. Rev. D.*, 74, 103509
- Jittoh, T., Kohri, K., Koike, M., Sato, J., Shimomura, T., & Yamanaka, M. 2008, *Phys. Rev. D.*, 78, 055007
- Kurucz, R. L. 2005, *Memorie della Società Astronomica Italiana Supplement*, 8, 14
- Korn, A. J., Grundahl, F., Richard, O., Barklem, P. S., Mashonkina, L., Collet, R., Piskunov, N., & Gustafsson, B. 2006, *Nature*, 442, 657
- Lind, K., Primas, F., Charbonnel, C., Grundahl, F., & Asplund, M. 2009, *A&A*, 503, 545
- Ludwig, H.-G., Behara, N. T., Steffen, M., & Bonifacio, P. 2009, *A&A (Letters)*, 502, 1
- Meléndez, J. & Ramírez, I. 2004, *ApJ (Letters)*, 615, 33
- Piau, L., Beers, T. C., Balsara, D. S., Sivarani, T., Truran, J. W., & Ferguson, J. W. 2006, *ApJ*, 653, 300
- Piau, L. 2008, *ApJ*, 689, 1279
- Richard, O., Michaud, G., & Richer, J. 2005, *ApJ*, 619, 538
- Spergel, D. N., Bean, R. Doré, O., Nolta, M. R., Bennett, C. L. *et al.* 2007 *ApJS*, 170, 377
- Spite, M. & Spite, F. 1982, *Nature*, 297, 483
- Straniero, O., Chieffi, A., & Limongi, M. 1997, *ApJ*, 490, 425
- Talon, S. & Charbonnel, C. 2004, *A&A*, 418, 1051
- Wedemeyer, S., Freytag, B., Steffen, M., Ludwig, H.-G., & Holweger, H. 2004, *A&A*, 414, 1121



Monique Spite & Katia Cunha



Corinne Charbonnel & Francesca Primas