

and (3) *C. difficile* negative. **Results:** A diagnosis of *C. difficile* adds significantly (>\$3,000) to unadjusted hospital cost compared to a negative result. Propensity-adjusted analyses demonstrated that *C. difficile* colonization was associated with significantly increased (median, \$5,000) hospital cost whereas any positive or true diagnoses of *C. difficile* were not associated with increased cost. Colonized patients also had significantly higher lengths of stay (1 day) and cost per length of stay (\$218 per day). **Conclusions:** This is the first *C. difficile* cost analysis to utilize PCR CT data to differentiate colonization. Surprisingly, patients with a high CT had disproportionately higher hospital costs compared to matched *C. difficile*-negative patients, which was not seen among patients with a low CT or with any positive result. We hypothesize that this unexpected finding may be due to misdiagnosis and mistreatment of diarrhea not caused by *C. difficile* or unadjusted factors associated with high cost and non-*C. difficile* diarrhea. In addition, the discrepantly high cost attributed to *C. difficile* diagnosis cited in the literature (\$3,000–11,000 per hospitalized case) could be explained by the common use of administrative data to identify *C. difficile* cases and controls as opposed to our study, which directly linked cost data to *C. difficile*-positive and -negative test results.

Funding: None

Disclosures: None

Doi:10.1017/ice.2020.908

Presentation Type:

Poster Presentation

Meta-analysis of Outcomes Using Ceftolozane-Tazobactam and Ceftazidime-Avibactam for Multidrug-Resistant Organism Infections

Geneva Wilson, University of Iowa; Margaret Fitzpatrick, Loyola University Chicago Stritch School of Medicine; Kyle Walding, Loyola University Medical Center; Beverly Gonzalez, Edward Hines Jr. VA; Charlesnika Evans, Northwestern University, Feinberg School of Medicine

Background: Ceftolozane-tazobactam (C/T) and ceftazidime-avibactam (C/A) are new β -lactam/ β -lactamase combination antibiotics that were approved by the FDA in 2014 and 2015, respectively, to treat complicated intra-abdominal and urinary tract infections. They are commonly used to treat multidrug-resistant *Pseudomonas aeruginosa* (MDRPA) and carbapenem-resistant *Enterobacteriaceae* (CRE) infections at any site. Both medications are also often used as salvage therapy when empiric therapy has failed or when the infectious organism tests resistant to all other available antibiotics. The purpose of this review is to present the clinical experience and reported clinical success rates of C/T and C/A. **Methods:** PubMed, EMBASE, and Google Scholar were searched from January 1, 2013, through October 1, 2019, for publications detailing clinical experience with C/T and C/A in patients with CRE and MDRPA infections. Included study designs were extended cases series and clinical observational studies. Information on infection type, bacterial agent, salvage therapy uses, clinical success, and resistance development during treatment were abstracted. Meta-regression analysis was used to determine the pooled effectiveness of C/T and C/A among included studies. **Results:** The literature search returned 1,645 publications. After exclusion criteria were applied, 16 publications representing 769 patients were retained. The study population was mostly male (pooled average, 62%). The major comorbidities represented in the pooled population were solid organ transplantation (20.0%),

kidney disease (19.5%), cardiovascular disease (15.3%), and diabetes (15.3%). Pneumonia was the predominant infection type (41.4%) and MDRPA was the pathogen most frequently evaluated (57.7%). The pooled clinical success rate was 70.2% (95% CI, 64.5%–75.3%). Also, 10 studies explicitly evaluated C/A or C/T as salvage therapy. The pooled clinical success rate for salvage therapy studies was 75.2% (95% CI, 69.7%–80.0%). Development of resistance to C/T or C/A during or after treatment was reported for 2.0% of the population. **Conclusion:** Overall, these medications have a high clinical success rate in patients with severe and complicated infections and limited treatment options. Pooled clinical success rates were high (70.2%) and the medications were particularly effective as salvage therapy. Resistance rates were low, although this could have been biased by the small percentage of studies that reported on this outcome. More longitudinal studies comparing the effectiveness of C/T and C/A against other antibiotic regimens are needed.

Funding: None

Disclosures: None

Doi:10.1017/ice.2020.909

Presentation Type:

Poster Presentation

Methicillin-Resistant *Staphylococcus aureus* Prevalence Among Healthcare Workers in Contact Tracings in a Dutch Hospital

Veronica Weterings, Amphia Hospital; Heidi Kievits, Amphia Hospital; Miranda van Rijen, Amphia Hospital; Jan Kluytmans, Amphia Hospital

Background: In The Netherlands, the national guidelines on Methicillin-Resistant *Staphylococcus aureus* (MRSA) prevention and control advocate screening of healthcare workers (HCWs) after unprotected exposure to MRSA carriers. Although this strategy is largely successful, contact tracing of staff is a time-consuming and costly component. We evaluated our contact tracing policy for HCWs over the years 2010–2018. **Methods:** A retrospective, observational study was performed in a Dutch teaching hospital. All HCWs who had unprotected contact with an MRSA carrier were included in contact tracing. When there had been a long period of unprotected admission prior to an MRSA finding, or when the index case was an HCW, the entire (nursing) team was tested. All samples of HCWs who were tested for MRSA carriage as part of contact tracing from 2010 until 2018 were included. A pooled nose, throat, and perineum swab was collected using the eSwab medium (Copan) and inoculated on chromID MRSA agar plates (bioMérieux) after enrichment in a broth. Molecular typing was performed using multiple-locus variable number of tandem repeat analysis (MLVA). **Results:** In total, we included 8,849 samples (range, 677–1,448 samples per year) from 287 contact tracings (range, 26–55 contact tracings per year). Overall, 32 HCWs were colonized with MRSA (0.36%; 95% CI, 0.26%–0.51%). None of them developed a clinical infection. Moreover, 8 HCWs (0.10%; 95% CI, 0.05%–0.19%) were colonized with the same MLVA type as the index case and were detected in 6 of 287 contact tracings (2%). In 4 of 8 of these cases, a positive HCW was the index for undertaking contact tracing. In 3 of 8 cases, it was clear that the HCW who was identified in the contact tracing was the source of the outbreak and was the cause of invasive MRSA infections in patients. Notably, a different MLVA type as the index case was found in 24 HCWs (0.27%; 95% CI, 0.18%–0.40%) of whom 7 of 24 HCWs (29.2%) were intermittent carriers. **Conclusions:** This study revealed a sustained low MRSA prevalence among

samples in contact tracing of HCWs over 9 years. Furthermore, it shows that when MRSA contact tracing is performed according to the national guideline, only 1 of 1,000 samples results in a secondary case. This is similar to the population carriage rate of MRSA in The Netherlands. More frequently, an unrelated strain is found. These findings raise questions regarding the efficacy of the current strategy to perform contact tracing after unprotected exposure.

Funding: None

Disclosures: None

Doi:10.1017/ice.2020.910

Presentation Type:

Poster Presentation

Middle East Respiratory Syndrome Coronavirus, Saudi Arabia, 2017 Outbreak: Super-Spreading Event and Control Measures
Hala Amer, National Research Center; Faisal Alaklobi; Ziad Memish; Hind Alzoman

Background: The hallmark of Middle East respiratory syndrome coronavirus (MERS-CoV) disease is the ability to cause major healthcare-associated outbreaks with superspreading events leading to massive transmission and excessive morbidity and mortality. This abstract provides overview of MERS-CoV multi-healthcare facilities outbreak in Riyadh in June 2017, with focus on cluster reported and the control measures taken at King Saud Medical City. The outbreak began with a patient who presented with acute renal failure requiring hemodialysis and became a MERS-CoV superspreader, igniting the cluster of cases in several hospitals in King Saud Medical City. **Methods:** For epidemiologic investigation, a case was defined as any patient with laboratory-confirmed MERS-CoV infection with connection to the affected healthcare facilities. Contact tracing and testing were performed according to the Ministry of Health (MOH) guidelines. MERS-CoV testing was recommended for HCWs who had unprotected close contact with a confirmed case. Considering the superspreading phenomena, contact tracing was included all persons attended the same area with the positive case either as a patient, an HCW, or a patient's visitor or companion. Laboratory confirmation was conducted using real-time RT-PCR. Genome sequencing and phylogenetic analysis were performed for available MERS-CoV rRT-PCR-positive samples by the CDC. The infection control measures applied included decreasing patient load through downsizing

Spider chart of Middle East respiratory syndrome coronavirus outbreak, King Saud Medical City cluster




Fig. 1.

Mapping of the patient's journey with numbers of contacts. ED, emergency department; HCW, health care worker, ICU, intensive care unit.




Fig. 2.

emergency department acceptance, maintaining low elective services, limiting inpatient admissions, and encouraging discharge. Early detection and quarantining of any suspected cases took place through extensive contact tracing, properly triaging all patients upon admission, consistent monitoring of inpatients and HCWs for any emerging acute respiratory illness, allocation of more single rooms inside the facility and staff dormitory, and extending the services of virology laboratory to get timely results. Further measures consisted of extensive education on infection control practices, monitoring healthcare worker adherence, reassuring the public by maintaining transparency of published reports, and launching a hotline to respond to HCW concerns. Moreover, travel restrictions applied to any person with a history of exposure to a confirmed MERS-CoV case during the 2-week monitoring period. **Results:** Overall, 44 cases of MERS-CoV infection were reported from 3 simultaneous clusters during the 2017 Riyadh outbreak, including 11 fatal cases. Among all of the cases, 29 cases were reported at King Saud Medical City. The outbreak at KSMC required 30 days to be controlled. **Conclusions:** High vigilance for early detection is a key control measure. To be more sensitive, point-of-care MERS-CoV testing is required because clinical suspicion is challenging in patients presenting with acute renal failure.

Funding: None

Disclosures: None

Doi:10.1017/ice.2020.911

Presentation Type:

Poster Presentation

Minimum Healthcare Worker Influenza Immunization Rates Required to Decrease Influenza Transmission in Acute-Care Inpatients

Victoria Williams, Sunnybrook Health Sciences Centre; Wendy Morgan, Sunnybrook Health Sciences Centre, Toronto, ON, Canada; Maja McGuire, Infection Prevention and Control, North York General Hospital, Toronto, ON Canada; Kevin Katz, North York General Hospital; Jerome Leis, University of Toronto

Background: Annual influenza immunization of healthcare workers (HCWs) is widely recommended to reduce the risk of healthcare-associated influenza (HAI). Although there is a clear association between