

ARTICLE

# Making $K_{r+1}$ -free graphs $r$ -partite

József Balogh<sup>1,†</sup>, Felix Christian Clemen<sup>2</sup>, Mikhail Lavrov<sup>2</sup>, Bernard Lidický<sup>3,\*,‡</sup> and Florian Pfender<sup>4,§</sup>

<sup>1</sup>Department of Mathematics, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA, and Moscow Institute of Physics and Technology, Russian Federation, <sup>2</sup>Department of Mathematics, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA, <sup>3</sup>Department of Mathematics, Iowa State University, Ames, IA 50011, USA and <sup>4</sup>Department of Mathematical and Statistical Sciences, University of Colorado Denver, CO 80217-3364, USA

\*Corresponding author. Email: lidicky@iastate.edu

(Received 13 December 2019; revised 26 September 2020; accepted 10 September 2020)

## Abstract

The Erdős–Simonovits stability theorem states that for all  $\varepsilon > 0$  there exists  $\alpha > 0$  such that if  $G$  is a  $K_{r+1}$ -free graph on  $n$  vertices with  $e(G) > \text{ex}(n, K_{r+1}) - \alpha n^2$ , then one can remove  $\varepsilon n^2$  edges from  $G$  to obtain an  $r$ -partite graph. Füredi gave a short proof that one can choose  $\alpha = \varepsilon$ . We give a bound for the relationship of  $\alpha$  and  $\varepsilon$  which is asymptotically sharp as  $\varepsilon \rightarrow 0$ .

2020 MSC Codes: 05C35

## 1. Introduction

Erdős asked how many edges need to be removed in a triangle-free graph on  $n$  vertices in order to make it bipartite. He conjectured that the balanced blow-up of  $C_5$  with class sizes  $n/5$  is the worst case, and hence  $n^2/25$  edges would always be sufficient. Together with Faudree, Pach and Spencer [6], he proved that one can remove at most  $n^2/18$  edges to make a triangle-free graph bipartite.

Further, Erdős, Györi and Simonovits [7] proved that for graphs with at least  $n^2/5$  edges, an unbalanced  $C_5$  blow-up is the worst case. For  $r \in \mathbb{N}$ , let  $D_r(G)$  denote the minimum number of edges which need to be removed to make  $G$   $r$ -partite.

**Theorem 1.1 (Erdős, Györi and Simonovits [7]).** *Let  $G$  be a  $K_3$ -free graph on  $n$  vertices with at least  $n^2/5$  edges. There exists an unbalanced  $C_5$  blow-up of  $H$  with  $e(H) \geq e(G)$  such that*

$$D_2(G) \leq D_2(H). \quad (1.1)$$

This proved the Erdős conjecture for graphs with at least  $n^2/5$  edges. A simple probabilistic argument (e.g. [7]) settles the conjecture for graphs with at most  $2/25n^2$  edges.

A related question was studied by Sudakov; he determined the maximum number of edges in a  $K_4$ -free graph which need to be removed in order to make it bipartite [16]. This problem for  $K_6$ -free graphs was solved by Hu, Lidický, Martins, Norin and Volec [11].

<sup>†</sup>Research is partially supported by NSF grant DMS-1764123, Arnold O. Beckman Research Award (UIUC Campus Research Board RB 18132) and the Langan Scholar Fund (UIUC).

<sup>‡</sup>Research of this author is partially supported by NSF grant DMS-1855653.

<sup>§</sup>Research of this author is partially supported by NSF grant DMS-1855622.

We will study the question of how many edges in a  $K_{r+1}$ -free graph need, at most, to be removed to make it  $r$ -partite. For  $n \in \mathbb{N}$  and a graph  $H$ , let  $\text{ex}(n, H)$  denote the Turán number, *i.e.* the maximum number of edges of an  $H$ -free graph. The Erdős–Simonovits theorem [8] for cliques states that for every  $\varepsilon > 0$  there exists  $\alpha > 0$  such that if  $G$  is a  $K_{r+1}$ -free graph on  $n$  vertices with  $e(G) > \text{ex}(n, K_{r+1}) - \alpha n^2$ , then  $D_r(G) \leq \varepsilon n^2$ .

Füredi [9] gave a nice short proof of the statement that a  $K_{r+1}$ -free graph  $G$  on  $n$  vertices with at least  $\text{ex}(n, K_{r+1}) - t$  edges satisfies  $D_r(G) \leq t$ , and thus provided a quantitative version of the Erdős–Simonovits theorem.

In [11] Füredi’s result was strengthened for some values of  $r$ . Roberts and Scott [15] showed that  $D_r(G) = O(t^{3/2}/n)$  when  $t \leq \delta n^2$ , and that this result is sharp up to a constant factor. They also proved a more general result for  $H$ -free graphs where  $H$  is an edge-critical graph. For small  $t$ , we will determine asymptotically how many edges are needed. For very small  $t$ , it is already known [4] that  $G$  has to be  $r$ -partite, as the following theorem shows.

**Theorem 1.2 (Brouwer [4]).** *Let  $r \geq 2$  and  $n \geq 2r + 1$  be integers. Let  $G$  be a  $K_{r+1}$ -free graph on  $n$  vertices with  $e(G) \geq \text{ex}(n, K_{r+1}) - \lfloor n/r \rfloor + 2$ . Then*

$$D_r(G) = 0. \tag{1.2}$$

This phenomenon was also studied in [1], [10], [12] and [18]. We will be studying  $K_{r+1}$ -free graphs on fewer edges. For these, our main result is the following theorem.

**Theorem 1.3.** *Let  $r \geq 2$  be an integer. Then, for all  $n \geq 3r^2$  and for all  $0 \leq \alpha \leq 10^{-7}r^{-12}$ , the following holds. Let  $G$  be a  $K_{r+1}$ -free graph on  $n$  vertices with*

$$e(G) \geq \text{ex}(n, K_{r+1}) - \alpha n^2. \tag{1.3}$$

Then

$$D_r(G) \leq \left( \frac{2r}{3\sqrt{3}} + o_\alpha(1) \right) \alpha^{3/2} n^2, \tag{1.4}$$

where  $o_\alpha(1)$  is a term converging to 0 for  $\alpha$  tending to 0.

Note that we did not try to optimize our bounds on  $n$  and  $\alpha$  in the theorem.

The blow-up of a graph  $G$  is obtained by replacing every vertex  $v \in V(G)$  with finitely many copies so that the copies of two vertices are adjacent if and only if the originals are.

For two graphs  $G$  and  $H$ , we define  $G \otimes H$  to be the graph on the vertex set  $V(G) \cup V(H)$  with  $gg' \in E(G \otimes H)$  if and only if  $gg' \in E(G)$ ,  $hh' \in E(G \otimes H)$  if and only if  $hh' \in E(H)$ , and  $gh \in E(G \otimes H)$  for all  $g \in V(G)$ ,  $h \in V(H)$ .

We will prove that Theorem 1.3 is asymptotically sharp by describing an unbalanced blow-up of  $K_{r-2} \otimes C_5$  that needs at least that many edges to be removed to make it  $r$ -partite. Our extremal example appeared first (with different class sizes) in a paper by Andrásfai, Erdős and Sós [2].

**Theorem 1.4.** *Let  $r, n \in \mathbb{N}$  and  $0 < \alpha < 1/(4r^4)$ . Then there exists a  $K_{r+1}$ -free graph  $G$  on  $n$  vertices with*

$$e(G) \geq \text{ex}(n, K_{r+1}) - \alpha n^2 + \frac{4r}{3\sqrt{3}} \alpha^{3/2} n^2 - \frac{2r(r-3)}{9} \alpha^2 n^2$$

and

$$D_r(G) \geq \frac{2r}{3\sqrt{3}} \alpha^{3/2} n^2.$$

In Kang and Pikhurko’s proof [12] of Theorem 1.2, the case  $e(G) = \text{ex}(n, K_{r+1}) - \lfloor n/r \rfloor + 1$  is studied. In this case they constructed a family of  $K_{r+1}$ -free non- $r$ -partite graphs, which includes our extremal graph, for that number of edges.

We conjecture that our extremal example needs the most edges removed to make it  $r$ -partite among all  $K_{r+1}$ -free graphs with many edges.

**Conjecture 1.5.** *Let  $r \geq 2$  be an integer and let  $n$  be sufficiently large. Then there exists  $\alpha_0 > 0$  such that for all  $0 \leq \alpha \leq \alpha_0$  the following holds. For every  $K_{r+1}$ -free graph  $G$  on  $n$  vertices there exists an unbalanced  $K_{r-2} \otimes C_5$  blow-up  $H$  on  $n$  vertices with  $e(H) \geq e(G)$  such that*

$$D_r(G) \leq D_r(H). \tag{1.5}$$

This conjecture can be seen as a generalization of Theorem 1.1. Note that Conjecture 1.5 was recently proved by Korándi, Roberts and Scott [13]. We recommend the interested reader to read the excellent survey [14] by Nikiforov. He gives a good overview on further related stability results, for example on guaranteeing large induced  $r$ -partite subgraphs of  $K_{r+1}$ -free graphs.

We organize the paper as follows. In Section 2 we prove Theorem 1.3 and in Section 3 we give the sharpness example, *i.e.* we prove Theorem 1.4.

## 2. Proof of Theorem 1.3

In this section we prove the following version of Theorem 1.3, which gives better control over the error term.

**Theorem 2.1.** *Let  $r \geq 2$  be an integer. Then, for all  $n \geq 3r^2$  and for all  $0 \leq \alpha \leq 10^{-7}r^{-12}$ , the following holds. Let  $G$  be a  $K_{r+1}$ -free graph on  $n$  vertices with*

$$e(G) \geq \text{ex}(n, K_{r+1}) - \alpha n^2. \tag{2.1}$$

Then

$$D_r(G) \leq \left( \frac{2r}{3\sqrt{3}} + 30r^3\alpha^{1/6} \right) \alpha^{3/2} n^2. \tag{2.2}$$

Let  $G$  be an  $n$ -vertex  $K_{r+1}$ -free graph with  $e(G) \geq \text{ex}(n, K_{r+1}) - t$ , where  $t = \alpha n^2$ . We will assume that  $n$  is sufficiently large. Furthermore, by Theorem 1.2 we can assume that

$$\alpha \geq \frac{\lfloor n/r \rfloor - 2}{n^2} \geq \frac{1}{2rn}. \tag{2.3}$$

This also implies that  $t \geq r$  because  $n \geq 3r^2$ . During our proof we will make use of Turán’s theorem and a version of Turán’s theorem for  $r$ -partite graphs on multiple occasions. Turán’s theorem [17] determines the maximum number of edges in a  $K_{r+1}$ -free graph.

**Theorem 2.2 (Turán [17]).** *Let  $r \geq 2$  and  $n \in \mathbb{N}$ . Then*

$$\frac{n^2}{2} \left( 1 - \frac{1}{r} \right) - \frac{r}{2} \leq \text{ex}(n, K_{r+1}) \leq \frac{n^2}{2} \left( 1 - \frac{1}{r} \right).$$

Let  $K(n_1, \dots, n_r)$  denote the complete  $r$ -partite graph whose  $r$  colour classes have sizes  $n_1, \dots, n_r$ , respectively. Turán’s theorem for  $r$ -partite graphs states the following.

**Theorem 2.3 (folklore).** *Let  $r \geq 2$  and  $n_1, \dots, n_r \in \mathbb{N}$  satisfying  $n_1 \leq \dots \leq n_r$ . For a  $K_r$ -free subgraph  $H$  of  $K(n_1, \dots, n_r)$ , we have*

$$e(H) \leq e(K(n_1, \dots, n_r)) - n_1 n_2.$$

For a proof of this folklore result see [3, Lemma 3.3], for example.

We denote the maximum degree of  $G$  by  $\Delta(G)$ . For two disjoint subsets  $U, W$  of  $V(G)$ , write  $e(U, W)$  for the number of edges in  $G$  with one endpoint in  $U$  and the other endpoint in  $W$ . We write  $e^c(U, W)$  for the number of non-edges between  $U$  and  $W$ , i.e.  $e^c(U, W) = |U||W| - e(U, W)$ .

Füredi [9] used Erdős’s degree majorization algorithm [5] to find a vertex partition with some useful properties. We include the proof for completeness.

**Lemma 2.1 (Füredi [9]).** *Let  $t, r, n \in \mathbb{N}$  and  $G$  be an  $n$ -vertex  $K_{r+1}$ -free graph with  $e(G) \geq \text{ex}(n, K_{r+1}) - t$ . Then there exists a vertex partition  $V(G) = V_1 \cup \dots \cup V_r$  such that*

$$\sum_{i=1}^r e(G[V_i]) \leq t, \quad \Delta(G) = \sum_{i=2}^r |V_i| \quad \text{and} \quad \sum_{1 \leq i < j \leq r} e^c(V_i, V_j) \leq 2t. \tag{2.4}$$

**Proof.** Let  $x_1 \in V(G)$  be a vertex of maximum degree. Define  $V_1 := V(G) \setminus N(x_1)$  and  $V_1^+ = N(x_1)$ . Iteratively, let  $x_i$  be a vertex of maximum degree in  $G[V_{i-1}^+]$ . Let  $V_i := V_{i-1}^+ \setminus N(x_i)$  and  $V_i^+ = V_{i-1}^+ \cap N(x_i)$ . Since  $G$  is  $K_{r+1}$ -free this process stops at  $i \leq r$  and thus gives a vertex partition  $V(G) = V_1 \cup \dots \cup V_r$ . Summing up the degrees of vertices in  $V_1$ , we have

$$2e(G[V_1]) + e(V_1, V_1^+) = \sum_{x \in V_1} \deg(x) \leq |V_1||V_1^+|,$$

and similarly for the other classes

$$2e(G[V_i]) + e(V_i, V_i^+) = \sum_{x \in V_i} \deg_{G[V_{i-1}^+]}(x) \leq |V_i||V_i^+|.$$

Adding up these inequalities we get

$$\text{ex}(n, K_{r+1}) - t + \sum_{i=1}^r e(G[V_i]) = e(G) + \sum_{i=1}^r e(G[V_i]) \leq \sum_{i=1}^{r-1} |V_i||V_i^+| \leq \text{ex}(n, K_{r+1}),$$

implying

$$\sum_{i=1}^r e(G[V_i]) \leq t.$$

By construction,

$$\sum_{i=2}^r |V_i| = |V_1^+| = |N(x_1)| = \Delta(G).$$

Let  $H$  be the complete  $r$ -partite graph with vertex set  $V(G)$  and all edges between  $V_i$  and  $V_j$  for  $1 \leq i < j \leq r$ . The graph  $H$  is  $r$ -partite and thus has at most  $\text{ex}(n, K_{r+1})$  edges. Finally, since  $G$  has at most  $t$  edges not in  $H$  and at least  $\text{ex}(n, K_{r+1}) - t$  edges in total, at most  $2t$  edges of  $H$  can be missing from  $G$ , giving us

$$\sum_{1 \leq i < j \leq r} e^c(V_i, V_j) \leq 2t$$

and proving the last inequality. □

For this vertex partition we can get bounds on the class sizes.

**Lemma 2.2.** For all  $i \in [r]$ ,

$$|V_i| \in \left\{ \frac{n}{r} - \frac{5}{2}\sqrt{\alpha n}, \frac{n}{r} + \frac{5}{2}\sqrt{\alpha n} \right\},$$

and thus also

$$\Delta(G) \leq \frac{r-1}{r}n + \frac{5}{2}\sqrt{\alpha n}.$$

**Proof.** We know that

$$\sum_{1 \leq i < j \leq r} |V_i||V_j| \geq e(G) - \sum_{i=1}^r e(G[V_i]) \geq \left(1 - \frac{1}{r}\right)\frac{n^2}{2} - \frac{r}{2} - 2t.$$

Also

$$\sum_{1 \leq i < j \leq r} |V_i||V_j| = \frac{1}{2} \sum_{i=1}^r |V_i|(n - |V_i|) = \frac{n^2}{2} - \frac{1}{2} \sum_{i=1}^r |V_i|^2.$$

Thus we can conclude that

$$\sum_{i=1}^r |V_i|^2 \leq \frac{n^2}{r} + r + 4t. \tag{2.5}$$

Now let  $x = |V_1| - n/r$ . Then

$$\begin{aligned} \sum_{i=1}^r |V_i|^2 &= \left(\frac{n}{r} + x\right)^2 + \sum_{i=2}^r |V_i|^2 \\ &\geq \left(\frac{n}{r} + x\right)^2 + \frac{(\sum_{i=2}^r |V_i|)^2}{r-1} \\ &\geq \left(\frac{n}{r} + x\right)^2 + \frac{(n(1 - 1/r) - x)^2}{r-1} \\ &\geq \frac{n^2}{r} + x^2. \end{aligned} \tag{2.6}$$

Combining this with (2.5), we get

$$|x| \leq \sqrt{r + 4t} \leq \frac{5}{2}\sqrt{t} = \frac{5}{2}\sqrt{\alpha n},$$

and thus

$$\frac{n}{r} - \frac{5}{2}\sqrt{\alpha n} \leq |V_1| \leq \frac{n}{r} + \frac{5}{2}\sqrt{\alpha n}.$$

In a similar way we get the bounds on the sizes of the other classes. □

**Lemma 2.3.** The graph  $G$  contains  $r$  vertices  $x_1 \in V_1, \dots, x_r \in V_r$  which form a  $K_r$ , and for every  $i$

$$\deg(x_i) \geq n - |V_i| - 5r\alpha n.$$

**Proof.** Let  $V_i^c := V(G) \setminus V_i$ . We call a vertex  $v_i \in V_i$  *small* if  $|N(v_i) \cap V_i^c| < |V_i^c| - 5r\alpha n$  and *big* otherwise. For  $1 \leq i \leq r$ , let  $B_i$  denote the set of big vertices inside class  $V_i$ . There are at most

$$\frac{4t}{5r\alpha n} = \frac{4}{5r}n$$

small vertices in total as otherwise (2.4) is violated. Thus in each class there are at least  $n/10r$  big vertices, i.e.  $|B_i| \geq n/10r$ . The number of missing edges between the sets  $B_1, \dots, B_r$  is at most  $2t < \frac{1}{100r^2}n^2$ . Thus, using Theorem 2.3, we can find a  $K_r$  with one vertex from each  $B_i$ .  $\square$

**Lemma 2.4.** *There exists a vertex partition  $V(G) = X_1 \cup \dots \cup X_r \cup X$  such that the  $X_i$  are independent sets,  $|X| \leq 5r^2\alpha n$  and*

$$\frac{n}{r} - 3\sqrt{\alpha n} \leq |X_i| \leq \frac{n}{r} + 3r\sqrt{\alpha n}$$

for all  $1 \leq i \leq r$ .

**Proof.** By Lemma 2.3 we can find vertices  $x_1, \dots, x_r$  forming a  $K_r$  and having  $\deg(x_i) \geq n - |V_i| - 5r\alpha n$ . Define  $X_i$  to be the common neighbourhood of  $x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_r$  and  $X = V(G) \setminus (X_1 \cup \dots \cup X_r)$ . Since  $G$  is  $K_{r+1}$ -free, the  $X_i$  are independent sets. Now we bound the size of  $X_i$  using the bounds on the sets  $V_i$ . Since every  $x_j$  has at most  $|V_j| + 5r\alpha n$  non-neighbours, we get

$$|X_i| \geq n - \sum_{\substack{1 \leq j \leq r \\ j \neq i}} (|V_j| + 5r\alpha n) \geq |V_i| - 5r^2\alpha n \geq \frac{n}{r} - 3\sqrt{\alpha n} \tag{2.7}$$

and

$$\sum_{i=1}^r \deg(x_i) \geq n(r-1) - 5r^2\alpha n. \tag{2.8}$$

A vertex  $v \in V(G)$  cannot be incident to all of the vertices  $x_1, \dots, x_r$ , because  $G$  is  $K_{r+1}$ -free. Further, every vertex from  $X$  is not incident to at least two of the vertices  $x_1, \dots, x_r$ . Thus

$$\sum_{i=1}^r \deg(x_i) \leq n(r-1) - |X|. \tag{2.9}$$

Combining (2.8) with (2.9), we conclude that

$$|X| \leq 5r^2\alpha n.$$

For the upper bound on the sizes of the sets  $X_i$  we get

$$|X_i| \leq n - \sum_{\substack{1 \leq j \leq r \\ j \neq i}} |X_j| \leq n - \frac{r-1}{r}n + 3r\sqrt{\alpha n} = \frac{n}{r} + 3r\sqrt{\alpha n}. \tag{2.10}$$

$\square$

We now bound the number of non-edges between  $X_1, \dots, X_r$ .

**Lemma 2.5.** *We have*

$$\sum_{1 \leq i < j \leq r} e^c(X_i, X_j) \leq t + e(X, X^c) + |X|^2 - \left(1 - \frac{1}{r}\right)n|X| + r.$$

**Proof.**

$$\begin{aligned} \frac{n^2}{2} \left(1 - \frac{1}{r}\right) - \frac{r}{2} - t &\leq e(G) \\ &= e(X, X^c) + e(X) + \sum_{1 \leq i < j \leq r} e(X_i, X_j) \\ &\leq e(X, X^c) + \frac{|X|^2}{2} + \left(1 - \frac{1}{r}\right) \left(\frac{(n - |X|)^2}{2}\right) - \sum_{1 \leq i < j \leq r} e^c(X_i, X_j). \end{aligned} \tag{2.11}$$

This gives the statement of the lemma. □

Let

$$\bar{X} = \left\{ v \in X \mid \deg_{X_1 \cup \dots \cup X_r}(v) \geq \frac{r-2}{r}n + 3\alpha^{1/3}n \right\} \quad \text{and} \quad \hat{X} := X \setminus \bar{X}.$$

Let  $d \in [0, 1]$  such that  $|\bar{X}| = d|X|$ . Further, let  $k \in [0, 5r^2]$  such that  $|X| = k\alpha n$ . Now we shall further develop the upper bound from Lemma 2.5.

**Lemma 2.6.** *We have*

$$\sum_{1 \leq i < j \leq r} e^c(X_i, X_j) \leq 20r^2\alpha^{4/3}n^2 + \left(1 - (1-d)\frac{1}{r}k\right)\alpha n^2.$$

**Proof.** By Lemma 2.5,

$$\begin{aligned} \sum_{1 \leq i < j \leq r} e^c(X_i, X_j) &\leq t + e(X, X^c) + |X|^2 - \left(1 - \frac{1}{r}\right)n|X| + r \\ &\leq t + d|X|\Delta(G) + (1-d)|X|\left(\frac{r-2}{r}n + 3\alpha^{1/3}n\right) + |X|^2 - \left(1 - \frac{1}{r}\right)n|X| + r \\ &\leq t + d|X|\left(n\frac{r-1}{r} + \frac{5}{2}\sqrt{\alpha n}\right) + (1-d)|X|\left(\frac{r-2}{r}n + 3\alpha^{1/3}n\right) \\ &\quad + |X|^2 - \left(1 - \frac{1}{r}\right)n|X| + r \\ &\leq \frac{5}{2}d|X|\sqrt{\alpha n} + 3(1-d)|X|\alpha^{1/3}n + |X|^2 + t + n|X|\frac{d-1}{r} + r \\ &\leq \frac{5}{2}k\alpha^{3/2}n^2 + 3k\alpha^{4/3}n^2 + |X|^2 + \left(1 - (1-d)\frac{1}{r}k\right)\alpha n^2 + r \\ &\leq \frac{25}{2}r^2\alpha^{3/2}n^2 + 15r^2\alpha^{4/3}n^2 + 25r^4\alpha^2n^2 + \left(1 - (1-d)\frac{1}{r}k\right)\alpha n^2 + r \\ &\leq 20r^2\alpha^{4/3}n^2 + \left(1 - (1-d)\frac{1}{r}k\right)\alpha n^2. \end{aligned} \tag{2.12}$$

□

Let

$$C := 20r^2\alpha^{4/3} + \left(1 - (1-d)\frac{1}{r}k\right)\alpha. \tag{2.13}$$

For every vertex  $u \in X$  there is no  $K_r$  in  $N_{X_1}(u) \cup \dots \cup N_{X_r}(u)$ . Thus, by applying Theorem 2.3 and Lemma 2.6, we get

$$\min_{i \neq j} |N_{X_i}(u)| |N_{X_j}(u)| \leq \sum_{1 \leq i < j \leq r} e^c(X_i, X_j) \leq Cn^2. \tag{2.14}$$

Bound (2.14) implies in particular that every vertex  $u \in X$  has degree at most  $\sqrt{Cn}$  to one of the sets  $X_1, \dots, X_r$ , that is,

$$\min_i |N_{X_i}(u)| \leq \sqrt{Cn}. \tag{2.15}$$

Therefore we can partition  $\hat{X} = A_1 \cup \dots \cup A_r$  such that every vertex  $u \in A_i$  has at most  $\sqrt{Cn}$  neighbours in  $X_i$ .

By the following calculation, for every vertex  $u \in \bar{X}$  the second smallest neighbourhood to the sets  $X_i$  has size at least  $\alpha^{1/3}n$ :

$$\min_{i \neq j} |N_{X_i}(u)| + |N_{X_j}(u)| \geq \frac{r-2}{r}n + 3\alpha^{1/3}n - (r-2) \left( \frac{n}{r} + 3r\sqrt{\alpha}n \right) \geq 2\alpha^{1/3}n, \tag{2.16}$$

where we used the definition of  $\bar{X}$  and Lemma 2.4. Combining the lower bound on the second smallest neighbourhood with (2.14), we can conclude that for every  $u \in \bar{X}$

$$\min_i |N_{X_i}(u)| \leq \frac{C}{\alpha^{1/3}}n. \tag{2.17}$$

Hence we can partition  $\bar{X} = B_1 \cup \dots \cup B_r$  such that every vertex  $u \in B_i$  has at most  $C\alpha^{-1/3}n$  neighbours in  $X_i$ . Consider the partition  $A_1 \cup B_1 \cup X_1, A_2 \cup B_2 \cup X_2, \dots, A_r \cup B_r \cup X_r$ . By removing all edges inside the classes, we end up with an  $r$ -partite graph. We have to remove at most

$$\begin{aligned} e(X) + d|X| \frac{C}{\alpha^{1/3}}n + (1-d)|X|\sqrt{Cn} &\leq 6r^2\alpha^{5/3}n^2 + (1-d)k\sqrt{C\alpha}n^2 \\ &\leq 6r^2\alpha^{5/3}n^2 + (1-d)k \left( \sqrt{20r^2\alpha^{4/3}} + \sqrt{\left(1 - (1-d)\frac{1}{r}k\right)\alpha} \right) \alpha n^2 \\ &\leq 6r^2\alpha^{5/3}n^2 + 5r^2\sqrt{20r^2\alpha^{4/3}}\alpha n^2 + (1-d)k\sqrt{\left(1 - (1-d)\frac{1}{r}k\right)\alpha} \alpha n^2 \\ &\leq 6r^2\alpha^{5/3}n^2 + 5\sqrt{20}r^3\alpha^{5/3}n^2 + \frac{2r}{3\sqrt{3}}\alpha^{3/2}n^2 \\ &\leq \left( \frac{2r}{3\sqrt{3}} + 30r^3\alpha^{1/6} \right) \alpha^{3/2}n^2 \end{aligned} \tag{2.18}$$

edges. We have used (2.15), (2.17) and the fact that

$$(1-d)k\sqrt{1 - (1-d)\frac{k}{r}} \leq \frac{2r}{3\sqrt{3}},$$

which can be seen by setting  $z = (1-d)k$  and finding the maximum of  $f(z) := z\sqrt{1-z/r}$ , which is obtained at  $z = 2r/3$ .

**3. Sharpness example**

In this section we will prove Theorem 1.4, i.e. that the leading term from Theorem 1.3 is best possible.






Figure 1. The graph  $G$ .

**Proof of Theorem 1.4.** Let  $G$  be the graph with vertex set  $V(G) = A \cup X \cup B \cup C \cup D \cup X_1 \cdots \cup X_{r-2}$ , where all classes  $A, X, B, C, D, X_1, \dots, X_{r-2}$  form independent sets;  $A, X, B, C, D$  form a complete blow-up of a  $C_5$ , where the classes are named in cyclic order; and for each  $1 \leq i \leq r - 2$ , every vertex from  $X_i$  is incident to all vertices from  $V(G) \setminus X_i$ . See Figure 1 for an illustration of  $G$ .

The sizes of the classes are

$$|X| = \frac{2r}{3}\alpha n, \quad |A| = |B| = \sqrt{\frac{\alpha}{3}}n, \quad |C| = |D| = \frac{1 - (2r/3)\alpha}{r}n - \sqrt{\frac{\alpha}{3}}n, \quad |X_i| = \frac{1 - (2r/3)\alpha}{r}n.$$

The smallest class is  $X$  and the second smallest are  $A$  and  $B$ . By deleting all edges between  $X$  and  $A$  ( $|X||A| = (2r/(3\sqrt{3}))\alpha^{3/2}n^2$ ), we get an  $r$ -partite graph. Since the classes  $A$  and  $X$  are the two smallest class sizes, the smallest canonical cut is of size  $(2r/(3\sqrt{3}))\alpha^{3/2}n^2$ . A result by Erdős, Györi and Simonovits [7, Theorem 7] states that there is a canonical ‘edge deletion’ achieving the minimum of  $D_r(G)$ . Hence

$$D_r(G) \geq \frac{2r}{3\sqrt{3}}\alpha^{3/2}n^2.$$

Let us now count the number of edges of  $G$ . The number of edges incident to  $X$  is

$$\begin{aligned} e(X, X^c) &= \left(\frac{2r}{3}\alpha\right)\left(2\sqrt{\frac{\alpha}{3}}n\right)^2 + \left(\frac{2r}{3}\alpha\right)\left(\frac{1 - (2r/3)\alpha}{r}(r - 2)\right)n^2 \\ &= \left(\frac{2}{3}(r - 2)\alpha + \frac{4r}{3\sqrt{3}}\alpha^{3/2} - \frac{4r(r - 2)}{9}\alpha^2\right)n^2. \end{aligned} \tag{3.1}$$

Using that  $|A| + |C| = |B| + |D| = |X_1|$ , we have that the number of edges inside  $A \cup B \cup C \cup D \cup X_1 \cup \dots \cup X_{r-2}$  is

$$\begin{aligned} e(X^c) &= |X_1|^2 \binom{r}{2} - |A||B| \\ &= \left(\frac{1 - (2r/3)\alpha}{r}n\right)^2 \binom{r}{2} - \frac{1}{3}\alpha n^2 \\ &= \frac{1}{r^2} \binom{r}{2} n^2 - \frac{4r}{3} \frac{1}{r^2} \alpha \binom{r}{2} n^2 + \frac{4}{9} \alpha^2 \binom{r}{2} n^2 - \frac{1}{3} \alpha n^2 \\ &= \left(1 - \frac{1}{r}\right) \frac{n^2}{2} - \frac{2}{3}(r - 1)\alpha n^2 - \frac{1}{3}\alpha n^2 + \frac{4}{9}\alpha^2 \binom{r}{2} n^2. \end{aligned} \tag{3.2}$$

Thus the number of edges of  $G$  is

$$\begin{aligned} e(G) &= e(X^c) + e(X, X^c) \\ &= \left(1 - \frac{1}{r}\right) \frac{n^2}{2} - \alpha n^2 + \frac{4r}{3\sqrt{3}} \alpha^{3/2} n^2 - \frac{2r(r-3)}{9} \alpha^2 n^2 \\ &\geq \text{ex}(n, K_{r+1}) - \alpha n^2 + \frac{4r}{3\sqrt{3}} \alpha^{3/2} n^2 - \frac{2r(r-3)}{9} \alpha^2 n^2, \end{aligned} \quad (3.3)$$

where we applied Turán's theorem in the last step.  $\square$

## Acknowledgements

We would like to thank an anonymous referee for useful comments.

## References

- [1] Amin, K., Faudree, J., Gould, R. J. and Sidorowicz, E. (2013) On the non- $(p-1)$ -partite  $K_p$ -free graphs. *Discuss. Math. Graph Theory* **33** 9–23.
- [2] Andrásfai, B., Erdős, P. and Sós, V. T. (1974) On the connection between chromatic number, maximal clique and minimal degree of a graph. *Discrete Math.* **8** 205–218.
- [3] Balogh, J., Morris, R., Samotij, W. and Warnke, L. (2016) The typical structure of sparse  $K_{r+1}$ -free graphs. *Trans. Amer. Math. Soc.* **368** 6439–6485.
- [4] Brouwer, A. E. (1981) *Some Lotto Numbers From an Extension of Turán's Theorem*, Vol. 152 of Afdeling Zuivere Wiskunde [Department of Pure Mathematics]. Mathematisch Centrum, Amsterdam.
- [5] Erdős, P. (1970) On the graph theorem of Turán. *Mat. Lapok* **21** 249–251.
- [6] Erdős, P., Faudree, R., Pach, J. and Spencer, J. (1988) How to make a graph bipartite. *J. Combin. Theory Ser. B* **45** 86–98.
- [7] Erdős, P., Győri, E. and Simonovits, M. (1992) How many edges should be deleted to make a triangle-free graph bipartite? In *Sets, Graphs and Numbers (Budapest, 1991)*, Vol. 60 of Colloquia Mathematica Societatis János Bolyai, pp. 239–263. North-Holland.
- [8] Erdős, P. and Simonovits, M. (1966) A limit theorem in graph theory. *Studia Sci. Math. Hungar* **1** 51–57.
- [9] Füredi, Z. (2015) A proof of the stability of extremal graphs, Simonovits' stability from Szemerédi's regularity. *J. Combin. Theory Ser. B* **115** 66–71.
- [10] Hanson, D. and Toft, B. (1991)  $k$ -saturated graphs of chromatic number at least  $k$ . *Ars Combin.* **31** 159–164.
- [11] Hu, P., Lidický, B., Martins, T., Norin, S. and Volec, J. Large multipartite subgraphs in  $H$ -free graphs. Manuscript.
- [12] Kang, M. and Pikhurko, O. (2005) Maximum  $K_{r+1}$ -free graphs which are not  $r$ -partite. *Mat. Stud.* **24** 12–20.
- [13] Korándi, D., Roberts, A. and Scott, A. (2020) Exact stability for Turán's theorem. [arXiv:2004.10685](https://arxiv.org/abs/2004.10685)
- [14] Nikiforov, V. (2011) Some new results in extremal graph theory. In *Surveys in Combinatorics 2011*, Vol. 392 of London Mathematical Society Lecture Note Series, pp. 141–181. Cambridge University Press.
- [15] Roberts, A. and Scott, A. (2018) Stability results for graphs with a critical edge. [arXiv:1610.08389](https://arxiv.org/abs/1610.08389). <https://doi.org/10.1016/j.ejc.2018.07.004>
- [16] Sudakov, B. (2007) Making a  $K_4$ -free graph bipartite. *Combinatorica* **27** 509–518.
- [17] Turán, P. (1941) Eine Extremalaufgabe aus der Graphentheorie. *Mat. Fiz. Lapok* **48** 436–452.
- [18] Tyomkyn, M. and Uzzell, A. J. (2015) Strong Turán stability. *Electron. J. Combin.* **22** P3-9.

**Cite this article:** Balogh J, Clemen FC, Lavrov M, Lidický B and Pfender F (2020). Making  $K_{r+1}$ -free graphs  $r$ -partite. *Combinatorics, Probability and Computing*. <https://doi.org/10.1017/S0963548320000590>