

P0110

Efficacy of Ziprasidone in hospitalized patients with severe mania

M.H. Allen¹, F.S. Mandel². ¹University of Colorado Health Sciences Center, Denver, CO, USA ²Pfizer Inc., New York, NY, USA

Background: Ziprasidone is effective and well tolerated in patients with bipolar mania, as demonstrated by 2 pivotal, placebo-controlled trials.

Methods: We evaluated pooled data for subjects with baseline Mania Rating Scale (MRS) scores in the highest 10% of the severity distribution at study baseline. All subjects in this post hoc analysis had an MRS score ≥ 36 , indicating marked severity of mania. Changes in MRS scores from baseline to days 2, 4, 7, 14, and 21 were analyzed using a Cochran-Mantel-Haenszel test to determine response, defined as a decrease in MRS score of $\geq 50\%$ from baseline, and remission, defined as an MRS score ≤ 10 .

Results: The proportion of subjects achieving response was significantly greater for subjects receiving ziprasidone compared with placebo from day 7 ($p = 0.03$) through to study end point ($p < 0.001$). The proportion of subjects achieving remission was significantly greater than placebo ($p = 0.01$) at study end point. Few subjects who received placebo achieved response, and none achieved remission at any timepoint during the study.

Conclusions: Ziprasidone is effective in the treatment of patients with severe manic symptoms. This conclusion is underscored by the relatively large drug-placebo differences observed in this patient subsample.

P0111

Meta-analysis of magnetic resonance imaging studies of the corpus callosum in bipolar disorder

D. Arnone¹, A.M. McIntosh², P. Chandra³, K.P. Ebmeier¹. ¹University Department of Psychiatry, Oxford, UK ²University of Edinburgh, Royal Edinburgh Hospital, Edinburgh, UK ³SUNY Downstate Medical Centre, New York, NY, USA

Objective: The corpus callosum plays a pivotal role in inter-hemispheric transfer and integration of information and is a relatively understudied structure in bipolar disorder. Magnetic resonance studies have reported callosal abnormalities in this condition but findings have been inconsistent. Structural changes affecting the CC may underlie functional abnormalities in bipolar disorder and could contribute to, or explain the pathophysiology of the condition.

Method: A systematic review was carried out to identify, appraise and summarise magnetic resonance studies which compared callosal areas in bipolar disorder with an unrelated control group. The findings were then synthesised using random effects meta-analysis. Consideration was given to a number of variables to explain heterogeneity.

Results: Five case-control studies were identified. Bipolar patients showed reduced callosal areas and the effect size revealed a statistically significant effect: (- 0.52, 95% CI = -0.82, - 0.21). Bipolar patients showed reduced callosal areas in comparison to healthy volunteers. There was no statistically significant heterogeneity in the studies included ($I^2 = 0.15$, $P = 0.3$) and no evidence of publication bias was detected (Egger test P value = 0.5). No significant effects were noted in meta-regression analysis with reference to age ($P = 0.9$), gender expressed as percentage of male subjects ($P = 0.6$), and year of publication ($P = 0.4$).

Conclusions: Findings from this study indicate that callosal areas are reduced in bipolar disorder and suggest that a failure to integrate

information across the hemispheres may contribute to the pathophysiology of the disorder.

P0112

Association between G72/30 haplotypes and bipolar disorder in a Swedish sample

L. Backlund¹, P. Nikamo², L. Rosvall², L. Frisen³, M. Landen¹, J. Ladanayi⁴, L. Träskman-Bendz⁵, H. Ågren¹, M. Schalling², U. Ösby^{1,3}. ¹Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden ²Neurogenetics, Karolinska Institutet, Stockholm, Sweden ³Psychiatry, Danderyds Hospital, Stockholm, Sweden ⁴Psychiatry South Area Stockholm, Stockholm, Sweden ⁵Division of Psychiatry, Department of Clin. Neurosciences, University Hospital, Lund, Sweden

Introduction: Bipolar Disorder type 1 (BP-1) is a severe and common psychiatric disorder with a strong genetic contribution to increased disease risk. Heritability is around 60% and at least a dozen different genes seems to be involved in the etiology. The DAOA-G72/G30 locus on chromosome 13p22-34 has been associated with BP-1 in at least six independent datasets, although there are different SNPs and haplotypes associated with increased disease risk in these studies.

Methods: 276 Swedish patients with BP-1, diagnosed according to DSM-IV, and 940 controls were analyzed with 36 SNPs from eight previously reported candidate genes for bipolar disorder (DAOA-G72/G30, P2RX7, COMT, BDNF, CAMKK2, GRK3, SLC6A4 and S100A10). Genotyping was carried out with Applied Biosystems TaqMan[®] using ABI PRISM[®] 7900HT Sequence Detecting System. Statistical analysis was performed with the Unphased program.

Results: None of the SNPs showed a significant association by itself. However, three SNPs in different haplotype blocks in the DAOA-G72/G30 genes were together associated to bipolar disorder ($p=0.0009$). The association remained when different SNPs in each haplotype block were analyzed.

Conclusions: This is the first study of the G72/30 gene in a homogeneous Swedish BP-1 sample. We found a strong association, which required SNPs from three different haplotype blocks. Thus, the association between the DAOA-G72/G30 genes and Swedish patients with BP-1 disorder was more complex than previous findings.

P0113

Relationship between psychological well-being and general health

A. Asghar Bayani. Islamic Azad University, Azadshahr, Iran

Aim: Traditionally, health researcher has focused on negative states, their determinants and consequences. Until relatively recently, few studies have been conducted on subjective moods or feelings of well-being and their determinants. The purpose of the present study was to examine the relationship between general health and psychological well-being.

Method: Participants were 145 undergraduate students of Islamic Azad university of Azadshahr. The mean age of the participants was 24.7 years ($SD = 4.52$) and ages ranged from 18 to 40 years old. They were 49 men and 96 women.

Measures: All participants completed a questionnaire booklet containing two self-report measures: The Scales of Psychological Well-Being (Ryff, 1995) and General Health Questionnaire (GHQ-28).

Results: The results of the present study demonstrate that: Correlation between psychological well-being and student's general health is meaningful and positive.

P0114

Quetiapine in the maintenance treatment of bipolar I disorder: Combined data from two long-term phase III studies

M. Brecher¹, H. Anderssen², B. Paulsson². ¹ AstraZeneca Pharmaceuticals LP, Wilmington, DE, USA ² AstraZeneca Pharmaceuticals, Sodertalje, Sweden

Background and Aims: Combined data are presented from two long-term, phase III studies (D1447C00126; D1447C00127) that examined the efficacy and safety of quetiapine in combination with lithium/divalproex in the prevention of mood events (manic, mixed, or depressed) in bipolar I disorder.

Methods: During a stabilization phase (12–36 weeks), patients received quetiapine (400–800 mg/day; open-label; flexible, divided doses) with lithium or divalproex (target serum concentrations 0.5–1.2 mEq/L and 50–125 µg/mL). Thereafter, patients were randomized to double-blind treatment with quetiapine (400–800 mg/day)+lithium/divalproex or placebo+lithium/divalproex for up to 104 weeks. The primary endpoint was time to the first mood event, defined by medication initiation, hospitalization for a mood event, YMRS or MADRS scores ≥ 20 at 2 consecutive assessments, or discontinuation due to a mood event.

Results: In total, 3414 patients entered the stabilization phase and 1326 were randomized and received ≥ 1 dose of study medication. Rates of recurrence were 19.3% vs 50.4% for quetiapine and placebo groups, respectively. The risk of recurrence of a mood event was significantly reduced in the quetiapine group relative to the placebo group (HR=0.30, $P < 0.0001$). HRs for the risk of recurrence of manic and depressed events were both 0.30 ($P < 0.0001$). Long-term treatment with quetiapine was generally well tolerated; safety data including glucose changes will also be presented.

Conclusions: Quetiapine in combination with lithium/divalproex is significantly more effective than lithium/divalproex alone in the prevention of mood episodes associated with bipolar I disorder, irrespective of the index episode.

Supported by funding from AstraZeneca Pharmaceuticals LP.

P0115

The using of lithium in the treatment of bipolar disorder - Changes in the time

D. Breznoscakova, E. Palova, J. Dragasek, K. Losonciová, J. Uhrin, P. Mirossay. *1st Department of Psychiatry of Medical School of University of P. J. Safarik, Kosice, Slovak Republic*

Introduction: Bipolar disorder (BD) is chronic psychiatric disorder characterized by remission and exacerbations of mood disturbances. Treatment of BD is often complicated and includes in the first step treatment with lithium, after with other "mood stabilizers", antipsychotics, antidepressants.

The aim of study:

- To assess the change of treatment with lithium over the period of time
- The number of hospitalizations in the patients with BD and differences in using of lithium, age and type of first episode

Patients and Methods:

- Retrospective survey of in-patient's files (1997 - 2007) with typical limitations for retrospective case survey:
- BD, n = 125 (67% women with BD, DSM-IV)

- Number of hospitalizations, average number hospitalizations/year, type of treatment, using of lithium

Results:

- The first episode was depressive in 62% of patients and average number of episodes: 6,2
- There was a steady decrease in use of mood stabilizers medication over the time (68% vs 82%)
- We observed increase in prescription of atypical antipsychotics (in 80% of patients; last 6 years = 93%)
- The using of lithium was in 52% of patients (in 41% patients during first hospitalization) with average dose 1250mg of lithium and average surface of lithium in blood: 0,64mmol/l

Conclusion: We find out trend in using of lithium in 52% patients with BD last time again with average dose 1250mg per day and average surface of lithium in blood 0,64mmol/l and increasing of prescription of atypical antipsychotics (last 6 years = 93%).

P0116

Self appreciation and affective temperaments in psychiatric nurses

R.C. Cordeiro^{1,2,3}. ¹ Instituto Politecnico de Portalegre, Portalegre, Portugal ² Escola Superior de Saude de Portalegre, Portalegre, Portugal ³ Centro de Estudos de Saúde Mental, Portalegre, Portugal

An analysis about the Self Appreciation and the Affective Temperament of Nurses in Psychiatric and Mental Health Services, have a crucial importance, because this fact will influence many phenomena's in nursing practice, like the capacity of develop relationships and create resistance to physical and psychological illness. A total of n=47 nurses, of both sexes, with average age of 38,57 years, working in Mental Health and Psychiatry Departments, in three portuguese Hospitals, were inquired through a Questionnaire of direct application. In the Questionnaire, were introduced measures like: TEMPS-A Scale (Akiskal, 1998), translated into Portuguese, by Figueira and Severino (1999) and Scale of Self Personal Appreciation Scale (Ribeiro, 2006). The obtained results indicate that the population, have, majority, a Hyperthymic Temperament. It was verified that the women presents a higher Self Personal Appreciation and his Self Appreciation is statistically related to the Cyclothymic and Anxious temperaments. The results also show that exist an association between the marital status and the Irritable Temperament, and between the Time of Service and the Cyclothymic and Hyperthymic Temperaments.

Affective temperament of nurses seems to be a good predictor of leadership capacity in violence situation at psychiatric services.

P0117

Overweight and obesity in bipolar disorder

L.C. Castro, V.G. Pais, J. Pereira. *Serviço de Matosinhos, Magalhaes Lemos Hospital, Porto, Portugal*

Aims of the study: To study Body Mass Index (BMI) and Blood Pressure (BP) in bipolar patients.

Methods: Retrospective study in bipolar inpatients of an acute psychiatric Portuguese department during a two years period. Of the 60 bipolar inpatients, 31 were selected. The patients without height or weight data were excluded.

Results: The mean age was 48.5 years. 66.7% of the patients were women. The most frequent pharmacological association (13.3%) was sodium valproate and olanzapine. Using INÉs (National Statistic Institute, 2007) criteria, 25.8% of these patients were overweight and