




Research Article

Isotopic and provenance analysis of Neolithic and Bronze Age shell disc beads from Ban Non Wat, north-east Thailand

Jelmer W. Eerkens^{1,*} , Charles Higham²  & Howard J. Spero³

¹ Department of Anthropology, University of California Davis, USA

² Department of Anthropology and Archaeology, University of Otago, New Zealand

³ Department of Earth and Planetary Sciences, University of California Davis, USA

* Author for correspondence ✉ jweerkens@ucdavis.edu



Small, disc-shaped shell beads are recorded as mortuary offerings in many Neolithic and Bronze Age burials in Southeast Asia. Yet the provenance of these artefacts is often obscure, as production processes involve the removal of diagnostic morphological features, negating taxonomic classification. Here, the authors report on the combined isotopic and morphological analysis of a subset of shell beads from the site of Ban Non Wat in north-east Thailand. In addition to identifying freshwater sources for nearly all the beads, the results suggest the presence of multiple shell production centres—each with access to distinct aqueous environments—and widespread exchange in the Bronze Age.

Keywords: Southeast Asia, Neolithic, Bronze Age, stable isotope analysis, provenance analysis, shell beads

Introduction

Charting the movement of material commodities between ancient societies continues to provide important frameworks for archaeologists seeking to understand the nature of archaeo-economies, especially the rise of sociopolitical complexity (Brumfiel & Earle 1987; Oka & Kusimba 2008; Abraham *et al.* 2016). Materials such as obsidian, glass and pottery are a mainstay within this field of study, where chemical signatures can reveal material sources and trace pathways to sites of consumption and deposition. Shell beads and ornaments are also common finds at many archaeological sites and, while taxonomic classification may sometimes identify the general bodies of water from which the shells were procured

Received: 5 May 2023; Revised: 25 August 2023; Accepted: 13 October 2023

© The Author(s), 2024. Published by Cambridge University Press on behalf of Antiquity Publications Ltd. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

(i.e. coastal versus freshwater), less consideration has been given to the geochemical provenancing of shell artefacts (though see Shackleton & Renfrew 1970; Shackleton & Elderfield 1990; Vanhaeren *et al.* 2004; Eerkens *et al.* 2010; Peacock *et al.* 2010; Bajnóczi *et al.* 2013).

Here, we report on the chemical analysis of a small sample of beads from Ban Non Wat, a settlement in the upper catchment of the Mun River in north-east Thailand (Figure 1), allowing an environmental insight into the production of shell disc beads in this area from *c.* 1750–450 BC. The inclusion of shell beads within burials was a widespread practice during the Neolithic and Bronze Age of Southeast Asia that reached an early apogee at Neolithic Khok Phanom Di where a woman was interred wearing approximately 121 000 shell specimens (Higham & Bannanurag 1990). While stone and glass beads, especially from the Iron Age, have been the subject of previous regional research, including application of provenance-based approaches (Francis 1991; Pilditch 1992; Klysubun *et al.* 2011; Bellina 2014; Carter 2015; Carter *et al.* 2021; Seman *et al.* 2021), shell beads have not figured prominently (Ciarla *et al.* 2017). Larger shell ornaments at Ban Non Wat include hundreds of *Trochus* and *Tridacna* bangles that are unquestionably marine in origin; a similar source is also generally assigned to the smaller disc beads but these have never been provenanced. Unfortunately, the morphological properties that might identify the original species of a shell are removed during the production of small disc beads. Thus, the taxonomic classification and indeed the number of species used to produce these items remains unknown. The present study aims to help identify if marine or freshwater shells were used and through this to examine the organisation of shell bead production, in order to illuminate possible sources and production centres, since no evidence for the manufacture of shell ornaments was found during the extensive excavations at Ban Non Wat.

Ban Non Wat

Ban Non Wat is a moated prehistoric settlement located in the upper Mun Valley, north-east Thailand. This is a strategic location, lying on a choke point for exchange to the east of a pass over the Phetchabun Upland, that divides the Khorat Plateau from the broad plains of the Chao Phraya River (Figure 1). The Mun River is a natural conduit for exchange as it flows east to its confluence with the Mekong. Many exotic items, including marine shell jewellery, copper-base artefacts and ornaments of marble, glass, silver, gold, carnelian and agate, have been recovered from Ban Non Wat over 10 seasons of excavations by a joint University of Otago and Fine Arts Department of Thailand research programme that began in 2002 (Higham & Kijngam 2009, 2010, 2012). Thirteen prehistoric periods of occupation have been identified through the analysis of mortuary practices and a programme of radiocarbon dating (Higham 2024). The site was initially occupied by hunter-gatherers *c.* 17 000 BC, though this occupation phase is not further considered here. A second period of hunter-gatherer occupation is contemporary with the first Neolithic settlement (beginning *c.* 1750 BC), during which interments of both cultures shared the same burial ground. Hunter-gatherer burials are distinguished from Neolithic contemporaries by the flexed position of the dead, a widespread characteristic of hunter-gatherer mortuary treatment in Southeast Asia. A second Neolithic occupation (not included in this study) was succeeded *c.* 1050 BC by the first of five Bronze Age phases, which conclude with an Iron Age occupation *c.* 450 BC. From the initial Neolithic settlement until the abandonment of the site (*c.* AD 400), there is a




Figure 1. Map showing the location of Ban Non Wat and places mentioned in the text (map by C.F.W. Higham, using GeoMapApp (www.geomapapp.org), after Ryan et al. 2009).

continuity of culture with no evidence for an incoming population (Higham & Higham 2009; Higham & Thosarat 2020).

From the combined mortuary record of almost 700 burials, 13 represent indigenous hunter-gatherers, while two Neolithic and five Bronze Age burial phases incorporate around 270 inhumations within which the dead were commonly interred with small disc-shaped shell beads (Table 1). The number of these beads varies markedly, from a solitary specimen in 31 Neolithic 1 graves to 83 055 examples in the 32 Bronze Age 2 burials. Based on their positioning within burials, beads were often strung and worn as belts and necklaces. Richly furnished burials also often included beads adhering to the skulls that might represent the presence of a shroud or head covering. A surge in the number of beads with Bronze Ages 2 and 3 is matched by an increase in the number of other mortuary offerings, including early copper-base axes, bells, anklets, chisels and awls, marine shell bangles and earrings, and marble ornaments. The presence of these offerings is interpreted as symbolic of a rapid but ephemeral rise of social aggrandisers with access to exchange networks (Higham 2024). Successful identification of the source of the shell beads, however, has been elusive; it is assumed that, like the bangles, they were probably marine.

The total number of beads from hunter-gatherer and Neolithic burials is relatively low ($n = 489$, Table 1). From Bronze Age 1 to Bronze Age 3, the total number of beads in burials increases dramatically, before decreasing again in Bronze Age 4 and 5. Both the percentage of burials that include beads as well as the average number of beads across all individuals in a time period peak in Bronze Age 3, where every individual has at least some disc beads and the average number of beads per individual tops 6000. Clearly, beads were of great importance in mortuary contexts during Bronze Age 2 and 3.

Table 1. Number of shell disc beads in each mortuary phase. Flexed: hunter-gatherer; Neo: Neolithic; BA: Bronze Age.

Phase	Approximate age range (cal BC)	Number of burials	Number of beads	Number of burials with beads	Percentage of burials with beads	Average number of beads per burial
Flexed	1750–1050	13	478	2	15.4	18
Neo 1	1650–1250	31	1	1	3.2	0.03
Neo 2	1250–1050	38	10	3	7.9	0.3
BA 1	1050–1000	7	3291	3	42.8	470
BA 2	1000–900	32	83 055	26	81.3	2596
BA 3	900–800	13	78 171	13	100.0	6013
BA 4	800–700	20	4335	7	35.0	216
BA 5	700–420	114	2785	66	57.9	24

The shell disc beads from Ban Non Wat are broadly similar in shape and vary categorically, rather than continuously, by size (e.g. Figure 2A shows large, medium and small beads). Those associated with hunter-gatherer and Neolithic individuals are often chipped and rough around the edges (see Figure 2B), occasionally producing polygonal or even rectangular outlines, whereas Bronze Age beads are ground smooth along the edges and more circular (Figure 2A).

Isotopic approach

Many molluscs precipitate calcium carbonate (CaCO_3) in sequential rings to form a protective exoskeleton by taking in calcium (Ca) and carbonate (CO_3^{2-}) or bicarbonate (HCO_3^-) from the waters in which they grow. The precipitated calcite or aragonite often does not alter after formation. Thus, the carbon and oxygen within the crystal structure of a shell records diachronic information about the growing environment of the organism via the stable isotopic ratios of these two elements ($^{13}\text{C}/^{12}\text{C}$ and $^{18}\text{O}/^{16}\text{O}$) (Urey *et al.* 1948; Mook & Vogel 1968). As the carbon and oxygen geochemistry of aqueous environments varies over both time and space, stable isotopic analyses of ancient shells can help archaeologists reconstruct aspects of the environment in which the organisms lived, such as temperature, elevation and the season shells were harvested (Deith 1895; Shackleton 1973; Kennett & Voorhies 1996; Stephens *et al.* 2008; Pérez *et al.* 2020; Skippington *et al.* 2021). By extension, such information may provide key details about the lives of people in the past.

Empirical data show that the oxygen isotope ratio ($\delta^{18}\text{O}$) of biogenic carbonate from freshwater shells is generally in equilibrium with the oxygen isotopic composition of ambient water ($\delta^{18}\text{O}_w$) (Grossman & Ku 1986; Dettman *et al.* 1999; Versteegh *et al.* 2010). The $\delta^{18}\text{O}_w$ in freshwater systems is strongly correlated with patterns of precipitation where $\delta^{18}\text{O}_w$ decreases with distance from the source of water (e.g. the ocean) and elevation. This trend is produced because water molecules containing the heavier ^{18}O isotope condense and precipitate at a higher rate than those containing the lighter ^{16}O isotope. With time, the $^{18}\text{O}/^{16}\text{O}$ ratio of water vapour in an air mass becomes depleted in ^{18}O relative to ^{16}O thereby



Figure 2. Examples of beads associated with A) Bronze Age burial 90 (n = 9; note different sizes) and B) hunter-gatherer burial 461 (n = 4; note rough and chipped outer margins) (figure by authors).

altering the oxygen isotope ratio of rainfall as a storm tracks further inland (Kendall & Coplen 2001). Local $\delta^{18}\text{O}_w$ can also vary as a function of humidity, air temperature and altitude, due to this isotopic rainout effect. In addition, water temperature has a predictable effect on the $\delta^{18}\text{O}$ of shell CaCO_3 ($\delta^{18}\text{O}_c$), where $\delta^{18}\text{O}_c$ decreases as temperature increases and vice versa (Grossman & Ku 1986). Together, these fractionation effects can lead to significant geographic and spatial variation in $\delta^{18}\text{O}_w$ and $\delta^{18}\text{O}_c$.

By contrast, the $^{13}\text{C}/^{12}\text{C}$ ratio of a shell ($\delta^{13}\text{C}_c$) is strongly correlated with the $^{13}\text{C}/^{12}\text{C}$ ratio of dissolved inorganic carbon (DIC) in water. $\delta^{13}\text{C}_{\text{DIC}}$ is largely governed by plant and algae photosynthesis where plants preferentially utilise dissolved carbon dioxide containing the ^{12}C isotope during growth, thereby increasing the $\delta^{13}\text{C}_{\text{DIC}}$ of the water. The decay of organic matter has the opposite effect on $\delta^{13}\text{C}_{\text{DIC}}$ because respiration releases ^{13}C -depleted carbon dioxide. In addition, water temperature and/or metabolic effects within an organism

Table 2. Beads sampled for stable isotope analysis from Ban Non Wat. Flexed: hunter-gatherer; Neo: Neolithic; BA: Bronze Age.

Time period	Approximate age range (cal BC)	Number of burials sampled	Burial numbers sampled	Number of beads	Number of isotopic samples
Flexed	1750–1050	2	438, 461	6	31
Neo 1	1650–1250	1	250	3	9
BA 1	1050–1000	2	446, 569	6	15
BA 2	1000–900	2	90, 105	6	22
BA 3A	900–800	2	260, 263	6	19
BA 4D	800–700	2	392, 430	6	20
BA 5	700–420	1	227	3	8
Total		12		36	124

can have a small but measurable effect on $\delta^{13}\text{C}_c$ (Grossman & Ku 1986; Dettman *et al.* 1999; Goewert *et al.* 2007). Aside from metabolic effects, these other factors may vary in their intensity or magnitude across different aquatic environments, for example due to differences in sunlight, water depth or flow rates. These environmental differences may lead to significant spatial variation in $\delta^{13}\text{C}_c$ across the landscape.

Provided variation over time within a particular body of water is smaller than that across different bodies of water, $\delta^{18}\text{O}_c$ and $\delta^{13}\text{C}_c$ can serve as markers of the original provenance of shells and their growing environment. We exploit this potential in our analysis of shell disc beads from Ban Non Wat.

Methods

Thirty-six beads associated with 12 individuals from Ban Non Wat were sampled for isotopic analysis (three beads per burial; see Table 2). Where possible (i.e. except in the case of very small beads), each specimen was sampled at multiple points. Prior to sampling, each bead was cleaned with deionised water and a toothbrush to remove any exogenous materials and left to dry. The outer surface of the bead was then lightly burred to remove the outermost layer of shell from the areas identified for sampling. Samples of carbonate were removed sequentially from the outer surface of the bead using a 0.5mm drill bit. Each sample was taken parallel to the orientation of shell growth rings to obtain estimates of water conditions at multiple points in the growth history of the organism. The linear distance between samples ranged from 1.0 to 2.0mm. Depth of sampling into the shell was restricted to the size of the drill bit, or approximately 0.5mm and powdered carbonate samples ranged from 60–100 μg in weight. Figure 3 shows a bead from burial 461 (which contained a Neolithic female) after drilling; the six areas from which isotopic samples were removed appear as lighter-coloured parallel bands. In total, 124 isotopic measurements were made from the 36 beads sampled (see Table 2), or an average of 3.4 samples per bead, with a minimum of one and a maximum of seven samples from an individual bead. It was possible to sample beads with larger diameters more intensively due to the greater surface area, while some small beads could only be sampled once.




Figure 3. Shell bead from burial 461, showing sampling strategy for isotopic analysis (figure by authors).

We additionally sampled three modern freshwater mussel shells from the Mun River near Ban Non Wat, one example each of *Hyriopsis bialata* ($n = 5$ isotopic samples), *Pilsbryconcha exilis* ($n = 6$ samples) and *Pseudodon inoscularis* ($n = 25$ samples). After removing the periostracum (the thin outer organic coating), these shells were also sampled parallel to the growth rings at 1mm intervals from the growing edge. Modern shell results are used to gain an estimate of isotopic variation within an individual shell, as well as the effects of modern

water conditions on shell isotopic composition near the site, though we recognise that modern conditions could be quite different from ancient water conditions.

Isotopic analysis was conducted at the University of California Davis on a Micromass Optima isotope ratio mass spectrometer (IRMS). Prior to analysis on the IRMS, powdered CaCO_3 samples were gently heated at 75°C *in vacuo* for 30 minutes to remove adsorbed water and subsequently reacted in 105% orthophosphoric acid at 90°C using an ISOCARB automated common acid bath system. The resulting carbon dioxide was purified through a series of cryotrap and introduced into the IRMS through a dual inlet system. Stable isotopic ratios are presented as the per mil (parts per thousand, ‰) deviation from the Pee Dee Belemnite (PDB) standard. External precision for $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values is ± 0.09 and ± 0.07 , respectively, based on multiple analyses of the calcite standards NBS-19 (limestone) and UCD-SM92 (marble).

Results

Stable isotope analysis

Figure 4 shows the isotopic results from the three modern shells. Samples were taken in the white-coloured area of each shell. The data do not display clear sinusoidal or cyclical shifts in isotopic ratios, as would be expected if our sampling covered annual shifts in water conditions, though it is probable that a full year's growth was not covered by the sampling of the *H. bialata* and *P. exilis* shells. Maximum isotopic variation within a modern shell is 3.5‰ for $\delta^{13}\text{C}$ and 4.5‰ for $\delta^{18}\text{O}$. All three shells overlap in their isotopic range for $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$, though the extent of the range of each differs between shells.

Figure 5 plots $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ for the beads coded by time period, together with an ellipse showing the range of the three modern shells. Every isotopic sample is plotted; thus, some beads are represented by multiple points while others only one (depending on how many samples were taken from a bead). The cluster of four points on the far right of Figure 5, for example, represents a single bead from burial 430 (Bronze Age 4, group D).

The three modern shells fall into the centre of the isotopic range of the prehistoric beads, suggesting that some of the beads from burials at Ban Non Wat could have been made from shells growing near the site, but that most were probably not. With the exception of the bead




Figure 4. $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ for three modern shells collected from the Mun River near Ban Non Wat (figure by authors).

from burial 430, the $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values of the Ban Non Wat shell beads fall within the range for freshwater shells (Keith *et al.* 1964). The high $\delta^{13}\text{C}$ values for the bead from burial 430 potentially indicate that this is the only shell in the whole dataset that was precipitated in marine or brackish water.

Figure 5 shows that the range in $\delta^{18}\text{O}$ (7.7‰) and especially $\delta^{13}\text{C}$ (14.9‰) within the beads is far greater than that observed in the modern shells (4.5‰ and 3.5‰, respectively). Even within a single burial, representing a short time span, values across beads have a range up to 4.8‰ in $\delta^{18}\text{O}$ and 12.8‰ in $\delta^{13}\text{C}$. Together, this suggests that the archaeological beads are drawn from a greater range of growing environments than the sample of modern shells. There is also little to no correlation between $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ within the bead data set (note that this result also holds for the modern shells). While some studies of freshwater shells show significant and positive correlation in these isotopic systems (e.g. Versteegh *et al.* 2010), others show weak or no correlation (Goewert *et al.* 2007). Lack of correlation here suggests that the two isotopic systems record independent aspects of the waters in which the shells grew in this region of Thailand. In particular, the distance to the coast (affecting mainly $\delta^{18}\text{O}$) is not correlated with the degree of plant and algal growth and/or decomposition (affecting mainly $\delta^{13}\text{C}$).

Some temporal trends are, however, apparent. Beads from hunter-gatherer and Neolithic burials fall towards the right side of the graph with generally higher $\delta^{13}\text{C}$ values, while many of the Bronze Age (and particularly the Bronze Age 5) beads fall towards the left side, with generally lower $\delta^{13}\text{C}$ values. $\delta^{18}\text{O}$ seems to be less correlated with time, though Bronze Age 2 and 3




Figure 5. $\delta^{18}\text{O}$ against $\delta^{13}\text{C}$ for samples from Ban Non Wat shell disc beads (figure by authors).

beads show more variation in $\delta^{18}\text{O}$ than the other time periods. These findings suggest two possible interpretations. First, it is possible that regional water conditions changed markedly over time. Decreasing $\delta^{13}\text{C}$ over time may indicate increasing organic load and decomposition (i.e. bacterial degradation of organics) within local river systems. This might occur, for example, if population levels were increasing regionally and more organic runoff from food processing, agricultural field fertilisation and human/animal defecation and the like were impacting local rivers. Second, decreasing $\delta^{13}\text{C}$ could represent a change in regional bead production and distribution networks. For example, Neolithic beads may have been made mostly in locations with access to faster-moving waters or areas with higher rates of aquatic plant growth, while Bronze Age beads may have been obtained mostly from locations with slower-moving water with higher organic decomposition. A shift in the importance of beads from these two environments, with different underlying water conditions, could also produce the results in Figure 5.

Shell bead production

To explore the potential for shifting production centres and distribution networks further, we measured the external diameter, thickness and perforation diameter of each bead (including beads not measured for stable isotopes) for the 12 burials included in this study ($n = 218$ beads total). During this process, we noted that beads seemed to cluster in distinct size categories and that burials often contained beads from multiple size categories. This observation is borne out in the metric data (Figure 6). In a comparison between the thickness and




Figure 6. Bead thickness against diameter for beads from three Ban Non Wat burials. Burials 90 and 105 are from Bronze Age 2; burial 263 is from Bronze Age 3 (figure by authors).

diameter of beads from three Bronze Age burials (90, 105 and 263), it can be seen that bead thickness varies in a continuous fashion, but that there are clear size classes for bead diameter—with small beads measuring between 2 and 3.5 mm in diameter, medium beads between 4 and 5 mm and large beads between 5.5 and 6.7 mm. Other burials show similar tri- or bi-modal distributions, although there is some overlap between size classes in different burials such that ‘medium’ sized beads in a different burial and/or time period might fall between the ‘medium’ and ‘large’ beads in Figure 6.

Consideration of average $\delta^{13}\text{C}$ against bead diameter reveals a pattern that seems to transcend time periods (Figure 7). In particular, the largest beads in all Bronze Age periods consistently have among the lowest $\delta^{13}\text{C}$ values. Note that the difference in average $\delta^{13}\text{C}$ for beads over 5.5 mm in diameter (avg. = -14%)—and for those less than 4.5 mm (avg. = -7%)—is twice as large as the largest absolute difference in $\delta^{13}\text{C}$ observed within a single modern shell (Figure 3) and more than twice as large as the maximum range observed within a single bead (2.9‰; average range within a bead is 0.8‰). This suggests that the difference in $\delta^{13}\text{C}$ among bead sizes is not due to random variations within shells but reflects the use of shells with different underlying $\delta^{13}\text{C}$ values. This pattern does not hold for bead thickness, nor for perforation size, which seem to vary continuously (i.e. unimodally) among different $\delta^{13}\text{C}$ values. No patterns are apparent when $\delta^{18}\text{O}$ values are plotted against bead diameter.




Figure 7. Average $\delta^{13}\text{C}$ plotted against bead diameter by time period at Ban Non Wat (figure by authors).

Our, admittedly small, sample of three modern shells in Figure 4 shows that different species growing in the same location overlap in their $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ isotopic composition. Therefore, the contrasting $\delta^{13}\text{C}$ values by bead size are unlikely to represent the exploitation of different shellfish species from the same location for producing beads of larger or smaller size. Likewise, because freshwater shellfish species often vary in the thickness of their shells, we would expect a similar correlation between $\delta^{13}\text{C}$ and bead thickness if exploitation of different species were responsible for the correlation with bead diameter.

Instead, we suggest that beads of different size were made from shells collected in different locations, especially during the Bronze Age when the pattern is most pronounced. The existence of multiple shell bead production centres near various bodies of water, each with different underlying conditions, could account for variation in $\delta^{13}\text{C}$ values of the shells. A propensity for producing beads of different sizes at these different centres—for example, due to local cultural transmission processes—could then lead to the pattern observed at Ban Non Wat. Perhaps different bead templates were used in different centres and handed down between craft workers. At the same time, there is no apparent correlation between $\delta^{13}\text{C}$ and bead perforation size. This suggests that a similar drilling technology may have been used across different bead production localities. For example, the use of stone drills made from blades removed from prepared cores could lead to similar perforation sizes (see Ciarla *et al.* 2017).

Assuming that beads of different sizes were made at different production centres, the inclusion of two or three sizes of beads within the same mortuary context indicates that the individuals buried with these items had access to multiple sources of beads during their lifetimes. This further points to the interconnectedness of Ban Non Wat and other

settlements during the Bronze Age. Our sample size of beads from the Neolithic is much smaller, but the same pattern does not seem to emerge during this time period. In this respect, the isotopic and bead size data point to the important social connections that developed across the landscape among different bead production centres in north-east Thailand, especially during the Bronze Age.

Mortuary wealth within the Bronze Age central burial ground at Ban Non Wat reached a level not seen elsewhere in Southeast Asia (Higham 2011, 2024). As well as large numbers of exotic shell disc beads, mortuary offerings included bangles of marine origin, copper-based axes sourced from the mines of the Khao Wong Prachan Valley in central Thailand (Pryce 2011) and marble probably derived from the Phetchabun Upland. Restricted access to these valuables—demonstrated by their absence from Early Bronze Age burials elsewhere in the settlement—has been ascribed to the rise of social aggrandisers who maintained elite status for about 150 years (Higham 2011, 2024).

The number of beads, as with other exotic valuables, fell sharply in the later Bronze Age 4, and again with the Bronze Age 5 where we find a tight range of $\delta^{13}\text{C}$ values consistent once again with a single and distinct source. Falling regional demand may have led to a reduction in the number of production centres that remained in use. No shell disc beads were found in any Iron Age grave, where glass, agate and carnelian beads took the place of shell disc beads as a form of mortuary offering and status symbol. As other sources of jewellery emerged, so shell ceased to be a preferred material.

Conclusions

All but one of the sampled beads from Ban Non Wat were made from shells from freshwater environments that persist today, across the low-lying wetlands of the Khorat Plateau. This contrasts with the presence in the same graves of bangles made from marine *Tridacna* and *Trochus* shells (Higham 2011). Despite the excavation of a considerable area, no evidence for local bead manufacture was encountered at Ban Non Wat; both the beads and the bangles were likely imported to the site.

Isotopic data, paired with morphological attribute data (bead diameter), provide compelling evidence for different shell sources and bead manufacturing locations over time. The earliest beads—from burial 461, a flexed hunter-gatherer interment, radiocarbon dated to 1521–1423 cal BC—are morphologically distinct from later beads, in retaining the external ridges of the shell and an uneven circumference (Figure 2). The $\delta^{13}\text{C}$ values of these early beads form a tight distribution, indicating that they all derive from the same production location. Yet they fall outside the isotopic range of modern shells, suggesting they were obtained from locations away from Ban Non Wat (or that local values were very different prior to modern times).

Relatively few shell disc beads were recovered from Neolithic 1 and 2 burials, but their numbers increase dramatically with the Early Bronze Age before declining again in the Late Bronze Age. Between *c.* 1050–800 BC, Early Bronze Age burials in the central part of the site reach a degree of mortuary wealth unmatched anywhere in Southeast Asia for the same period. Our data suggest that the individuals within these burials wore beads from several distinct production centres. As demand for shell disc beads

rose regionally, it is likely that more centres of production emerged, with some producing distinctively sized items. Some of these centres must have been next to bodies of water with different underlying aquatic environments, resulting in a relationship between $\delta^{13}\text{C}$ values and bead size.

Shell beads have been recovered at many other sites in Southeast Asia, and beyond, including evidence of Bronze Age production at the site of Tha Kae in central Thailand (Ciarla *et al.* 2017). Our work demonstrates the wealth of information contained within these artefacts and their potential for unlocking new insights into production and exchange in prehistoric societies. Application of similar analyses at other sites will strengthen our ability to identify and link potential water sources, production centres and consumer sites and thereby expand our understanding of exchange routes, links between regions and the role that shell beads played in projecting social status. Additional work should also seek to apply other isotopic systems, such as strontium, to shell bead provenance analysis, to cross-check the carbon and oxygen isotopic results.

Acknowledgements

We thank Jim Cassidy for initiating the study and connecting the authors, Susan Talcott, Brittany Elizondo, Tiffany Hirokawa and Roxanne Lamson for assistance in various steps of the stable isotope and bead measurement analysis, and two anonymous reviewers of an earlier draft. The excavation of Ban Non Wat was co-directed by Charles Higham and Rachanie Thosarat. We thank the National Research Council of Thailand and the Fine Arts Department for permitting us to undertake the fieldwork, the villagers of Ban Non Wat for their hospitality and contribution to the excavations, and our many colleagues who brought their expertise to the analysis of the site.

Funding statement

Funding for this work came from a University of California Davis Academic Senate Small Grant in Aid of Research to JWE. Excavations at Ban Non Wat were funded by the Marsden Fund and Earthwatch and its Research Corps.

Supplementary material

To view supplementary material for this article, please visit <https://doi.org/10.15184/aqy.2024.86>.

References

- ABRAHAM, S.A., P. GULLAPALLI, T.P. RACZEK & U.Z. RIZVI (ed.). 2016. Connections and complexity: new approaches to the archaeology of South Asia, in *Connections and complexity: new approaches to the archaeology of South Asia*: 15–34. Abingdon: Routledge.
- BAJNÓCZI, B. *et al.* 2013. Tracing the source of Late Neolithic *Spondylus* shell ornaments by stable isotope geochemistry and cathodoluminescence microscopy. *Journal of Archaeological Science* 40: 874–82. <https://doi.org/10.1016/j.jas.2012.09.022>
- BELLINA, B. 2014. Maritime Silk Roads' ornament industries: socio-political practices and cultural transfers in the South China Sea. *Cambridge Archaeological Journal* 24: 345–77. <https://doi.org/10.1017/S0959774314000547>
- BRUMFIEL, E. & T. EARLE (ed.). 1987. Specialization, exchange, and complex societies:

- an introduction, in *Specialization, exchange, and complex societies*: 1–9. Cambridge: Cambridge University Press.
- CARTER, A.K. 2015. Beads, exchange networks and emerging complexity: a case study from Cambodia and Thailand (500 BCE–CE 500). *Cambridge Archaeological Journal* 25: 733–57. <https://doi.org/10.1017/S0959774315000207>
- CARTER, A.K., L. DUSSUBIEUX, M.T. STARK & H.A. GILG. 2021. Angkor Borei and protohistoric trade networks: a view from the glass and stone bead assemblage. *Asian Perspectives* 60: 32–70. <https://doi.org/10.1353/asi.2020.0036>
- CIARLA, R., F. RISPOLI & P. YUKONGDI. 2017. Shell personal ornaments craft at the site of Tha Kae, Lopburi Province, central Thailand. Tracing the southward dispersal of the drilling technique. *Journal of Indo-Pacific Archaeology* 41: 30–65. <https://doi.org/10.7152/jipa.v41i0.15011>
- DEITH, M.R. 1985. Seasonality from shells: an evaluation of two techniques for seasonal dating of marine molluscs, in N.R.J. Fieller, D.D. Gilbertson & N.G.A. Ralph (ed.) *Palaeobiological investigations research design, methods and data analysis* (British Archaeological Reports International Series 266): 119–130. Oxford: BAR.
- DETTMAN, D.L., A.K. REISCHE & K.C. LOHMANN. 1999. Controls on the stable isotope composition of seasonal growth bands in aragonitic fresh-water bivalves (Unionidae). *Geochimica et Cosmochimica Acta* 63(7–8): 1049–57. [https://doi.org/10.1016/S0016-7037\(99\)00020-4](https://doi.org/10.1016/S0016-7037(99)00020-4)
- EERKENS, J.W., J.S. ROSENTHAL, N.E. STEVENS, A. CANNON, E.L. BROWN & H.J. SPERO. 2010. Stable isotope provenance analysis of *Olivella* shell beads from the Los Angeles Basin and San Nicolas Island. *Journal of Island & Coastal Archaeology* 5: 105–19. <https://doi.org/10.1080/15564890902955327>
- FRANCIS JR, P. 1991. Beadmaking at Arikamedu and beyond. *World Archaeology* 23: 28–43.
- GOEWERT, A., D. SURGE, S.J. CARPENTER & J. DOWNING. 2007. Oxygen and carbon isotope ratios of *Lampsilis cardium* (Unionidae) from two streams in agricultural watersheds of Iowa, USA. *Palaeogeography, Palaeoclimatology, Palaeoecology* 252: 637–48. <https://doi.org/10.1016/j.palaeo.2007.06.002>
- GROSSMAN, E.L. & T-L KU. 1986. Oxygen and carbon isotope fractionation in biogenic aragonite: temperature effects. *Chemical Geology* 59: 59–74. [https://doi.org/10.1016/0168-9622\(86\)90057-6](https://doi.org/10.1016/0168-9622(86)90057-6)
- HIGHAM, C.F.W. 2011. The Bronze Age of Southeast Asia: new insight on social change from Ban Non Wat. *Cambridge Archaeological Journal* 21: 365–89. <https://doi.org/10.1017/S0959774311000424>
- 2024. Aggrandisers and the first copper-base metallurgy in Southeast Asia. *Antiquity* 98: 433–53. <https://doi.org/10.15184/aqy.2024.11>
- HIGHAM, C.F.W. & R. BANNANURAG. 1990. *The excavation of Khok Phanom Di, volume 1: the excavation, chronology and human burials*. London: Society of Antiquaries of London.
- HIGHAM, C.F.W. & T.F.G. HIGHAM. 2009. A new chronological framework for prehistoric Southeast Asia, based on a Bayesian model from Ban Non Wat. *Antiquity* 83: 125–44. <https://doi.org/10.1017/S0003598X00098136>
- HIGHAM, C.F.W. & A. KIJNGAM. (ed.) 2009. *The origins of the civilization of Angkor, volume 3: the excavation Ban Non Wat. Part one: introduction*. Bangkok: The Fine Arts Department of Thailand.
- HIGHAM, C.F.W. & A. KIJNGAM. 2010. *The origins of the civilization of Angkor, volume 4: the excavation Ban Non Wat. Part two: the Neolithic occupation*. Bangkok: The Fine Arts Department of Thailand.
- 2012. *The origins of the civilization of Angkor, volume 5: the excavation Ban Non Wat. Part three: the Bronze Age*. Bangkok: The Fine Arts Department of Thailand.
- HIGHAM, C.F.W. & R. THOSARAT. 2020. An early hunter-gatherer site at Ban Non Wat, Northeast Thailand. *Journal of Indo-Pacific Archaeology* 43: 93–96. <https://doi.org/10.7152/jipa.v43i0.15413>
- KEITH, M.L., G.M. ANDERSON & R. EICHLER. 1964. Carbon and oxygen isotope composition of mollusk shells from marine and fresh-water environments. *Geochimica et Cosmochimica Acta* 28: 1757–86. [https://doi.org/10.1016/0016-7037\(64\)90021-3](https://doi.org/10.1016/0016-7037(64)90021-3)
- KENDALL, C. & T.B. COPLEN. 2001. Distribution of oxygen-18 and deuterium in river waters across the United States. *Hydrological Processes* 15: 1363–93. <https://doi.org/10.1002/hyp.217>
- KENNETT, D.J. & B. VOORHIES. 1996. Oxygen isotopic analysis of archaeological shells to detect

- seasonal use of wetlands on the southern Pacific coast of Mexico. *Journal of Archaeological Science* 23: 689–704.
<https://doi.org/10.1006/jasc.1996.0065>
- KLYSUBUN, W., Y. THONGKAM, S. PONGKRAPAN, K. WON-IN, J.T. THIENPRASERT & P. DARARUTANA. 2011. XAS study on copper red in ancient glass beads from Thailand. *Analytical and Bioanalytical Chemistry* 399: 3033–40.
<https://doi.org/10.1007/s00216-010-4219-1>
- MOOK, W.G. & J.C. VOGEL. 1968. Isotopic equilibrium between shells and their environment. *Science* 159: 874–5.
<https://doi.org/10.1126/science.159.3817.874>
- OKA, R. & C.M. KUSIMBA. 2008. The archaeology of trading systems, part 1: towards a new trade synthesis. *Journal of Archaeological Research* 16: 339–95.
<https://doi.org/10.1007/s10814-008-9023-5>
- PEACOCK, E., A.R. PALMER, T. XIA, W. BACON-SCHULTE, B. CARLOCK & J. SMITH. 2010. Chemical sourcing of a prehistoric freshwater shell artifact using laser ablation-inductively coupled plasma-mass spectrometry. *Archaeology of Eastern North America* 38: 91–99.
- PÉREZ, A.E., D.A. BATRES, I. ROCCHETTA, M.R. EPPIS, M.L. BIANCHI & C.M. LUQUET. 2020. Paleoenvironmental reconstruction using stable isotopes and trace elements from archaeological freshwater bivalve shell fragments in Northwest Patagonia, Argentina. *Quaternary International* 547: 22–32.
<https://doi.org/10.1016/j.quaint.2020.02.029>
- PILDITCH, J.S. 1992. The glass beads of Ban Bon Noen, central Thailand. *Asian Perspectives* 31: 171–81.
- PRYCE, T.O. 2011. Technical analysis of Bronze Age Ban Non Wat copper-base artefacts, in C.F.W. Higham & A. Kijngam (ed.) *The origins of the civilization of Angkor, volume 5: the excavation Ban Non Wat. Part three: the Bronze Age*: 489–97. Bangkok: The Fine Arts Department of Thailand.
- RYAN, W.B.F. *et al.* 2009. Global multi-resolution topography (GMRT) synthesis data set. *Geochemistry, Geophysics, Geosystems* 10(3).
<https://doi.org/10.1029/2008GC002332>
- SEMAN, S., L. DUSSUBIEUX, C. CLOQUET & T.O. PRYCE. 2021. Strontium isotope analysis in ancient glass from South Asia using portable laser ablation sampling. *Archaeometry* 63: 88–104.
<https://doi.org/10.1111/arc.12618>
- SHACKLETON, N.J. 1973. Oxygen isotope analysis as a means of determining season of occupation of prehistoric midden sites. *Archaeometry* 15: 133–41.
<https://doi.org/10.1111/j.1475-4754.1973.tb00082.x>
- SHACKLETON, N. & H. ELDERFIELD. 1990. Strontium isotope dating of the source of Neolithic European *Spondylus* shell artefacts. *Antiquity* 64: 312–15.
<https://doi.org/10.1017/S0003598X00077942>
- SHACKLETON, N. & C. RENFREW. 1970. Neolithic trade routes re-aligned by oxygen isotope analyses. *Nature* 228: 1062–65.
- SKIPPINGTON, J., T. MANNE & P. VETH. 2021. Isotopic indications of late Pleistocene and Holocene paleoenvironmental changes at Boodie Cave archaeological site, Barrow Island, Western Australia. *Molecules* 26.
<https://doi.org/10.3390/molecules26092582>
- STEPHENS, M., D. MATTEY, D.D. GILBERTSON & C.V. MURRAY-WALLACE. 2008. Shell-gathering from mangroves and the seasonality of the Southeast Asian Monsoon using high-resolution stable isotopic analysis of the tropical estuarine bivalve (*Geloina erosa*) from the Great Cave of Niah, Sarawak: methods and reconnaissance of molluscs of early Holocene and modern times. *Journal of Archaeological Science* 35: 2686–97.
<https://doi.org/10.1016/j.jas.2008.04.025>
- UREY, H.C., S. EPSTEIN, C. MCKINNEY & J. MCCREA. 1948. Method for measurement of paleotemperatures. *Bulletin of the Geological Society of America* 62: 1359–60.
- VANHAEREN, M., F. D'ERRICO, I. BILLY & F. GROUSSET. 2004. Tracing the source of Upper Palaeolithic shell beads by strontium isotope dating. *Journal of Archaeological Science* 31: 1481–88.
<https://doi.org/10.1016/j.jas.2004.03.011>
- VERSTEEGH, E.A., H.B. VONHOF, S.R. TROELSTRA, R.J. KAANDORP & D. KROON. 2010. Seasonally resolved growth of freshwater bivalves determined by oxygen and carbon isotope shell chemistry. *Geochemistry, Geophysics, Geosystems* 11.
<https://doi.org/10.1029/2009GC002961>