

Increased Range—

OXOID

COMPLETE CULTURE MEDIA

TABLETS

For single tubes and small quantities of media

GRANULES

For larger scale preparation

| | |
|------------------------------------|--|
| MacCONKEY BROTH (Official Formula) | AGAR-AGAR |
| MacCONKEY BROTH (Purple) | GLUCOSE PEPTONE BROTH |
| MacCONKEY AGAR | NUTRIENT AGAR |
| GLUCOSE PEPTONE AGAR | NUTRIENT BROTH |
| LAB-LEMCO BROTH | YEAST EXTRACT AGAR |
| LAB-LEMCO AGAR | EDWARDS AGAR (Modified) |
| THIOGLYCOLLATE MEDIUM (Brewer) | MILK AGAR (Official Formula) |
| SABOURAUD AGAR | MILK AGAR (For Roll Tubes) |
| PEPTONE WATER | TETRATHIONATE BROTH BASE |
| BRILLIANT GREEN BILE (2%) BROTH | WILSON & BLAIR MEDIUM (Bismuth Sulphite Agar) |

For those who prefer to prepare their own media,
the following **OXOID** preparations are available:—

| | |
|-------------------------------|-------------------------|
| BACTERIOLOGICAL LIVER EXTRACT | AGAR-AGAR |
| DESICCATED OX BILE | DESICCATED OX HEART |
| BILE SALTS | LAB-LEMCO |
| MYCOLOGICAL PEPTONE | BACTERIOLOGICAL PEPTONE |
| BACTERIOLOGICAL YEAST EXTRACT | |

FULL INFORMATION ON ANY OF THE ABOVE IS READILY AVAILABLE FROM

OXO LIMITED (Medical Dept.)

Thames House, Queen St. Place, London, E.C.4. Tel. CENTral 9781

CONTENTS

(All rights reserved)

| | PAGE |
|---|------|
| MAITLAND, MARY L. COWAN. The rapid detection of <i>Mycobacterium tuberculosis</i> by microscopic examination of milk from individual cows and from groups of ten cows | 397 |
| LANCASTER, H. O. Statistical control in haematology. (With 1 Figure in the Text) | 402 |
| MASON, J. H. Alkali-dissolved diphtheria toxoid-antitoxin floccules adsorbed on aluminium carriers. Preparation, and immunity experiments in animals | 418 |
| ALLEN, L. A., THOMAS, G., THOMAS, M. C. C., WHEATLAND, A. B., THOMAS, H. N., JONES, E. E., HUDSON, J. and SHERWOOD, H. P. Repeated re-use of sea water as a medium for the functioning and self-cleansing of molluscan shellfish. (With 8 Figures in the Text) | 431 |
| MYNORS, L. S., HEARD, D. H. and COOMBS, R. R. A. An experimental study of haemolytic disease of the newborn due to isoimmunization of pregnancy. II. Report of an attempt to produce anti-red cell isoantibodies in guinea-pigs, together with a note on the blood picture of one-day-old guinea-pigs | 458 |
| REILLY, LILIAN V. Human tuberculosis of bovine origin in Northern Ireland | 464 |
| SMITH, J., GALLOWAY, W. H. and SPEIRS, A. L. Infantile gastro-enteritis with special reference to the specific serological type O55 B5 H6 (<i>Beta</i> type) of <i>Bacterium coli</i> | 472 |
| COOMBS, R. R. A., BLOMFIELD, A. M. and ROBERTS, G. FULTON. The conglutination phenomenon. VII. A study of the interaction of complement components and conglutinin in the process of conglutination | 484 |
| PICKETT, M. J. and NELSON, ERIC L. <i>Brucella</i> bacteriophage | 500 |
| WILLIAMS, R. E. O. and HIRCH, ANN. The detection of streptococci in air. (With Plate 7) | 504 |

The Journal of Hygiene is issued in four parts to a volume.

Quotations can be given for back volumes on application.

Papers for publication should be sent to Professor E. T. C. SPOONER, London School of Hygiene and Tropical Medicine, Keppel Street, Gower Street, London, W.C. 1. Other communications should be addressed to the University Press, Cambridge.

Papers forwarded to the Editor for publication are understood to be offered to *The Journal of Hygiene* alone, unless the contrary is stated.

Contributors receive twenty copies of their papers free. Twenty-five additional copies may be had at cost price: these should be ordered when the final proof is returned.

The subscription price is £3. 15s. 0d. per volume (post-free), payable in advance; single numbers 20s. net. Subscriptions may be sent to any Bookseller, or to The Cambridge University Press, Bentley House, 200 Euston Road, London, N.W. 1.

Enquiries from the U.S.A. should be addressed to the American Branch of the Press at 51 Madison Avenue, New York, 10.