

consciousness “+” (a- / hyperkinetic mutism with emotional reactions, understanding of addressed speech);

2nd - 67 (32%) patients had manifestations of physical and cognitive abilities with minimal consciousness “-” (a- / hyperkinetic mutism without emotional manifestations and understanding of addressed speech);

3rd - 95 (40%) patients had only the manifestation of physical capabilities at the exit from the vegetative status.

4th - 11 (10%) patients had a low manifestation of mental activity in the form of physical capabilities with a vegetative status.

Conclusions: 4 variants of mental activity in children after acute severe brain damage have been identified: from minimal involuntary reactions or their absence in vegetative status to voluntary actions according to the instructions of an adult in minimal consciousness “+”. Taking into account the variability of mental activity helps to differentiate the methods of psychiatric and psychological-pedagogical assistance in the recovery of children already in the early stages of rehabilitation.

Disclosure of Interest: None Declared

EPP0108

Schizophrenia and Polycythemia Vera: A Case Report

N. El Moussaoui*, S. Bahetta, I. Belabbes, F. Laboudi and A. Ouanass

Arrazi University Psychiatric Hospital, Salé, Morocco

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.331

Introduction: Schizophrenia is a severe mental disorder marked by abnormal interpretations of reality, often resulting in hallucinations, delusions, and disordered thinking that significantly impairs daily functioning and can be disabling. Lifelong treatment is necessary, and early intervention can help manage symptoms and improve long-term outcomes.

Polycythemia Vera (PV) is a chronic myeloproliferative neoplasm causing an excess of red blood cells in the peripheral blood (polyglobulia). While the disease typically presents with symptoms, it can also be asymptomatic and discovered incidentally during routine laboratory tests, leading to a diagnosis of polycythemia when no secondary cause is apparent.

While early 20th-century literature linked PV to intense neurological and psychiatric symptoms, contemporary studies rarely make such references.

Objectives: The aim of this study is to explore, through a clinical case of a patient undergoing treatment for treatment-resistant schizophrenia with clozapine, and concurrently diagnosed with Polycythemia Vera, the potential causes of this condition. We seek to discern whether it represents mere comorbidities or if Polycythemia Vera is an adverse effect of antipsychotic treatment, particularly with clozapine.

Methods: A 41-year-old patient, with a history of cranial trauma at the age of 5 and 19 years of treatment for schizophrenia, also has a tobacco use disorder. While hospitalized for the management of symptomatic reactivation of schizophrenia, despite being on clozapine, the patient underwent various therapeutic combinations with no observed clinical improvement. A few months later, follow-up blood tests indicated an elevation in all blood cell lines.

An internal medicine consultation was sought, resulting in the diagnosis of Polycythemia Vera.

Results: The evaluations conducted led us to the conclusion that there are two distinct nosological entities, with the treatment of the psychiatric condition revealing true polycythemia. Even after reducing the doses of clozapine and changing the atypical antipsychotic, all subsequent evaluations showed no effectiveness in managing the psychiatric disorder or improvement in the hematological condition.

Conclusions: In summary, schizophrenia is a severe and lifelong mental disorder requiring early intervention for symptom management. Polycythemia Vera (PV), a myeloproliferative disorder, typically presents with symptoms but can also be asymptomatic.

While early literature linked PV to intense neurological and psychiatric symptoms, contemporary studies seldom reference such associations. The coexistence of schizophrenia and PV in a patient underscores the need for comprehensive and interdisciplinary care to address the complex interplay between mental and physical health. Further research is needed to deepen our understanding of concurrent psychiatric and hematological conditions.

Disclosure of Interest: None Declared

EPP0109

The Impact of Internet Use on the Parameters of Attention in Adults

A. Mihai^{1*} and M. Rebeca-Isabela²

¹Psychiatry, University of Medicine, Pharmacy, Science and Technology and ²Mureş County Hospital, Targu-Mures, Romania

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.332

Introduction: Internet use in the adult population is growing at alarming rates. The latest statistical data show an average internet usage time of 6 hours and 58 minutes (2023), an increase of 1% compared to 2021. Research studies on the influence of the excess use of internet on attention is in its prime years, and clear steps need to be made in an attempt to clarify current hypotheses and to find effective methods for prevention. Nowadays, one of the most powerful influences on attention is the use of the internet, which, more often than not, crosses the line of addiction.

Objectives: The initial hypothesis is that in the event of exposure to a high number of stimuli, the ability to switch attention to a single task may only be possible at a superficial level. The aim of this study was to assess the impact that excess internet use has on the ability to maintain attention in the adult population. The present study aims to sketch a well-established structure and direction of research in the field of attention and its effects on human functioning.

Methods: Using the DSM 5-TR diagnostic criteria for pathological Internet gaming disorder we enrolled 60 people who expressed their consent to participate in the study. We check psychiatric comorbidities using SCID II. As a method for evaluating changes in the level of attention, we used of the Stroop test. The results were analysed with the SPSS program (version 23).

Results: The results showed a marked decrease in the ability to maintain attention, without increasing the number of stimuli. Although excessive Internet use leads to changes in attention parameters, research in this area is scarce and incomplete.

Currently, most of the published studies focus on a causal relationship between the pathological use of the Internet and the appearance of attention deficit/hyperkinetic disorder, especially in children and adolescents. Although the results are promising, we cannot neglect the multitude of additional consequences of excess Internet use, which these studies targeting a single pathology overlook. Moreover, using the Internet involves exposure to an ever-increasing number of stimuli, which is why switching attention and maintaining it is currently an insufficiently researched parameter. Regarding the impact of Internet use on individual functioning, there is a relatively modest number of studies in the literature that outline a correlation between excess Internet use and various psychiatric comorbidities.

Conclusions: The impact of the research on the general population could be an increased awareness of negative effects and the development of prevention programs.

Disclosure of Interest: None Declared

COVID-19 and related topics

EPP0111

Psychometric Properties of the Depression, Anxiety, Stress Scales-21 (DASS-21) in a Portuguese Sample during the early stage of the COVID-19 pandemic

C. Laranjeira^{1,2*}, M. A. Dixe^{1,2} and A. I. Querido^{1,2}

¹School of Health Sciences and ²ciTechCare, Polytechnic University of Leiria, Leiria, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.333

Introduction: The COVID-19 global crisis has resulted in significant disruptions in the lives of students in higher education, leading to negative consequences for their academic achievements and general psychological well-being.

Objectives: In this study, we sought to examine the psychometric properties of the Depression Anxiety Stress Scale-21 (DASS-21) among students in Portuguese higher education institutions during the initial phase of the COVID-19 pandemic and its efficacy in capturing mental health symptoms due to a global health crisis.

Methods: In this cross-sectional study, a convenience sampling method was used to enlist a total of 1522 participants. The sample consisted of 75.1% women and 79.2% undergraduate students. Participants completed an electronic survey that was designed using the Depression Anxiety Stress Scale-21 (DASS-21) — a self-report instrument measuring anxiety, depression, and stress.

Results: The findings of the study indicated a significant occurrence of depressive symptoms [≥ 10] (N = 434, 28.5%), anxiety symptoms [≥ 7] (N = 551, 36.2%), and stress symptoms [≥ 11] (N = 544, 35.7%). Based on the collected data, a Confirmatory Factor Analysis (CFA) was conducted in order to examine the factor structure of the scale. The analysis revealed a three-factor solution that corresponded to the three subscales of the DASS-21. The Heterotrait-Monotrait (HTMT) correlation ratio was then used to assess the discriminant validity, with good results. Results showed that the DASS21 has satisfactory reliability indexes (Cronbach's $\alpha > 0.90$).

Conclusions: In light of the notable changes in living conditions brought by the COVID-19 pandemic, the present study has shown that the DASS-21 instrument has maintained its reliability and validity. Consequently, this finding supports the appropriateness of using the DASS-21 as a screening tool for assessing mental health among students in Portugal. Moreover, it is recommended that academics and healthcare practitioners use the DASS-21 as a tool for assessing the levels of psychological distress experienced by students. Additional validation studies of this scale are required, using bigger and more representative populations.

Disclosure of Interest: None Declared

EPP0112

The neuro-psychological manifestations of COVID-19 in healthcareworkers

A. Ghenim¹, I. Kacem^{1*}, A. Chouchane¹, A. Aloui¹, C. Sridi², A. Fekih², M. Hafsia², M. Maoua¹, M. Kahloul³ and N. Mrizak¹

¹Occupational Medicine Department, Farhat Hached Academic hospital; ²Occupational Medicine Department and ³Anesthesia and Intensive Care Department, Sahloul Academic hospital, sousse, Tunisia

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.334

Introduction: At the beginning of the Covid-19 pandemic, respiratory expression of SARS-CoV-2 infection was the most worrying one. Later, other symptoms appeared to be more disturbing such as neurological and psychiatric manifestations, which may be due to direct or indirect effects of this virus on the central nervous system.

Objectives: To assess the prevalence of neuropsychological manifestations of covid-19 in healthcareworkers and to identify their risk factors.

Methods: This is a cross-sectional descriptive epidemiological study, carried out in the teaching hospitals of Farhat Hached and Sahloul of Sousse. All healthcareworkers, having tested positive for SARS-COV 2 during the period from 01/09/2020 to 28/02/2021 were enrolled. The collection of socio-professional and medical data was based on a pre-established synoptic form completed during an interview with the participants.

Results: A total of 953 COVID-19 patients were enrolled in this study. The mean age was 40.1 ± 10.5 years, with a sex ratio of 0.32. In our sample, 37.9% of patients had comorbidities such as psychiatric history (4.9%) and neurological history (2.4%). The prevalence of neuropsychological manifestations of covid-19 was 72.6%. The main neuropsychological manifestations were headache (50.3%), anosmia (40.7%), dysgeusia (29.9%), sleep disturbances (0.5%), dizziness (1, 2%) and paresthesia (0.3%). Neuropsychological symptoms of long covid were dominated by memory impairment (10.7%), anosmia (8.5%), headache (7.3%), dizziness (3.4%) and sleep disturbances (3.1%).

The occurrence of neuropsychological manifestations was significantly associated with age (OR=1.6; $p < 10^{-3}$), male gender (OR=0.57; $p=0.03$), smoking (OR=1.7; $p=0.033$), history of hypertension (OR=1.6; $p=0.038$), history of diabetes (OR=2.4; $p < 10^{-3}$) and hospitalization (OR=4.03 ; $p < 10^{-3}$).