## INSTRUCTIONS TO CONTRIBUTORS

The *Journal of Helminthology* publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance, but only in exceptional circumstances will systematic or taxonomic studies be acceptable.

Manuscripts, which must be in English or French (with an English summary) should be accompanied by a letter signed by *all* the authors and should be addressed to:

The Editors, Journal of Helminthology, London School of Hygiene and Tropical Medicine, 395A Hatfield Road, St Albans, Herts AL4 0XQ. England.

Three copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief Introduction; Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Research Notes should also be preceded by a brief abstract. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration will be printed is  $12.0 \times 20.0$  cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space, should be indicated. Some indication of scale (preferably a scale bar) should normally be given on the figure. Photocopies of illustrations should be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size  $12 \cdot 0 \times 20 \cdot 0$  cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

ASHFORD, R. W. & BARNISH, G. (1989) Strongyloides fuelleborni and similar parasites in animals and man. In: Strongyloidiasis: a major roundworm infection of man (editor, D. I. Grove) pp. 271–286. Taylor & Francis: London.

GARSIDE, P. & BEHNKE, J. M. (1989) Primary infection of hamsters with *Ancylostoma ceylanicum*. *Parasitology*, **98**, 283–289.

## **Contents**

The effects of crowding on adults of <i>Philophthalmus nocturnus</i> grown in domestic chicks. V. G. M. SWARNAKUMARI and R. MADHAVI  255—  Comparison of restriction fragment length polymorphisms of ribosomal DNA between <i>Diphyllobothrium nihonkaiense</i> and <i>D. latum.</i> T. MATSUURA, G. BYLUND and K. SUGANE  261—  Ultrastructure of the spermatid and spermatozoon of <i>Macracanthorhynchus hirudinaceus.</i> B. ZHAO and B. LIU  267—  The effect of catecholamines and catecholamine antagonists on the third larval moult of <i>Dirofilaria immitis</i> in vitro. E. V. WARBRICK and S. A. WARD  273—  The use of 35S-labelled <i>Toxocara canis</i> to determine larval numbers in tissues. K. C. CARTER  279—  Treatment of <i>Nippostrongylus brasiliensis</i> in normal and SPF rats using tetramisole loaded into zeolite. S. K. SHAKER, A. DYER and D. M. STOREY  288—  Efficacy of ivermectin against <i>Parastrongylus malaysiensis</i> infection in rats. S. AMBU, J. W. MAK and C. S. NG  Immunodiagnosis of human trichinellosis using excretory-secretory (ES) antigen. P. MAHANNOP, W. CHAICUMPA, P. SETASUBAN, N. MORAKOTE and		Pages
Comparison of restriction fragment length polymorphisms of ribosomal DNA between Diphyllobothrium nihonkaiense and D. latum. T. MATSUURA, G. BYLUND and K. SUGANE  Ultrastructure of the spermatid and spermatozoon of Macracanthorhynchus hirudinaceus. B. ZHAO and B. LIU  The effect of catecholamines and catecholamine antagonists on the third larval moult of Dirofilaria immitis in vitro. E. V. WARBRICK and S. A. WARD  The use of 35S-labelled Toxocara canis to determine larval numbers in tissues. K. C. CARTER  Treatment of Nippostrongylus brasiliensis in normal and SPF rats using tetramisole loaded into zeolite. S. K. SHAKER, A. DYER and D. M. STOREY  Efficacy of ivermectin against Parastrongylus malaysiensis infection in rats. S. AMBU, J. W. MAK and C. S. NG  Immunodiagnosis of human trichinellosis using excretory-secretory (ES) antigen. P. MAHANNOP, W. CHAICUMPA, P. SETASUBAN, N. MORAKOTE and	planorbid snail, Indoplanorbis exustus. T. S. DUNN, P. H. DANG, R. E. B.	43–254
between Diphyllobothrium nihonkaiense and D. latum. T. MATSUURA, G. BYLUND and K. SUGANE  Ultrastructure of the spermatid and spermatozoon of Macracanthorhynchus hirudinaceus. B. ZHAO and B. LIU  267–  The effect of catecholamines and catecholamine antagonists on the third larval moult of Dirofilaria immitis in vitro. E. V. WARBRICK and S. A. WARD  The use of 35S-labelled Toxocara canis to determine larval numbers in tissues. K. C. CARTER  Treatment of Nippostrongylus brasiliensis in normal and SPF rats using tetramisole loaded into zeolite. S. K. SHAKER, A. DYER and D. M. STOREY  Efficacy of ivermectin against Parastrongylus malaysiensis infection in rats. S. AMBU, J. W. MAK and C. S. NG  Immunodiagnosis of human trichinellosis using excretory-secretory (ES) antigen. P. MAHANNOP, W. CHAICUMPA, P. SETASUBAN, N. MORAKOTE and	s of crowding on adults of <i>Philophthalmus nocturnus</i> grown in domestic V. G. M. SWARNAKUMARI and R. MADHAVI	55–259
The effect of catecholamines and catecholamine antagonists on the third larval moult of <i>Dirofilaria immitis</i> in vitro. E. V. WARBRICK and S. A. WARD  The use of <sup>35</sup> S-labelled <i>Toxocara canis</i> to determine larval numbers in tissues. K. C. CARTER  Treatment of <i>Nippostrongylus brasiliensis</i> in normal and SPF rats using tetramisole loaded into zeolite. S. K. SHAKER, A. DYER and D. M. STOREY  Efficacy of ivermectin against <i>Parastrongylus malaysiensis</i> infection in rats. S. AMBU, J. W. MAK and C. S. NG  Immunodiagnosis of human trichinellosis using excretory-secretory (ES) antigen. P. MAHANNOP, W. CHAICUMPA, P. SETASUBAN, N. MORAKOTE and	en Diphyllobothrium nihonkaiense and D. latum. T. MATSUURA,	61–266
moult of <i>Dirofilaria immitis</i> in vitro. E. V. WARBRICK and S. A. WARD  273–  The use of <sup>35</sup> S-labelled <i>Toxocara canis</i> to determine larval numbers in tissues. K. C. CARTER  279–  Treatment of <i>Nippostrongylus brasiliensis</i> in normal and SPF rats using tetramisole loaded into zeolite. S. K. SHAKER, A. DYER and D. M. STOREY  288–  Efficacy of ivermectin against <i>Parastrongylus malaysiensis</i> infection in rats. S. AMBU, J. W. MAK and C. S. NG  293–  Immunodiagnosis of human trichinellosis using excretory-secretory (ES) antigen. P. MAHANNOP, W. CHAICUMPA, P. SETASUBAN, N. MORAKOTE and	ture of the spermatid and spermatozoon of Macracanthorhynchus naceus. B. ZHAO and B. LIU	67–272
K. C. CARTER  279  Treatment of Nippostrongylus brasiliensis in normal and SPF rats using tetramisole loaded into zeolite. S. K. SHAKER, A. DYER and D. M. STOREY  288  Efficacy of ivermectin against Parastrongylus malaysiensis infection in rats. S. AMBU, J. W. MAK and C. S. NG  293  Immunodiagnosis of human trichinellosis using excretory-secretory (ES) antigen. P. MAHANNOP, W. CHAICUMPA, P. SETASUBAN, N. MORAKOTE and	of catecholamines and catecholamine antagonists on the third larval of <i>Dirofilaria immitis</i> in vitro. E. V. WARBRICK and S. A. WARD	73–278
Efficacy of ivermectin against <i>Parastrongylus malaysiensis</i> infection in rats. S. AMBU, J. W. MAK and C. S. NG  Immunodiagnosis of human trichinellosis using excretory-secretory (ES) antigen. P. MAHANNOP, W. CHAICUMPA, P. SETASUBAN, N. MORAKOTE and		79–287
S. AMBU, J. W. MAK and C. S. NG  293–  Immunodiagnosis of human trichinellosis using excretory-secretory (ES) antigen. P. MAHANNOP, W. CHAICUMPA, P. SETASUBAN, N. MORAKOTE and	of Nippostrongylus brasiliensis in normal and SPF rats using tetramisole d into zeolite. S. K. SHAKER, A. DYER and D. M. STOREY	88–292
P. MAHANNOP, W. CHAICUMPA, P. SETASUBAN, N. MORAKOTE and		93–296
	AHANNOP, W. CHAICUMPA, P. SETASUBAN, N. MORAKOTE and	97–304
Radiolabelling of the excretory-secretory and somatic antigens of <i>Anisakis simplex</i> larvae. K. SUGANE, S. SUN and T. MATSUURA 305–	lling of the excretory-secretory and somatic antigens of <i>Anisakis simplex</i> e. K. SUGANE, S. SUN and T. MATSUURA	05–309
Isoenzymatic pattern and structure of glutamate dehydrogenase from Ascaris suum.  C. CUTILLAS, B. RODRIGUEZ and D. C. GUEVARA  310–		10–312
Subject Index 313-	dex 3	13–315
Author Index 316-	dex 3	16–318
Contents of Volume	of Volume	i–iv