

APERIODIC FLUX VARIABILITY IN A 0535+262

MARK H. FINGER

*Compton Observatory Science Support Center/USRA
Goddard Space Flight Center*

ROBERT B. WILSON AND B. ALAN HARMON
NASA/Marshall Space Flight Center

AND

WILLIAM S. PACIESAS
University of Alabama Huntsville

Abstract. A “giant” outburst of A 0535+262, a transient X-ray binary pulsar, was observed in 1994 February and March with the Burst and Transient Source Experiment (BATSE) onboard the Compton Gamma-Ray Observatory. During the outburst power spectra of the hard X-ray flux contained a QPO-like component with a FWHM of approximately 50% of its center frequency. Over the course of the outburst the center frequency rose smoothly from 35 mHz to 70 mHz and then fell to below 40 mHz. We compare this QPO frequency with the neutron star spin-up rate, and discuss the observed correlation in terms of the beat frequency and Keplerian frequency QPO models in conjunction with the Ghosh-Lamb accretion torque model.

1. Introduction

A 0535+262 is a 103 s X-ray pulsar in a binary system with the Be star HDE 245770. Since its initial discovery in 1975 (Rosenberg *et al.* 1975), the source has been frequently observed to undergo transient outbursts. The outbursts show a range of peak intensities, with the largest reaching 3 Crab in the 2–10 keV band. For a review of previous observations see Giovannelli & Graziati (1992).

A major outburst of A 0535+262 occurred in 1994 February and March (Wilson *et al.* 1994a). Hard X-ray observations with BATSE were made continuously during the 50 day duration of the outburst. At the peak of

the outburst the intrinsic spin up rate determined from pulse timing was approximately $1.2 \times 10^{-11} \text{ Hz s}^{-1}$ (Wilson *et al.* 1994b), clearly indicating the presence of an accretion disk. The formation of a transient accretion disk during “giant” outbursts has previously been inferred from optical and UV observations (Motch *et al.* 1991).

During 27 days of the outburst a broad quasi-periodic oscillation (QPO) like feature appeared in Fourier power spectra of the flux (Finger *et al.* 1994b). One possible explanation for this feature is the beat frequency model (Alpar & Shaham 1985; Lamb *et al.* 1985). In this model blobs of matter, in the process of being entrained in the neutron star’s magnetic field, orbit the neutron star at approximately the Keplerian frequency of the inner edge of the accretion disk, accreting at a rate that is modulated by the magnetic field. This produces a peak in the power spectra at the beat frequency between the Keplerian and the pulsar spin frequencies. Another possible model is that the inner edge of the accretion disk contains structures that persist for a few cycles around the neutron star, and modulate the observed flux by obscuration (Van der Klis *et al.* 1987). In this case, the power spectral feature should be located near the Keplerian frequency of the disk inner edge.

Both of these models predict a simple relationship between the QPO frequency and the rate of mass accretion through the disk. The accretion rate also determines the torque on the neutron star. We compare the observed relationship between the neutron star spin up rate and the QPO frequency with the predictions of the beat frequency or Keplerian frequency models combined with the Ghosh & Lamb (1979) accretion torque model. The beat frequency model does best at predicting the trend of the data, but the Keplerian frequency model is not ruled out by the observations.

2. Observations and Analysis

BATSE (Fishman *et al.* 1989) has eight unshielded planar scintillation detectors oriented in the corner directions of a cube. Its full sky field of view allows the continuous monitoring of transient outbursts.

Pulsed hard X-ray emission from A 0535+262 was first detected with BATSE on January 28th (TJD 9380). The flux initially remained at a low level, but then on February 3 the flux began to rise quickly (Finger *et al.* 1994a), peaking on February 18 (Wilson *et al.* 1994a,b). The last BATSE pulsed flux detection was on March 20, fifty days after the first detection. The light curve of the February-March outburst as determined by the Earth occultation method is shown in panel A of Fig. 1.

A pulse timing analysis was performed to determine the intrinsic spin-up rate of A 0535+262. This analysis used the binary orbital parameters




Figure 1. A) shows the A 0535+262 flux history determined from Earth occultations. Also shown is the interval of QPO detections. B) shows the intrinsic spin-up of A 0535+262 during the outburst.

recently determined by Finger *et al.* (1993), based on a series of weak outbursts of A 0535+262 that occurred near periastron passage in the three orbits previous to the February-March outburst. The spin-up rate during the outburst is shown in panel B of Fig. 1.

Aperiodic variability in the source flux was noticed early in the outburst (Finger *et al.* 1994b). Daily average power spectra were made from the DISCLA channel 1 rates (20–50 keV, 1.024 s resolution) after subtraction of the daily mean pulse profile. The power spectrum for February 19 is shown in panel A of Fig. 2. These power spectra consist of an approximately $1/f$ power-law component extending from at least 5 mHz to 0.5 Hz, and a significant concentration of noise power in a bump centered in the 35–70 mHz range. For convenience we will call this bump in the power spectra a QPO, although the bump typically has a FWHM/center frequency of 50% or slightly greater, and therefore does not strictly meet the QPO definition used in connection with LMXBs. Typical fractional r.m.s. amplitudes of the coherent, power-law (10–500 mHz), and QPO components were 20%, 15%, and 9%, respectively. The QPO was detectable for an interval of 27 days indicated in panel A of Fig. 1. Panel B of Fig. 2 shows the history of the QPO center frequency, obtained by fitting the daily power spectra with the sum of Gaussian and power-law models. As the outburst progressed the center frequency rose smoothly from 35 mHz to 70 mHz near the peak of the outburst, and returned slowly to 35 mHz.




Figure 2. A) shows the mean power spectrum for February 19. The best fit model is superposed within the fit interval. B) shows the center frequency of the Gaussian QPO component of the power spectral model.




Figure 3. Comparison of intrinsic spin-up rate of A 0535+262 during the outburst and the QPO center frequency. Diamond symbols are used during the rise of the outburst and circles during the decline.

Fig. 3 compares the QPO center frequency and the neutron star spin up rate. For the plot, pairs of center frequencies were interpolated to the midpoint of the spin up rate measurements. The QPO frequency and the spin up rate are seen to be highly correlated. The tracks from the rise and the fall of the outburst are amazingly close, with spin up on the rise being only slightly higher than spin up on the fall at the same QPO frequency.

3. Discussion

Simple theoretical predictions of the relationship between the QPO center frequency and the spin up rate can be obtained by combining either the beat frequency model, or the Keplerian frequency model with the Ghosh & Lamb (1979) accretion torque model. The Ghosh & Lamb model gives the accretion torque as

$$N = \dot{M} \sqrt{GM r_0} n(\omega_s) \quad (1)$$

where \dot{M} is the accretion rate, M is the neutron star mass, and r_0 is the radius of the inner edge of the accretion disk. The dimensionless torque function $n(\omega_s)$ depends only on the fastness parameter ω_s which is the ratio of the neutron star spin frequency ν_{ns} to the Keplerian frequency at the inner edge of the accretion disc ν_K . The radius of the inner edge of the accretion disk is given by

$$r_0 = (GM)^{1/3} (2\pi\nu_K)^{-2/3} = \eta\mu^{4/7} (2GM)^{-1/7} \dot{M}^{-2/7} \quad (2)$$

where η is a geometry-dependent constant that Ghosh & Lamb computed to be 0.52, and μ is the neutron star magnetic moment. If we assume that the torque N acts on the solid body moment of inertia of $2/5MR^2$, the spin-up rate may be written as

$$\dot{\nu}_{\text{ns}} = \aleph n(\omega_s) \nu_K^2 \quad \text{where} \quad \aleph = \frac{5\pi\eta^{3.5}\mu^2}{\sqrt{2GM^2R^2}} \quad (3)$$

The dimensionless constant \aleph has a value of 2.1×10^{-9} for the representative values of $\eta = 1$, $\mu = 10^{31} \text{ G cm}^3$, $M = 1.4 M_\odot$, and $R = 10^6 \text{ cm}$.

A plot of $\dot{\nu}_{\text{ns}}/\nu_K^2$ versus the fastness parameter ω_s will therefore yield a measurement of $\aleph n(\omega_s)$. This is shown in Fig. 4 for both the beat frequency and Keplerian frequency models. Also shown on the plots is the theoretical relationship for several values of \aleph using the approximation for $n(\omega_s)$ given by Ghosh & Lamb (1979). For both the beat frequency model and the Keplerian frequency model the approximate scaling of $\dot{\nu}_{\text{ns}} \propto \nu_K^2$ is born out by the data. For the beat frequency model, the observations are consistent with $\aleph n(\omega_s)$ being a slowly varying function that decreases with increasing ω_s (decreasing \dot{M}), although detailed agreement between observation and model prediction is not achieved. For the Keplerian frequency model, the observed form of $\aleph n(\omega_s)$ rises at low ω_s and then falls at higher ω_s . The agreement between observations and model prediction is worse than in the beat frequency model case. The observations therefore favour the beat frequency model, for which we estimate a value of $\aleph = (2.5 \pm 0.5) \times 10^{-9}$.

For assumed values of $\eta = 1$, $R = 10^6 \text{ cm}$, and $M = 1.4 M_\odot$, we compute a polar magnetic field of $B = 2\mu/R^3 = 2 \times 10^{13} \text{ G}$. Observations with




Figure 4. The ratio of the neutron star spin-up rate to the Keplerian frequency squared for the beat frequency (panel A) and Keplerian frequency (panel B) models. The curves give the Ghosh-Lamb prediction, for values of the dimensionless constant \aleph of $2.0 \cdot 10^{-9}$ (solid), $3.0 \cdot 10^{-9}$ (dotted), and $4.0 \cdot 10^{-9}$ (dashed).

OSSE during this same outburst of A 0535+262 revealed a cyclotron line at an energy of 110 keV (Grove *et al.* 1994), or a polar magnetic field of $B = 9.5 \cdot 10^{12}$ G. Given the number of poorly known parameters, this rough agreement is encouraging.

Acknowledgements. M.H.F. would like to thank J. van Paradijs, M. van der Klis, and L. Bildsten for useful discussions.

References

- Alpar, M.A. & Shaham, J. 1985, *Nat* 316, 239
 Finger, M.H. *et al.* 1993, in *Evolution of X-ray Binaries*, S.S. Holt & C.S. Day (Eds.), AIP Conf. Proc. Vol. 408, p. 459
 Finger, M.H., Wilson, R.B. & Hagedorn, K.S. 1994a, *IAU Circ.* 5931
 Finger, M.H., Wilson, R.B. & Harmon, B.A. 1994b, *IAU Circ.* 5934
 Fishman, G.J. *et al.* 1989, *Proceedings GRO Science Workshop, GSFC*, p. 2–39
 Ghosh, P. & Lamb, F.K. 1979, *ApJ* 234, 296
 Giovannelli, F. & Graziati, L.S. 1992, *Space Sci. Rev.* 59, 1
 Grove, E. *et al.* 1994, *ApJ* (in press)
 Lamb, F.K. *et al.* 1985, *Nat* 317, 681
 Motch, C. *et al.* 1991, *ApJ* 369, 490
 Rosenberg, F.D. *et al.* 1975, *Nat* 256, 628
 Van der Klis, M. *et al.* 1987, *ApJ* 313, L19
 Wilson, R.B., Harmon, B.A. & Finger, M.H. 1994a, *IAU Circ.* 5933
 Wilson, R.B. *et al.* 1994b, *IAU Circ.* 5945

Discussion

P. Ghosh: Your determination of $\frac{d \ln \nu_K}{d \ln M}$ (which is nominally 2/7, as verified by early QPO work on LMXB) is very likely going to be the best measurement so far. This has very important implications for the state of the inner accretion disk in X-ray pulsars. The value is in agreement with the usual one-temperature and gas-pressure dominated disk, and not with some other dark models.

A. Alpar: The observation of the beat frequency does not depend on the existence of beaming or polar caps if the field is weak enough, as in LMXBs.

M. Finger: This is a HMXB with beaming and the observation of ν_{rotation} is expected.

A. Alpar: To see the inclination dependence comparatively and to discuss ν_{rotation} , we hope for QPO observations of this quality from other HMXBs.

J. van Paradijs: This is a response to the remark by P. Ghosh. One can derive relations between $\dot{\nu}$ and ν_K for different disk models. It turns out that the only disk model (from a sample of models published by P. Ghosh) that fits the approximately quadratic dependence of $\dot{\nu}$ on ν_K is that of a classical gas-pressure dominated disk; a radiation-pressure dominated disk, or a two-temperature disk, lead to very different $\dot{\nu}(\nu_K)$ relations, inconsistent with the one observed for A 0535+262.