
INSTRUCTIONS TO CONTRIBUTORS

The *Journal of Helminthology* publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance.

Manuscripts, which must be in English or French (with an English summary), should be addressed to:

The Editor, Journal of Helminthology
London School of Hygiene and Tropical Medicine
Keppel Street, London WC1E 7HT
England

Two copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief introduction (unheaded); Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration may be printed is 13.5 × 18 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space should be indicated. Some indication of scale should normally be given on the figure. Photocopies of illustrations should also be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size 13.5 × 18 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

DUKE, B. O. L. (1971) The ecology of onchocerciasis in man and animals. In *Ecology and physiology of parasites* (ed A. M. Fallis) pp. 213-222. Adam Hilger Ltd.: London.

JAMES, C. and WEBBE, G. (1973) A comparison of Egyptian and East African strains of *Schistosoma haematobium*. *Journal of Helminthology*, **47**, 49-59.

50 offprints are provided free of charge; additional copies may be ordered at the proof stage.

Contents

	Pages
BEHNKE, J. M. and WAKELIN, D. <i>Nematospiroides dubius</i> : stimulation of acquired immunity in inbred strains of mice	167—176
DARGIE, J. D., BERRY, C. I., HOLMES, P. H., TAYLOR, M. G., JAMES, E. R. and NELSON, G. S. Studies on the pathogenesis of a strain of <i>Schistosoma mattheei</i> maintained in hamsters	177—188
EVANS, N. A. The occurrence of <i>Sphaerostoma bramae</i> (Digenea: Allocreadiidae) in the roach from the Worcester—Birmingham canal	189—196
EVANS, N. A. The site preferences of two digeneans, <i>Asymphylodora kubanicum</i> and <i>Sphaerostoma bramae</i> , in the intestine of the roach	197—204
PALMIERI, J. R. and SULLIVAN, J. T. <i>Mesocoelium malayanum</i> sp. n. (Digenea: Brachycoeliidae) in a frog, <i>Rana macrodon</i> Dumeril and Bibron 1941, from Malaysia	205—208
GIBBONS, L. M. and KHALIL, L. F. Research Note. <i>Longistrongylus curvispiculum</i> (Gibbons, 1973) (Nematoda: Trichostrongylidae) in <i>Oryx tao</i> in England	209
WEST, B., WILSON, P. and HATCH, C. Research Note. <i>Aelurostrongylus abstrusus</i> in the Cheetah	210—211
KAJUBIRI, V. and HOHORST, W. Research Note. Increasing incidence of <i>Dicrocoelium hospes</i> (Looss, 1907) (Trematoda Digenea) in Uganda	212—214
CHERNIN, J. Common host antigens in laboratory rats infected with the metacestodes of <i>Taenia crassiceps</i>	215—220
RICKARD, M. D., MACKINLAY, L. M., KANE, G. J., MATOSSIAN, R. M. and SMYTH, J. D. Studies on the mechanism of lysis of <i>Echinococcus granulosus</i> protoscoleces incubated in normal serum	221—228
SUPPLEMENTARY REVIEW ARTICLES	
PITCHFORD, R. J. A check list of definitive hosts exhibiting evidence of the genus <i>Schistosoma</i> Weinland, 1858 acquired naturally in Africa and the Middle East	229—252
SPRENT, J. F. A. Ascaridoid nematodes of amphibians and reptiles: <i>Dujardinascaris</i>	253—287