

## VERY SLOW GROW-UP OF SOLUTIONS OF A SEMI-LINEAR PARABOLIC EQUATION

MAREK FILA<sup>1</sup>, JOHN R. KING<sup>2</sup>, MICHAEL WINKLER<sup>3</sup> AND EIJI YANAGIDA<sup>4\*</sup>

<sup>1</sup>*Department of Applied Mathematics and Statistics, Comenius University,  
84248 Bratislava, Slovakia (fila@fmph.uniba.sk)*

<sup>2</sup>*Division of Theoretical Mechanics, University of Nottingham,  
Nottingham NG7 2RD, UK (etzjrk@maths.nottingham.ac.uk)*

<sup>3</sup>*Fachbereich Mathematik, Universität Duisburg-Essen,  
45117 Essen, Germany (michael.winkler@uni-due.de)*

<sup>4</sup>*Mathematical Institute, Tohoku University, Sendai 980-8578,  
Japan (yanagida@math.tohoku.ac.jp)*

(Received 13 October 2009)

*Abstract* We consider large-time behaviour of global solutions of the Cauchy problem for a parabolic equation with a supercritical nonlinearity. It is known that the solution is global and unbounded if the initial value is bounded by a singular steady state and decays slowly. In this paper we show that the grow-up of solutions can be arbitrarily slow if the initial value is chosen appropriately.

*Keywords:* grow-up; semi-linear parabolic equation; comparison principle

2010 *Mathematics subject classification:* Primary 35K57  
Secondary 35B40

### 1. Introduction

This paper is a continuation of our research project on the large-time behaviour of global classical solutions of the Cauchy problem

$$\left. \begin{aligned} u_t &= \Delta u + u^p, & x \in \mathbb{R}^N, t > 0, \\ u(x, 0) &= u_0(x), & x \in \mathbb{R}^N, \end{aligned} \right\} \quad (1.1)$$

where we assume that  $u_0$  is continuous,  $N \geq 11$  and

$$p > p_c := \frac{(N-2)^2 - 4N + 8\sqrt{N-1}}{(N-2)(N-10)}.$$

\* Present address: Department of Mathematics, Tokyo Institute of Technology, Meguro-ku, Tokyo 152-8551, Japan.

Problem (1.1) has been studied as a typical super-linear problem and as a canonical problem of more general super-linear equations after taking a scaling limit. In spite of its simple appearance, (1.1) is known to have a rich mathematical structure and has been studied extensively by many authors. The exponent  $p_c$  appeared for the first time in [13] and recent studies have revealed that it is an important critical exponent for the dynamics of solutions (see [17] and the references therein).

So far, we have studied grow-up [1, 2, 4, 5], the convergence of solutions to regular steady states [6, 12], the decay to the trivial solution [3, 7] and the convergence to self-similar solutions [8]. For some previous related results we refer the reader to [9–11, 20]. It is shown in [15] that the solution of (1.1) exists globally in time but becomes unbounded if the initial value satisfies

$$0 \leq u_0(x) \leq \varphi_\infty(|x|) := L|x|^{-m}, \quad x \in \mathbb{R}^N \setminus \{0\}, \quad (1.2)$$

and

$$|x|^{m+\lambda_1} |\varphi_\infty(|x|) - u_0(x)| \rightarrow 0 \quad \text{as } |x| \rightarrow \infty, \quad (1.3)$$

where

$$m = \frac{2}{p-1}, \quad L = \{m(N-2-m)\}^{1/(p-1)}$$

and  $\lambda_1$  is the smaller positive root of

$$\lambda^2 - (N-2-2m)\lambda + 2(N-2-m) = 0. \quad (1.4)$$

We note that this equation has two distinct positive roots if  $p > p_c$ .

In our previous papers [1, 5], given a specific decay rate of  $u_0$  as  $|x| \rightarrow \infty$ , we determined the exact grow-up rate of solutions. More precisely, if the initial value satisfies (1.2) and

$$c_1|x|^{-l} < \varphi_\infty(|x|) - u_0(x) < c_2|x|^{-l}, \quad |x| > R,$$

with some positive constants  $c_1, c_2, R$  and  $l \in (m + \lambda_1, m + \lambda_2 + 2)$ , where  $\lambda_2$  is the larger positive root of (1.4), then the solution of (1.1) satisfies

$$C_1(t+1)^{m(l-m-\lambda_1)/2\lambda_1} < \|u(\cdot, t)\|_{L^\infty} < C_2(t+1)^{m(l-m-\lambda_1)/2\lambda_1}, \quad t > 0, \quad (1.5)$$

with some positive constants  $C_1, C_2$  (see [2] for the critical case  $p = p_c$  and [14] for the optimality of this result, and see also [16] for other types of global unbounded solutions).

In particular, (1.5) shows that, in (1.1), arbitrarily slow grow-up occurs *in terms of algebraic rates*: as the deviation of  $u_0$  from the steady state  $\varphi_\infty$  approaches the critical spatial decay rate  $|x|^{-m-\lambda_1}$ , the temporal growth of the corresponding solution takes place at arbitrarily small positive powers of  $t$ . We investigate whether grow-up can occur at even smaller rates than any positive power. Accordingly, we assume that the initial value satisfies (1.2) and

$$b_1|x|^{-m-\lambda_1}\omega(|x|) < \varphi_\infty(|x|) - u_0(x) < b_2|x|^{-m-\lambda_1}\omega(|x|), \quad |x| > R, \quad (1.6)$$

with some positive constants  $b_1, b_2$  and  $R$ . Here  $\omega \in C^2([0, \infty))$  is a function satisfying

$$\omega(z) > 0, \quad \omega'(z) < 0 \quad \text{and} \quad \omega''(z) \geq 0 \quad \text{for all } z \geq 0, \tag{1.7}$$

and representing slow decay at infinity in the sense that

$$\frac{z\omega'(z)}{\omega(z)} \rightarrow 0 \quad \text{as } z \rightarrow \infty. \tag{1.8}$$

Moreover, for technical reasons we will also require the regularity property

$$\left| \frac{z\omega''(z)}{\omega'(z)} \right| \leq C_\omega \quad \text{for all } z \geq 0 \tag{1.9}$$

with some constant  $C_\omega > 0$ . Note that, as a consequence of (1.8) and (1.9), we also see that

$$\frac{z^2\omega''(z)}{\omega(z)} \rightarrow 0 \quad \text{as } z \rightarrow \infty. \tag{1.10}$$

Under the above assumptions, the initial value satisfies (1.2) and (1.3) so that the solution of (1.1) is global and unbounded in time.

The main result of this paper is as follows.

**Theorem 1.1.** *Let  $N \geq 11$  and  $p > p_c$ . Suppose that the initial value satisfies (1.2) and (1.6). Then the solution of (1.1) satisfies*

$$C_1\omega^{-m/\lambda_1}(t^{1/2}) \leq \|u(\cdot, t)\|_{L^\infty} \leq C_2\omega^{-m/\lambda_1}(t^{1/2}) \quad \text{for all } t > 0.$$

with some constants  $C_1, C_2 > 0$ .

This theorem implies that the solution grows up arbitrarily slowly if  $u_0$  is chosen appropriately. For example, the function

$$\omega(z) = [\log(\log(\cdots(\log(z + z_0))\cdots))]^{-\alpha}, \quad \alpha > 0,$$

satisfies our assumptions if  $z_0 > 0$  is sufficiently large.

After the first draft of this paper was completed, our result was extended in [18] to very slow convergence to zero and in [19] to very slow convergence to positive steady states.

This paper is organized as follows. In §2 we give a lower bound of radial solutions by constructing a suitable subsolution. In §3 we give an upper bound of radial solutions by constructing a suitable super-solution. In §4 we prove Theorem 1.1 by using these estimates for radial solutions. In the following sections, we assume  $N > 10$  and  $p > p_c$  throughout.

2. Lower bound

In this section and the next we consider radially symmetric solutions  $u = u(r, t)$ ,  $r := |x|$ , of (1.1). Then we may write (1.1) as

$$\left. \begin{aligned} u_t &= u_{rr} + \frac{N-1}{r}u_r + u^p, & r > 0, t > 0, \\ u(r, 0) &= u_0(r), & x \in \mathbb{R}^N, \end{aligned} \right\} \tag{2.1}$$

where  $u_0(r)$  is assumed to satisfy (1.2) and (1.6). We shall construct a subsolution of (2.1) that inherits the asymptotic behaviour of the initial value, at least in an *outer domain* that will be specified by an inequality of the form  $r \geq B(T + 1)^{1/2}$  with  $B > 0$  in Corollary 2.2.

**Lemma 2.1.** *For any  $\theta \in (0, \min\{1, \frac{1}{2}(\lambda_2 - \lambda_1)\})$  and  $b_1 > 0$ , there exists  $b_2 > 0$  such that*

$$u_{\text{out}}^-(r, t) := \max\{0, Lr^{-m} - b_1r^{-m-\lambda_1}\omega(r) - b_2r^{-m-\lambda_1-2\theta}\omega(r)(t + 1)^\theta\}$$

defines a subsolution of (2.1) for all  $r \geq 0$  and  $t \geq 0$ .

**Proof.** Let  $\theta \in (0, \min\{1, \frac{1}{2}(\lambda_2 - \lambda_1)\})$  and  $b_1 > 0$  be given, and fix  $\delta > 0$  such that

$$0 < \delta \leq \frac{\theta(\lambda_2 - \lambda_1 - 2\theta)}{|N - 1 - 2m - 2\lambda_1 - 4\theta| + 1}. \tag{2.2}$$

In view of (1.8) and (1.10), we may choose  $z_0 > 0$  so large that

$$\left| \frac{z\omega'(z)}{\omega(z)} \right| \leq \delta \quad \text{and} \quad \left| \frac{z^2\omega''(z)}{\omega(z)} \right| \leq \delta \quad \text{for all } z \geq z_0. \tag{2.3}$$

We now take  $b_2 > 0$  such that

$$b_2 \geq \frac{Lz_0^{\lambda_1+2\theta}}{\omega(z_0)} \tag{2.4}$$

and

$$b_2 \geq \frac{b_1(|N - 1 - 2m - 2\lambda_1| + 1)\delta}{\min\{\theta, \theta(\lambda_2 - \lambda_1 - 2\theta)\}}. \tag{2.5}$$

Then, at each point from the positivity set

$$S := \{(r, t) \in [0, \infty)^2 \mid u_{\text{out}}^-(r, t) > 0\}$$

of  $u_{\text{out}}^-$ , we have

$$Lr^{-m} > b_2r^{-m-\lambda_1-2\theta}\omega(r)(t + 1)^\theta \geq b_2r^{-m-\lambda_1-2\theta}\omega(r),$$

and hence, by (2.4),

$$\frac{r^{\lambda_1+2\theta}}{\omega(r)} > \frac{b_2}{L} > \frac{z_0^{\lambda_1+2\theta}}{\omega(z_0)}.$$

Since  $r \mapsto r^{\lambda_1+2\theta}/\omega(r)$  is strictly increasing on  $(0, \infty)$  in view of (1.7), this implies

$$r > z_0 \quad \text{for all } (r, t) \in S. \tag{2.6}$$

Moreover, if  $(r, t) \in S$ , then

$$\begin{aligned} & \mathcal{P}u_{\text{out}}^- \\ & := (u_{\text{out}}^-)_t - (u_{\text{out}}^-)_{rr} - \frac{N-1}{r}(u_{\text{out}}^-)_r - (u_{\text{out}}^-)^p \\ & = -b_2\theta r^{-m-\lambda_1-2\theta}\omega(r)(t+1)^{\theta-1} \\ & \quad - \left\{ (Lr^{-m})_{rr} + \frac{N-1}{r}(Lr^{-m})_r \right\} \\ & \quad + b_1 \left\{ (r^{-m-\lambda_1}\omega(r))_{rr} + \frac{N-1}{r}(r^{-m-\lambda_1}\omega(r))_r \right\} \\ & \quad + b_2 \left\{ (r^{-m-\lambda_1-2\theta}\omega(r))_{rr} + \frac{N-1}{r}(r^{-m-\lambda_1-2\theta}\omega(r))_r \right\} (t+1)^\theta - (u_{\text{out}}^-)^p \\ & = -b_2\theta r^{-m-\lambda_1-2\theta}\omega(r)(t+1)^{\theta-1} + (Lr^{-m})^p \\ & \quad + b_1 \{ (m+\lambda_1)(m+\lambda_1+2-N)r^{-m-\lambda_1-2}\omega(r) \\ & \quad \quad + (N-1-2m-2\lambda_1)r^{-m-\lambda_1-1}\omega'(r) + r^{-m-\lambda_1}\omega''(r) \} \\ & \quad + b_2 \{ (m+\lambda_1+2\theta)(m+\lambda_1+2\theta+2-N)r^{-m-\lambda_1-2\theta-2}\omega(r) \\ & \quad \quad + (N-1-2m-2\lambda_1-4\theta)r^{-m-\lambda_1-2\theta-1}\omega'(r) + r^{-m-\lambda_1-2\theta}\omega''(r) \} (t+1)^\theta \\ & \quad \quad \quad - (u_{\text{out}}^-)^p. \end{aligned} \tag{2.7}$$

By the convexity of  $z \mapsto (1-z)^p$  for  $z < 1$ , we have, using  $(p-1)m = 2$  and  $pL^{p-1} = (m+2)(N-2-m)$ ,

$$\begin{aligned} (u_{\text{out}}^-)^p & \geq (Lr^{-m})^p - pL^{p-1}r^{-(p-1)m}[b_1r^{-m-\lambda_1}\omega(r) - b_2r^{-m-\lambda_1-2\theta}\omega(r)(t+1)^\theta] \\ & = (Lr^{-m})^p - b_1(m+2)(N-2-m)r^{-m-\lambda_1-2}\omega(r) \\ & \quad - b_2(m+2)(N-2-m)r^{-m-\lambda_1-2\theta-2}\omega(r)(t+1)^\theta \end{aligned}$$

for all  $(r, t) \in S$ . Therefore, for all  $(r, t) \in S$ ,

$$\begin{aligned} & \mathcal{P}u_{\text{out}}^- \\ & \leq -b_2\theta r^{-m-\lambda_1-2\theta}\omega(r)(t+1)^{\theta-1} \\ & \quad + b_1 \{ [(m+\lambda_1)(m+\lambda_1+2-N) + (m+2)(N-2-m)]r^{-m-\lambda_1-2}\omega(r) \\ & \quad \quad + (N-1-2m-2\lambda_1)r^{-m-\lambda_1-1}\omega'(r) + r^{-m-\lambda_1}\omega''(r) \} \\ & \quad + b_2 \{ [(m+\lambda_1+2\theta)(m+\lambda_1+2\theta+2-N) \\ & \quad \quad + (m+2)(N-2-m)]r^{-m-\lambda_1-2\theta-2}\omega(r) \\ & \quad \quad + (N-1-2m-2\lambda_1-4\theta)r^{-m-\lambda_1-2\theta-1}\omega'(r) + r^{-m-\lambda_1-2\theta}\omega''(r) \} (t+1)^\theta. \end{aligned}$$

Here we observe that, by the definition of  $\lambda_1$ ,

$$\begin{aligned} (m + \lambda_1)(m + \lambda_1 + 2 - N) + (m + 2)(N - 2 - m) \\ = \lambda_1^2 - (N - 2 - 2m)\lambda_1 + 2(N - 2 - m) \\ = 0 \end{aligned}$$

and, consequently,

$$\begin{aligned} (m + \lambda_1 + 2\theta)(m + \lambda_1 + 2\theta + 2 - N) + (m + 2)(N - 2 - m) \\ = 2\theta(m + \lambda_1 + 2\theta + 2 - N) + (m + \lambda_1)2\theta \\ = 2\theta(2m + 2\lambda_1 + 2\theta + 2 - N) \\ = 2\theta[2\theta - (\lambda_2 - \lambda_1)], \end{aligned}$$

where we have used the equalities

$$2m + 2\lambda_1 = N - 2 - \sqrt{(N - 2 - 2m)^2 - 2(N - 2 - m)}$$

and

$$\lambda_2 - \lambda_1 = \sqrt{(N - 2 - 2m)^2 - 2(N - 2 - m)}.$$

Accordingly,

$$\begin{aligned} \mathcal{P}u_{\text{out}}^- \\ \leq -b_2\theta r^{-m-\lambda_1-2\theta}\omega(r)(t+1)^{\theta-1} \\ + b_1\{(N-1-2m-2\lambda_1)r^{-m-\lambda_1-1}\omega'(r) + r^{-m-\lambda_1}\omega''(r)\} \\ + b_2\{-2\theta(\lambda_2-\lambda_1-2\theta)r^{-m-\lambda_1-2\theta-2}\omega(r) \\ + (N-1-2m-2\lambda_1-4\theta)r^{-m-\lambda_1-2\theta-1}\omega'(r) + r^{-m-\lambda_1-2\theta}\omega''(r)\}(t+1)^\theta \\ = b_2r^{-m-\lambda_1-2\theta}\omega(r)(t+1)^{\theta-1} \\ \times \left\{ -\theta + \frac{b_1}{b_2} \left[ (N-1-2m-2\lambda_1)\frac{r\omega'(r)}{\omega(r)} + \frac{r^2\omega''(r)}{\omega(r)} \right] \left( \frac{t+1}{r^2} \right)^{1-\theta} \right. \\ \left. - 2\theta(\lambda_2-\lambda_1-2\theta)\frac{t+1}{r^2} \right. \\ \left. + \left[ (N-1-2m-2\lambda_1-4\theta)\frac{r\omega'(r)}{\omega(r)} + \frac{r^2\omega''(r)}{\omega(r)} \right] \frac{t+1}{r^2} \right\} \end{aligned}$$

for all  $(r, t) \in S$ . Using the trivial estimate

$$\left( \frac{t+1}{r^2} \right)^{1-\theta} \leq \max \left\{ 1, \frac{t+1}{r^2} \right\} \leq 1 + \frac{t+1}{r^2}$$

and recalling (2.6), we obtain from (2.3) that

$$\begin{aligned} \mathcal{P}u_{\text{out}}^- &\leq b_2 r^{-m-\lambda_1-2\theta} \omega(r) (t+1)^{\theta-1} \\ &\quad \times \left\{ -\theta + \frac{b_1}{b_2} [N-1-2m-2\lambda_1|\delta+\delta] \left(1 + \frac{t+1}{r^2}\right) \right. \\ &\quad \left. - 2\theta(\lambda_2 - \lambda_1 - 2\theta) \frac{t+1}{r^2} + [N-1-2m-2\lambda_1-4\theta|\delta+\delta] \frac{t+1}{r^2} \right\} \\ &= b_2 r^{-m-\lambda_1-2\theta} \omega(r) (t+1)^{\theta-1} \\ &\quad \times \left\{ -\theta + \frac{b_1}{b_2} [N-1-2m-2\lambda_1+1]\delta \right. \\ &\quad \left. - \left( 2\theta(\lambda_2 - \lambda_1 - 2\theta) - \frac{b_1}{b_2} [N-1-2m-2\lambda_1+1]\delta \right. \right. \\ &\quad \left. \left. - [N-1-2m-2\lambda_1-4\theta+1]\delta \right) \frac{t+1}{r^2} \right\}, \end{aligned}$$

and hence, in view of (2.2) and (2.5), we conclude that  $\mathcal{P}u_{\text{out}}^- < 0$  for  $(r, t) \in S$ . Since  $u \equiv 0$  is evidently a subsolution, this completes the proof.  $\square$

**Corollary 2.2.** *Suppose that*

$$u_0(r) \geq Lr^{-m} - b_- r^{-m-\lambda_1} \omega(r) \quad \text{for all } r > 0 \tag{2.8}$$

*holds with some  $b_- > 0$ . Then, for all  $B > 0$ , there exists  $b_0 > 0$  such that the solution  $u$  of (2.1) satisfies*

$$u(r, t) \geq Lr^{-m} - b_0 r^{-m-\lambda_1} \omega(r) \quad \text{for all } t \geq 0 \text{ and } r \geq B(t+1)^{1/2}. \tag{2.9}$$

**Proof.** We apply Lemma 2.1 to  $b_1 := b_-$  and any  $\theta \in (0, \min\{1, \frac{1}{2}(\lambda_2 - \lambda_1)\})$  to obtain some  $b_2 > 0$  such that  $u_{\text{out}}^-$  as given in Lemma 2.1 is a subsolution of (2.1). Our lower estimate (2.8) for  $u_0$ , in conjunction with the fact that  $u_0$  is non-negative, implies that  $u_{\text{out}}^-(r, 0) \leq u_0(r)$  for all  $r \geq 0$ . Therefore, the maximum principle shows that  $u_{\text{out}}^- \geq u$  for all  $r \geq 0$  and  $t \geq 0$ . In particular, if  $B > 0$  is given, then, for all  $t \geq 0$  and  $r \leq B(t+1)^{1/2}$ , we find

$$\begin{aligned} u(r, t) &\geq u_{\text{out}}^-(r, t) \\ &\geq Lr^{-m} - b_1 r^{-m-\lambda_1} \omega(r) - b_2 r^{-m-\lambda_1-2\theta} \omega(r) (t+1)^\theta \\ &\geq Lr^{-m} - b_1 r^{-m-\lambda_1} \omega(r) - b_2 B^{-2\theta} r^{-m-\lambda_1} \omega(r), \end{aligned}$$

which proves (2.9).  $\square$

We proceed to derive an estimate from below in a corresponding inner region. In preparation, let us recall some facts about the solutions  $\psi$  and  $\Psi$  of the initial-value problems

$$\left. \begin{aligned} \psi_{\xi\xi} + \frac{N-1}{\xi} \psi_\xi + \psi^p &= 0, \quad \xi > 0, \\ \psi(0) &= 1, \quad \psi_\xi(0) = 0 \end{aligned} \right\} \tag{2.10}$$

and

$$\left. \begin{aligned} \Psi_{\xi\xi} + \frac{N-1}{\xi}\Psi_{\xi} + p\psi^{p-1}\Psi &= \psi + \frac{1}{m}\xi\psi_{\xi} + \chi(\xi), \quad a\xi > 0, \\ \Psi(0) &= -1, \quad \Psi_{\xi}(0) = 0, \end{aligned} \right\} \quad (2.11)$$

respectively, where  $\chi(\xi) := 1/(1 + \xi^{-m-\lambda_1})$ . More specifically, it is known [5] that there exist  $a_1 > 0$  and  $K > 0$  such that

$$\psi(\xi) \simeq L\xi^{-m} - a_1\xi^{-m-\lambda_1}, \quad (2.12)$$

$$\Psi(\xi) \simeq K\xi^{2-m-\lambda_1}, \quad (2.13)$$

$$\Psi_{\xi}(\xi) \simeq (2 - m - \lambda_1)K\xi^{1-m-\lambda_1}, \quad (2.14)$$

as  $\xi \rightarrow \infty$ . In fact, in what follows we shall refer neither to the prescribed explicit value of  $\Psi(0)$  nor to the precise form of  $\chi$  as introduced above, for which (2.13) and (2.14) were proved in [5]. Both formulae would remain unchanged for any value of  $\Psi(0)$  and any smooth positive decreasing  $\chi$  satisfying  $\xi^{m+\lambda_1}\chi(\xi) \rightarrow A \geq 0$  as  $\xi \rightarrow \infty$ .

**Lemma 2.3.** *Fix an arbitrary  $\kappa > 2/m$ . Then there exists  $\mu_0 > 0$  such that if*

$$\sigma(t) := \varepsilon\omega^{-m/\lambda_1}((t + \mu^{-\kappa})^{1/2}), \quad \varepsilon := \mu\omega^{m/\lambda_1}(\mu^{-\kappa/2}), \quad r \geq 0, \quad t \geq 0, \quad (2.15)$$

with

$$\xi(r, t) := \sigma^{1/m}(t)r, \quad r \geq 0, \quad t \geq 0, \quad (2.16)$$

and some  $\mu < \mu_0$ , then

$$u_{\text{in}}^-(r, t) := \max \left\{ 0, \sigma \left( \psi(\xi) + \frac{\sigma_t}{\sigma^p} \Psi(\xi) \right) \right\} \quad (2.17)$$

defines a subsolution of (2.1) for all  $r \geq 0$  and  $t \geq 0$ .

**Proof.** Since  $u \equiv 0$  is a subsolution, we only need to consider those points where  $u_{\text{in}}^-$  is positive.

By (2.13) and (2.14), there exists  $\xi_0 > 0$  such that

$$\Psi(\xi) \geq 0 \quad \text{and} \quad \Psi_{\xi}(\xi) \leq 0 \quad \text{for all } \xi \geq \xi_0. \quad (2.18)$$

Then

$$\Psi(\xi) \leq C \quad \text{and} \quad |\xi\Psi_{\xi}(\xi)| \leq C \quad \text{for all } \xi \geq \xi_0 \quad (2.19)$$

with some  $C > 0$ . Next we take  $\delta > 0$  so small that

$$\frac{m + \lambda_1 - 2}{\lambda_1} \omega^{2/\lambda_1}(0)\delta \leq 1 \quad (2.20)$$

and then, according to (1.8), we take  $z_0$  large with the property that

$$\left| \frac{z\omega'(z)}{\omega(z)} \right| \leq \delta \quad \text{for all } z \geq z_0. \quad (2.21)$$



We finally fix  $\mu_0 > 0$  small enough to satisfy

$$\mu_0 \leq z_0^{-2/\kappa} \tag{2.22}$$

and

$$\mu_0 \leq \left( \frac{\chi(\xi_0)}{C[(m + \lambda_1 - 1)/2\lambda_1]\delta + \frac{1}{2} + \frac{1}{2}C_\omega} \right). \tag{2.23}$$

With these choices of constants, we take  $\mu < \mu_0$  and let  $u_{\text{in}}^-$  be defined by (2.17). Regarding  $\mathcal{P}u_{\text{in}}^-$  with  $\mathcal{P}$  as defined in the proof of Lemma 2.1, it can easily be checked using the convexity of  $z \mapsto z^p$  for  $z > 0$  that, at each point where  $u_{\text{in}}^-$  is positive, we have

$$\begin{aligned} \mathcal{P}u_{\text{in}}^- &= \sigma_t \left( \psi + \frac{1}{m} \xi \psi_\xi \right) + \left( \frac{\sigma_t}{\sigma^{p-1}} \Psi \right)_t \\ &\quad - \sigma_t \left( \Psi_{\xi\xi} + \frac{N-1}{\xi} \Psi_\xi \right) - \sigma^p \left\{ \left( \psi + \frac{\sigma_t}{\sigma^p} \Psi \right)^p - \psi^p \right\} \\ &\leq \left( \frac{\sigma_t}{\sigma^{p-1}} \Psi \right)_t + \sigma_t \left( -\Psi_{\xi\xi} - \frac{N-1}{\xi} \Psi_\xi - p\psi^{p-1}\Psi + \psi + \frac{1}{m} \xi \psi_\xi \right) \\ &= \left( \frac{\sigma_t}{\sigma^{p-1}} \Psi(\xi) \right)_t - \sigma_t \chi(\xi). \end{aligned}$$

Suppressing the argument  $(t + \mu^{-\kappa})^{1/2}$  in  $\omega$ , we compute

$$\begin{aligned} \sigma_t &= -\frac{m}{2\lambda_1} \varepsilon \omega^{(-m-\lambda_1)/\lambda_1} \omega'(t + \mu^{-\kappa})^{-1/2}, \\ \frac{\sigma_t}{\sigma^{p-1}} &= -\frac{m}{2\lambda_1} \varepsilon^{(m-2)/m} \omega^{(-m-\lambda_1+2)/\lambda_1} \omega'(t + \mu^{-\kappa})^{-1/2}, \\ \frac{\sigma_t^2}{\sigma^p} &= \frac{m^2}{4\lambda_1^2} \varepsilon^{(m-2)/m} \omega^{(-m-2\lambda_1+2)/\lambda_1} \omega'^2(t + \mu^{-\kappa})^{-1}. \end{aligned}$$

Hence, using

$$\xi = \frac{1}{m} \sigma^{1/m-1} \sigma_t r = \frac{1}{m} \frac{\xi \sigma_t}{\sigma},$$

we obtain that

$$\begin{aligned} \mathcal{P}u_{\text{in}}^- &\leq \left( \frac{\sigma_t}{\sigma^{p-1}} \right)_t \Psi(\xi) + \frac{1}{m} \frac{\sigma_t^2}{\sigma^p} \xi \Psi_\xi + \frac{m}{2\lambda_1} \varepsilon \omega^{(-m-\lambda_1)/\lambda_1} \omega'(t + \mu^{-\kappa})^{-1/2} \chi(\xi) \\ &= \varepsilon^{(m-2)/m} \left\{ \frac{m(m + \lambda_1 - 2)}{4\lambda_1^2} \omega^{(-m-2\lambda_1+2)/\lambda_1} \omega'^2(t + \mu^{-\kappa})^{-1} \right. \\ &\quad \left. + \frac{m}{4\lambda_1} \omega^{(-m-\lambda_1+2)/\lambda_1} \omega'(t + \mu^{-\kappa})^{-3/2} - \frac{m}{4\lambda_1} \omega^{(-m-\lambda_1+2)/\lambda_1} \omega''(t + \mu^{-\kappa})^{-1} \right\} \Psi(\xi) \\ &\quad + \frac{m}{4\lambda_1^2} \varepsilon^{(m-2)/m} \omega^{(-m-2\lambda_1+2)/\lambda_1} \omega'^2(t + \mu^{-\kappa})^{-1} \xi \Psi_\xi(\xi) \\ &\quad + \frac{m}{2\lambda_1} \varepsilon \omega^{(-m-\lambda_1)/\lambda_1} \omega'(t + \mu^{-\kappa})^{-1/2} \chi(\xi). \tag{2.24} \end{aligned}$$

Now, for  $(r, t)$  such that  $\xi(r, t) \geq \xi_0$ , (2.18) in combination with the monotonicity and convexity of  $\omega$  and the positivity of  $\chi$  implies that

$$\mathcal{P}u_{\text{in}}^- \leq \varepsilon^{(m-2)/m} \left\{ \frac{m(m + \lambda_1 - 2)}{4\lambda_1^2} \omega^{(-m-2\lambda_1+2)/\lambda_1} \omega'^2 (t + \mu^{-\kappa})^{-1} + \frac{m}{4\lambda_1} \omega^{(-m-\lambda_1+2)/\lambda_1} \omega' (t + \mu^{-\kappa})^{-3/2} \right\} \Psi(\xi).$$

Here, in view of (2.22), we have  $\mu^{-\kappa} \geq z_0^2$  and hence, by (2.21) and (2.20),

$$\begin{aligned} & \frac{[m(m + \lambda_1 - 2)/4\lambda_1^2] \omega^{(-m-2\lambda_1+2)/\lambda_1} \omega'^2 (t + \mu^{-\kappa})^{-1}}{[m/4\lambda_1] \omega^{(-m-\lambda_1+2)/\lambda_1} |\omega'| (t + \mu^{-\kappa})^{-3/2}} \\ &= \frac{m + \lambda_1 - 2}{\lambda_1} \omega^{2/\lambda_1} ((t + \mu^{-\kappa})^{1/2}) \frac{(t + \mu^{-\kappa})^{1/2} |\omega'| ((t + \mu^{-\kappa})^{1/2})}{\omega((t + \mu^{-\kappa})^{1/2})} \\ &\leq \frac{m + \lambda_1 - 2}{\lambda_1} \omega^{2/\lambda_1} (0) \delta \\ &\leq 1, \end{aligned}$$

which yields

$$\mathcal{P}u_{\text{in}}^- \leq 0 \quad \text{if } u_{\text{in}}^-(r, t) > 0 \text{ and } \xi(r, t) \geq \xi_0. \quad (2.25)$$

On the other hand, if  $\xi < \xi_0$ , then, due to (2.24), (2.19) and the fact that  $\omega$  and  $\chi$  are decreasing, we have

$$\begin{aligned} & \frac{\mathcal{P}u_{\text{in}}^-}{\varepsilon \omega^{(-m-\lambda_1)/\lambda_1} |\omega'| (t + \mu^{-\kappa})^{-1/2}} \\ &\leq \varepsilon^{-2/m} \left\{ \frac{m(m + \lambda_1 - 2)}{4\lambda_1^2} \omega^{(-\lambda_1+2)/\lambda_1} |\omega'| (t + \mu^{-\kappa})^{-1/2} + \frac{m}{4\lambda_1} \omega^{2/\lambda_1} (t + \mu^{-\kappa})^{-1} + \frac{m}{4\lambda_1} \omega^{2/\lambda_1} \frac{|\omega''|}{|\omega'|} (t + \mu^{-\kappa})^{-1/2} \right\} C \\ &\quad + \frac{m}{4\lambda_1^2} \varepsilon^{-2/m} \omega^{(-\lambda_1+2)/\lambda_1} |\omega'| (t + \mu^{-\kappa})^{-1/2} C - \frac{m}{2\lambda_1} \chi(\xi_0) \\ &= C \mu^{-2/m} \omega^{-2/\lambda_1} (\mu^{-\kappa/2}) \omega^{2/\lambda_1} ((t + \mu^{-\kappa})^{1/2}) (t + \mu^{-\kappa})^{1/2} \\ &\quad \times \left\{ \frac{m(m + \lambda_1 - 2)}{4\lambda_1^2} \left| \frac{(t + \mu^{-\kappa})^{1/2} \omega'((t + \mu^{-\kappa})^{1/2})}{\omega((t + \mu^{-\kappa})^{1/2})} \right| + \frac{m}{4\lambda_1} + \frac{m}{4\lambda_1} \left| \frac{(t + \mu^{-\kappa})^{1/2} \omega''((t + \mu^{-\kappa})^{1/2})}{\omega'((t + \mu^{-\kappa})^{1/2})} \right| + \frac{m}{4\lambda_1^2} \left| \frac{(t + \mu^{-\kappa})^{1/2} \omega'((t + \mu^{-\kappa})^{1/2})}{\omega((t + \mu^{-\kappa})^{1/2})} \right| \right\} - \frac{m}{2\lambda_1} \chi(\xi_0) \\ &\leq C \mu^{\kappa-2/m} \left\{ \frac{m(m + \lambda_1 - 2)}{4\lambda_1^2} \delta + \frac{m}{4\lambda_1} + \frac{m}{4\lambda_1} C_\omega + \frac{m}{4\lambda_1^2} \delta \right\} - \frac{m}{2\lambda_1} \chi(\xi_0) \\ &\leq 0 \end{aligned}$$

by (2.23), where we also have used (2.21), (1.9) and (2.22). This proves the desired subsolution property.  $\square$

In order to compare  $u$  in a suitable inner region with one of the functions  $u_{\text{in}}^-$  that we just constructed, we need to show that  $u_{\text{in}}^- \leq u$  holds at the corresponding ‘lateral’ boundary. We prepare for this with the next lemma.

**Lemma 2.4.** *Let  $\kappa > 2/m$  and  $b_0 > 0$ . Then there exists  $\mu_1 > 0$  such that if  $\mu \leq \mu_1$ , then the function  $u_{\text{in}}^-$  defined in Lemma 2.3 satisfies*

$$u_{\text{in}}^- \leq Lr^{-m} - b_0r^{-m-\lambda_1}\omega(r) \quad \text{for all } (r, t) \in P,$$

where

$$P := \{(r, t) \in [0, \infty)^2 \mid r = (t + \mu^{-\kappa})^{1/2}\}.$$

**Proof.** According to (2.12) and (2.13), we can find large  $\xi_1 > 0$  such that

$$\psi(\xi) \leq L\xi^{-m} - \frac{1}{2}a_1\xi^{-m-\lambda_1} \quad \text{for all } \xi \geq \xi_1 \tag{2.26}$$

and

$$\Psi(\xi) \leq 2K\xi^{2-m-\lambda_1} \quad \text{for all } \xi \geq \xi_1. \tag{2.27}$$

With large  $z_1 > 0$  such that

$$\left| \frac{z\omega'(z)}{\omega(z)} \right| \leq \frac{a_1\lambda_1}{4Km} \quad \text{for all } z \geq z_1, \tag{2.28}$$

we let  $\mu_1 > 0$  be so small that

$$\mu_1 \leq \xi_1^{-2/(\kappa-2/m)}, \tag{2.29}$$

$$\mu_1 \leq z_1^{-2/\kappa} \tag{2.30}$$

and

$$\mu_1 \leq \left( \frac{a_1\omega(0)}{4b_0} \right)^{m/\lambda_1}. \tag{2.31}$$

Then, for any  $\mu \leq \mu_1$ , (2.29) guarantees that if  $t \geq 0$  and  $r = (t + \mu^{-\kappa})^{1/2}$ , then  $\xi$  as given by (2.16) and (2.15) satisfies

$$\begin{aligned} \xi(r, t) &= \sigma^{1/m}(t)r \\ &\geq \sigma^{1/m}(t)\mu^{-\kappa/2} \\ &= \mu^{1/m}\omega^{1/\lambda_1}(\mu^{-\kappa/2})\omega^{-1/\lambda_1}((t + \mu^{-\kappa})^{1/2})\mu^{-\kappa/2} \\ &\geq \mu^{1/m}\mu^{-\kappa/2} \\ &\geq \xi_1. \end{aligned}$$

Hence, from (2.26), (2.27) and (2.30) we obtain that, at  $r = (t + \mu^{-\kappa})^{1/2}$ ,

$$\begin{aligned} u_{\text{in}}^- &\leq \sigma \left( L\xi^{-m} - \frac{1}{2}a_1\xi^{-m-\lambda_1} + \frac{\sigma t}{\sigma^p} 2K\xi^{2-m-\lambda_1} \right) \\ &= Lr^{-m} - \frac{1}{2}a_1\sigma^{-\lambda_1/m}r^{-m-\lambda_1} + 2K\sigma^{(-m-\lambda_1)/m}\sigma_t r^{2-m-\lambda_1} \\ &= Lr^{-m} - \frac{1}{2}a_1\varepsilon^{-\lambda_1/m}\omega((t + \mu^{-\kappa})^{1/2})r^{-m-\lambda_1} \\ &\quad - 2K\frac{m}{2\lambda_1}\varepsilon^{-\lambda_1/m}\omega'((t + \mu^{-\kappa})^{1/2})(t + \mu^{-\kappa})^{-1/2} \cdot r^{2-m-\lambda_1} \\ &= Lr^{-m} - \varepsilon^{-\lambda_1/m} \left\{ \frac{1}{2}a_1 - \frac{Km}{\lambda_1} \left| \frac{r\omega'(r)}{\omega(r)} \right| \right\} r^{-m-\lambda_1}\omega(r) \\ &\leq Lr^{-m} - \frac{1}{4}a_1\varepsilon^{-\lambda_1/m}r^{-m-\lambda_1}\omega(r). \end{aligned}$$

Since

$$\varepsilon^{-\lambda_1/m} = \mu^{-\lambda_1/m}\omega^{-1}(\mu^{-\kappa/2}) \geq \mu^{-\lambda_1/m}\omega^{-1}(0)$$

due to the fact that  $\omega$  decreases on  $(0, \infty)$ , the restriction (2.31) on  $\mu_1$  yields the desired inequality. □

**Lemma 2.5.** *Suppose that  $u_0 = u_0(r)$  is continuous and positive for  $r \geq 0$  and that it satisfies*

$$u_0(r) \geq Lr^{-m} - b_-r^{-m-\lambda_1}\omega(r) \quad \text{for all } r > 0$$

with some positive constant  $b_-$ . Then there exists  $\mu_2 > 0$  such that, whenever  $\mu \leq \mu_2$ , the function  $u_{\text{in}}^-$  introduced in Lemma 2.3 satisfies

$$u_{\text{in}}^-(r, 0) \leq u_0(r) \quad \text{for all } r \in [0, \mu^{-\kappa/2}]. \tag{2.32}$$

**Proof.** In a similar way to the proof of Lemma 2.4, we first choose  $\xi_1 \geq 0$  such that (2.26) and (2.27) hold. Since  $\psi$  and  $\Psi$  are continuous and  $0 < \psi(\xi) < L\xi^{-m}$  for all  $\xi \geq 0$ , we can then fix  $C > 0$  satisfying

$$\frac{\Psi(\xi)}{\psi(\xi)} \leq C \quad \text{for all } \xi \leq \xi_1 \tag{2.33}$$

and find that

$$\nu := \frac{1}{2} \min(L - \xi^m\psi(\xi)) \tag{2.34}$$

is positive. Next we let  $r_0 > 0$  be large enough that

$$\frac{r^{\lambda_1}}{\omega(r)} \geq \frac{b_-}{\nu} \quad \text{for all } r \geq r_0 \tag{2.35}$$

and set

$$\delta := \min\{u_0(r) \mid r \leq r_0\}, \tag{2.36}$$

which is greater than zero because  $u_0$  is positive. By (1.8), we can find  $z_2 > 0$  satisfying

$$\left| \frac{z\omega'(z)}{\omega(z)} \right| \leq \min \left\{ \frac{a_1\lambda_1}{4Km}, 1 \right\} \quad \text{for all } z \geq z_2, \tag{2.37}$$

and, finally, we take  $\mu_2 > 0$  so small that

$$\mu_2 \leq z_2^{-2/\kappa}, \tag{2.38}$$

$$\mu_2 \leq \left( \frac{2\lambda_1}{mC} \frac{\nu}{L - 2\nu} \right)^{1/(\kappa - (2/m))}, \tag{2.39}$$

$$\mu_2 \leq \frac{L - 2\nu}{L - \nu} \delta, \tag{2.40}$$

$$\mu_2 \leq \left( \frac{a_1}{4b - \omega(0)} \right)^{m/\lambda_1}. \tag{2.41}$$

In deriving (2.32), we may evidently assume that  $u_{\text{in}}^-(r, 0) > 0$  and first consider those  $r \leq \mu^{-\kappa/2}$  for which  $\xi = \sigma^{1/m}(0)r = \mu^{1/m}r \geq \xi_1$  holds. At such points, from (2.26), (2.27) and (2.37) we obtain

$$\begin{aligned} u_{\text{in}}^-(r, 0) &\leq \sigma(0) \left\{ L\xi^{-m} - \frac{1}{2}a_1\xi^{-m-\lambda_1} + \frac{\sigma_t(0)}{\sigma^p(0)}\xi^{2-m-\lambda_1} \right\} \\ &= Lr^{-m} - \frac{1}{2}a_1\sigma^{-\lambda_1/m}(0)r^{-m-\lambda_1} + 2K\sigma^{(-m-\lambda_1)/\lambda_1}(0)\sigma_t(0)r^{2-m-\lambda_1} \\ &= Lr^{-m} - \frac{1}{2}a_1\varepsilon^{-\lambda_1/m}\omega(\mu^{-\kappa/2})r^{-m-\lambda_1} \\ &\quad - \frac{Km}{\lambda_1}\varepsilon^{-\lambda_1/m}\omega'(\mu^{-\kappa/2})r^{-m-\lambda_1}\mu^{\kappa/2}r^{2-m-\lambda_1} \\ &= Lr^{-m} - \mu^{-\lambda_1/m} \left\{ \frac{1}{2}a_1 - \frac{Km}{\lambda_1} \left| \frac{\mu^{-\kappa/2}\omega'(\mu^{-\kappa/2})}{\omega(\mu^{-\kappa/2})} \right| \mu^\kappa r^2 \right\} r^{-m-\lambda_1} \\ &\leq Lr^{-m} - \frac{1}{4}a_1\mu^{-\lambda_1/m}r^{-m-\lambda_1} \\ &\leq Lr^{-m} - \frac{a_1}{4\omega(0)}\mu^{-\lambda_1/m}r^{-m-\lambda_1}\omega(r) \\ &\leq Lr^{-m} - b_-r^{-m-\lambda_1}\omega(r), \end{aligned}$$

because  $\omega$  is decreasing. Hence,

$$u_{\text{in}}^-(r, 0) \leq u_0(r) \quad \text{if } \mu^{-1/m} \leq r \leq \mu^{-\kappa/2}. \tag{2.42}$$

Next, if  $\xi < \xi_1$ , then by (2.33), (2.37)–(2.39),

$$\begin{aligned} \frac{[\sigma_t(0)/\sigma^p(0)]\Psi(\xi)}{\psi(\xi)} &\leq C \frac{\sigma_t(0)}{\sigma^p(0)} \\ &= -\frac{mC}{2\lambda_1}\varepsilon^{-2/m}\omega^{(2-\lambda_1)/\lambda_1}(\mu^{-\kappa/2})\omega'(\mu^{-\kappa/2}) \\ &= \frac{mC}{2\lambda_1}\mu^{\kappa-2/m} \left| \frac{\mu^{-\kappa/2}\omega'(\mu^{-\kappa/2})}{\omega(\mu^{-\kappa/2})} \right| \\ &\leq \frac{mC}{2\lambda_1}\mu^{\kappa-2/m} \\ &\leq \frac{\nu}{L - 2\nu}. \end{aligned} \tag{2.43}$$

Since (2.34) implies that  $\psi(\xi) \leq (L - 2\nu)\xi^{-m}$  for all  $\xi < \xi_1$ , we thus obtain

$$\begin{aligned} u_{\text{in}}^-(r, 0) &= \sigma(0)\psi(\xi) \left\{ 1 + \frac{[\sigma_t(0)/\sigma^p(0)]\Psi(\xi)}{\psi(\xi)} \right\} \\ &= \frac{L - \nu}{L - 2\nu} \sigma(0)\psi(\xi) \\ &\leq \frac{L - \nu}{L - 2\nu} \sigma(0)(L - 2\nu)\xi^{-m} \\ &= (L - \nu)r^{-m} \quad \text{for all } r \leq \mu^{-1/m}\xi_1. \end{aligned} \quad (2.44)$$

By definition (2.35) of  $r_0$ , however, in the case where  $r \geq r_0$ , we have

$$\begin{aligned} u_0(r) &\geq Lr^{-m} - b_-r^{-m-\lambda_1}\omega(r) \\ &\geq Lr^{-m} - \nu r^{-m}, \end{aligned}$$

which, combined with (2.44), yields

$$u_{\text{in}}^-(r, 0) \leq u_0(r) \quad \text{if } r_0 \leq r < \mu^{-1/m}\xi_1, \quad (2.45)$$

so that we are left with small  $r$  satisfying  $r < r_0$ . With regard to these, we recall (2.36) and use (2.43) and the trivial estimate  $\psi(\xi) \leq 1$  to obtain

$$\begin{aligned} u_{\text{in}}^-(r, 0) &\leq \sigma(0) \left( 1 + \frac{\nu}{L - 2\nu} \right) \psi(\xi) \\ &\leq \frac{L - \nu}{L - 2\nu} \sigma(0) \\ &= \frac{L - \nu}{L - 2\nu} \mu \\ &\leq \delta \\ &\leq u_0(r) \quad \text{for } r < r_0. \end{aligned}$$

Together with (2.42) and (2.45), this proves (2.32).  $\square$

Combining the above estimates, we can now derive a lower bound of radial solutions.

**Proposition 2.6.** *Assume that  $u_0 = u_0(r)$  is a continuous and positive function of  $r \geq 0$  satisfying*

$$u_0(r) \geq Lr^{-m} - b_-r^{-m-\lambda_1}\omega(r) \quad \text{for all } r > 0$$

*with some  $b_- > 0$ . Then there exists  $c > 0$  such that the solution  $u$  of (2.1) satisfies*

$$u(0, t) \geq c\omega^{-m/\lambda_1}(t^{1/2}) \quad \text{for all } t > 0. \quad (2.46)$$

**Proof.** Let  $b_0 > 0$  be the constant provided by Corollary 2.2, and take any  $\mu > 0$  satisfying  $\mu < \min\{1, \mu_0, \mu_1, \mu_2\}$  with  $\mu_0, \mu_1$  and  $\mu_2$  taken from Lemmas 2.3, 2.4 and 2.5, respectively. Then the function  $u_{\text{in}}^-$  defined by (2.17) satisfies  $u_{\text{in}}^- \geq u$  for  $r = (t + \mu^{-\kappa})^{1/2}$ ,  $t \geq 0$ , by Corollary 2.2 and Lemma 2.4, whereas Lemma 2.5 guarantees that  $u_{\text{in}}^- \geq u$  also

at  $t = 0$ . Since  $u_{\text{in}}^-$  is a subsolution of (2.1) by Lemma 2.3, the comparison principle shows that  $u_{\text{in}}^- \geq 0$  holds for all  $t \geq 0$  and  $r \leq (t + \mu^{-\kappa})^{1/2}$ . In particular,

$$\begin{aligned} u(0, t) &\geq u_{\text{in}}^-(0, t) \\ &= \varepsilon \omega^{-m/\lambda_1} ((t + \mu^{-\kappa})^{1/2}) \\ &\geq \varepsilon \omega^{-m/\lambda_1} (t^{1/2}) \quad \text{for all } t > 0, \end{aligned}$$

because  $\omega$  is decreasing. □

### 3. Upper bound

In this section we give an upper bound for the solution of (2.1) by constructing a suitable super-solution of (2.1). We first consider an appropriate outer region.

**Lemma 3.1.** *Suppose that*

$$u_0(r) \leq Lr^{-m} \quad \text{for all } r > 0 \tag{3.1}$$

and

$$u_0(r) \leq Lr^{-m} - b_+ r^{-l} \omega(r) \quad \text{for all } r \geq 1 \tag{3.2}$$

hold with a positive constant  $b_+$ . Then there exists  $B > 0$  such that the solution  $u$  of (2.1) satisfies

$$u(r, t) \leq Lr^{-m} - \frac{1}{2} b_+ r^{-m-\lambda_1} \omega(r) \quad \text{for all } t \geq 0 \text{ and } r \geq B(t + 1)^{1/2}. \tag{3.3}$$

**Proof.** We let  $C > 0$  satisfy

$$\left| \frac{z\omega'(z)}{\omega(z)} \right| \leq C \quad \text{and} \quad \left| \frac{z^2\omega''(z)}{\omega(z)} \right| \leq C \quad \text{for all } z \geq 0, \tag{3.4}$$

which is possible in view of (1.8) and (1.9). We next fix  $b_2 > 0$  such that

$$b_2 \geq 2b_+ [(m + \lambda_1)|m + \lambda_1 + 2 - N| + |N - 1 - 2m - 2\lambda_1|C + C] \tag{3.5}$$

and, finally, we take  $B > 0$  so large that

$$B \geq \sqrt{2} \sqrt{(m + \lambda_1 + 2)|m + \lambda_1 + 4 - N| + |N - 5 - 2m - 2\lambda_1|C + C} \tag{3.6}$$

and

$$B \geq \sqrt{\frac{2b_2}{b_+}}. \tag{3.7}$$

Then

$$u_{\text{out}}^+(r, t) := \min\{Lr^{-m}, Lr^{-m} - b_+ r^{-m-\lambda_1} \omega(r) + b_2 r^{-m-\lambda_1-2} \omega(r)(t + 1)\}$$

satisfies

$$u_{\text{out}}^+(r, 0) \geq u_0(r) \quad \text{for all } r \geq 0 \tag{3.8}$$

by (3.1) and (3.2). Moreover, at each point  $(r, t)$  where  $u_{\text{out}}^+(r, t) < Lr^{-m}$ , we have  $(u_{\text{out}}^+)^p < (Lr^{-m})^p$  and, thus, repeating the computation in (2.7), we find

$$\begin{aligned} \mathcal{P}u_{\text{out}}^+ &= b_2r^{-m-\lambda_1-2}\omega(r) + (Lr^{-m})^p \\ &\quad + b_1\{(m + \lambda_1)(m + \lambda_1 + 2 - N)r^{-m-\lambda_1-2}\omega(r) \\ &\quad\quad + (N - 1 - 2m - 2\lambda_1)r^{-m-\lambda_1-1}\omega'(r) + r^{-m-\lambda_1}\omega''(r)\} \\ &\quad - b_2\{(m + \lambda_1 + 2)(m + \lambda_1 + 4 - N)r^{-m-\lambda_1-4}\omega(r) \\ &\quad\quad + (N - 5 - 2m - 2\lambda_1)r^{-m-\lambda_1-3}\omega'(r) + r^{-m-\lambda_1-2}\omega''(r)\}(t + 1) - (u_{\text{out}}^+)^p \\ &> b_2r^{-m-\lambda_1-2}\omega(r) \\ &\quad \times \left\{ 1 + \frac{b_1}{b_2} \left[ (m + \lambda_1)(m + \lambda_1 + 2 - N) \right. \right. \\ &\quad\quad \left. \left. + (N - 1 - 2m - 2\lambda_1)\frac{r\omega'(r)}{\omega(r)} + \frac{r^2\omega''(r)}{\omega(r)} \right] \right. \\ &\quad \left. - \left[ (m + \lambda_1 + 2)(m + \lambda_1 + 4 - N) \right. \right. \\ &\quad\quad \left. \left. + (N - 5 - 2m - 2\lambda_1)\frac{r\omega'(r)}{\omega(r)} + \frac{r^2\omega''(r)}{\omega(r)} \right] \frac{t + 1}{r^2} \right\}. \end{aligned}$$

Using (3.4)–(3.6), for all  $(r, t)$  satisfying  $r \geq B(t + 1)^{1/2}$  and  $u_{\text{out}}^+(r, t) < Lr^{-m}$ , we obtain

$$\begin{aligned} \mathcal{P}u_{\text{out}}^+ &> b_2r^{-m-\lambda_1-2}\omega(r) \\ &\quad \times \left\{ 1 - \frac{b_1}{b_2} [(m + \lambda_1)|m + \lambda_1 + 2 - N| + |N - 1 - 2m - 2\lambda_1|C + C] \right. \\ &\quad\quad \left. - [(m + \lambda_1 + 2)|m + \lambda_1 + 4 - N| + |N - 5 - 2m - 2\lambda_1|C + C] \frac{1}{B^2} \right\} \\ &\geq b_2r^{-m-\lambda_1-2}\omega(r) \left\{ 1 - \frac{1}{2} - \frac{1}{2} \right\} \\ &= 0. \end{aligned}$$

Since  $(r, t) \mapsto Lr^{-m}$  is a solution of (2.1), it follows that  $u_{\text{out}}^+$  is a super-solution for all  $r \geq 0$  and  $t \geq 0$ , and therefore, by (3.8), the comparison principle implies  $u \leq u_{\text{out}}^+$  for all  $r \geq 0$  and  $t \geq 0$ . In particular, recalling (3.7), we have

$$\begin{aligned} u(r, t) &\leq u_{\text{out}}^+(r, t) \\ &\leq Lr^{-m} - b_+r^{-m-\lambda_1}\omega(r) + \frac{1}{2}b_+B^2r^{-m-\lambda_1-2}\omega(r)(t + 1) \\ &\leq Lr^{-m} - \frac{1}{2}b_+r^{-m-\lambda_1}\omega(r) \end{aligned}$$

for all  $t \geq 0$  and  $r \geq B(t + 1)^{1/2}$ , which proves (3.3). □

We also need the following elementary property of  $\omega$ , which, along with (1.8), is a simple consequence of its positivity and monotonicity.

**Lemma 3.2.** *For any  $\Lambda > 0$ , there exists  $z_\Lambda > 0$  such that*

$$\omega(\Lambda z) \geq \frac{1}{2}\omega(z) \quad \text{for all } z \geq z_\Lambda.$$



**Proof.** We evidently may assume  $\Lambda > 1$ . We define  $z_\Lambda$  as any sufficiently large number satisfying

$$\left| \frac{z\omega'(z)}{\omega(z)} \right| \leq \frac{1}{2(\Lambda - 1)} \quad \text{for all } z \geq z_\Lambda. \tag{3.9}$$

Then

$$\omega'(z) \geq -\frac{1}{2(\Lambda - 1)} \frac{\omega(z)}{z} \quad \text{for all } z \geq z_\Lambda \tag{3.10}$$

and thus

$$\begin{aligned} \omega(\Lambda z) - \omega(z) &= \int_z^{\Lambda z} \omega'(s) \, ds \\ &\geq -\frac{1}{2(\Lambda - 1)} \int_z^{\Lambda z} \frac{\omega(s)}{s} \, ds \quad \text{for all } z \geq z_\Lambda. \end{aligned}$$

Since  $s \mapsto \omega(s)/s$  decreases on  $(0, \infty)$ , we obtain

$$\begin{aligned} \omega(\Lambda z) - \omega(z) &\geq -\frac{1}{2(\Lambda - 1)} (\Lambda z - z) \frac{\omega(z)}{z} \\ &= -\frac{1}{2} \omega(z) \quad \text{for all } z \geq z_\Lambda, \end{aligned}$$

which proves the lemma. □

We are now in a position to give an upper bound for radial solutions. The proof closely follows that of [2, Lemma 4.3], but we include a complete proof here for convenience.

**Proposition 3.3.** *Suppose that  $u_0$  satisfies (3.1) and (3.2) and that, for each  $\alpha > u_0(0)$ ,  $u_0$  intersects  $\varphi_\alpha$  exactly once in  $(0, \infty)$ . Then there exists  $C > 0$  such that the solution  $u$  of (2.1) satisfies*

$$u(0, t) \leq C\omega^{-m/\lambda_1}(t^{1/2}) \quad \text{for all } t > 0. \tag{3.11}$$

**Proof.** We let  $\sigma(t) := u(0, t)$  and we may assume that  $\sigma$  is unbounded, since otherwise (3.11) is trivial. Thus, there exists  $t_0 > 0$  such that  $\sigma(t_0) > \sigma(0)$ . Then, for each  $t > t_0$ ,  $u(\cdot, t)$  does not intersect  $\varphi_{\sigma(t_0)}$  because the number of intersections of  $u(\cdot, t)$  with the equilibrium  $\varphi_{\sigma(t_0)}$  initially equals 1 and drops at time  $t_0$ . Since  $\sigma$  is unbounded, this means that  $u(\cdot, t) > \varphi_{\sigma(t_0)}$  for all  $t > t_0$ . In particular,  $\sigma(t) > \sigma(0)$  for all  $t > t_0$  and, hence, we may repeat the above argument with  $t_0$  replaced by any  $t_1 \geq t_0$  to obtain  $u(\cdot, t) > \varphi_{\sigma(t_1)}$  for all  $t > t_1$ . Taking  $t \searrow t_1$ , we infer that

$$u(r, t) \geq \varphi_{\sigma(t)}(r) \quad \text{for all } t \geq t_0 \text{ and } r \geq 0.$$

By (2.12) and evident scaling properties of  $\varphi_\alpha$ , there exists  $M > 0$  such that if  $\alpha^{1/m}r \geq M$ , then

$$\begin{aligned} \varphi_\alpha(r) &= \alpha\varphi_1(\alpha^{1/m}r) \\ &\geq \alpha\{L(\alpha^{1/m}r)^{-m} - 2a_1(\alpha^{1/m}r)^{-m-\lambda_1}\} \\ &= Lr^{-m} - 2a_1\alpha^{-\lambda_1/m}r^{-m-\lambda_1}. \end{aligned}$$

Thus, if

$$T := \left( \frac{M}{B\sigma^{1/m}(0)} \right)^2 - 1$$

with  $B$  as provided by Lemma 3.1, for all  $t \geq \max\{T, t_0\}$  and  $r = B(t+1)^{1/2}$  we have

$$\sigma^{1/m}(t)r \geq \sigma^{1/m}(0)B(T+1)^{1/2} = M$$

and therefore

$$u(r, t) \geq \varphi_{\sigma(t)}(r) \geq Lr^{-m} - 2a_1\sigma^{-\lambda_1/m}(t)r^{-m-\lambda_1} \quad \text{at } r = B(t+1)^{1/2} \quad (3.12)$$

for such  $t$ . On the other hand, from Lemma 3.1, we see that

$$u(r, t) \leq Lr^{-m} - \frac{1}{2}b_+r^{-m-\lambda_1}\omega(r) \quad \text{at } r = B(t+1)^{1/2} \text{ for all } t \geq 0. \quad (3.13)$$

Combining (3.12) with (3.13) and solving with respect to  $\sigma(t)$ , we obtain

$$\sigma(t) \leq \left( \frac{4a_1}{b_+} \right)^{m/\lambda_1} \omega^{-m/\lambda_1}(B(t+1)^{1/2}) \quad \text{for all } t \geq \max\{T, t_0\}. \quad (3.14)$$

Now the observation that

$$B(t+1)^{1/2} \leq \sqrt{2}Bt^{1/2}$$

in conjunction with Lemma 3.2, applied to  $\Lambda := \sqrt{2}B$ , yields

$$\omega(B(t+1)^{1/2}) \geq \omega(\sqrt{2}Bt^{1/2}) \geq \frac{1}{2}\omega(t^{1/2}) \quad \text{for all } t \geq z_\Lambda^2,$$

and (3.14) thereby easily leads to (3.11).  $\square$

#### 4. Proof of Theorem 1.1

In this section we complete a proof of Theorem 1.1 by using the upper and lower estimates of radial solutions.

Given an initial value  $u_0(x)$  satisfying (1.2) and (1.6), we define radially symmetric functions by

$$\underline{u}_0(r) := \min\{u_0(x) : |x| \leq r\}, \quad r \geq 0,$$

and

$$\bar{u}_0(r) := \max\{u_0(x) : |x| \geq r\}, \quad r \geq 0.$$

Then

- (i)  $\underline{u}_0(r)$  and  $\bar{u}_0(r)$  are continuous and decreasing in  $r \geq 0$ ,
- (ii)  $0 \leq \underline{u}_0(|x|) \leq u_0(x) \leq \bar{u}_0(|x|) \leq \varphi_\infty(|x|)$  for all  $x \in \mathbb{R}^N \setminus \{0\}$  and
- (iii)  $\underline{u}_0(|x|)$  and  $\bar{u}_0(|x|)$  satisfy (1.6).

Let  $\underline{u}(r, t)$  and  $\bar{u}(r, t)$  denote the solutions of (2.1) with the initial values  $u_0(r)$  and  $\bar{u}_0(r)$ , respectively. Then the solutions exist globally in time and are decreasing in  $r$  for all  $t > 0$ . Moreover, by the comparison principle, the solution of (1.1) satisfies

$$\underline{u}(|\cdot|, t)\|_{L^\infty} \leq \|u(\cdot, t)\|_{L^\infty} \leq \|\bar{u}(|\cdot|, t)\|_{L^\infty}, \quad x \in \mathbb{R}^N,$$

for all  $t > 0$ . Since  $\underline{u}(r, t)$  and  $\bar{u}(r, t)$  are decreasing in  $r$  for each  $t > 0$ , since

$$\underline{u}(0, t) \geq c\omega^{-m/\lambda_1}(t^{1/2}) \quad \text{for all } t > 0$$

by Proposition 2.6, and since

$$\bar{u}(0, t) \leq C\omega^{-m/\lambda_1}(t^{1/2}) \quad \text{for all } t > 0$$

by Proposition 3.3, we obtain the desired estimates of the grow-up rate of the solution of (1.1).

**Acknowledgements.** M.F. was supported by the Slovak Research and Development Agency under contract APVV-0414-07. E.Y. was partly supported by the Grant-in-Aid for Scientific Research (A) (19204014) from the Ministry of Education, Science, Sports and Culture.

## References

1. M. FILA, J. R. KING, M. WINKLER AND E. YANAGIDA, Optimal lower bound of the grow-up rate for a supercritical parabolic equation, *J. Diff. Eqns* **228** (2006), 339–356.
2. M. FILA, J. R. KING, M. WINKLER AND E. YANAGIDA, Grow-up of solutions of a semilinear parabolic equation with a critical exponent, *Adv. Diff. Eqns* **12** (2007), 1–26.
3. M. FILA, J. R. KING, M. WINKLER AND E. YANAGIDA, Linear behaviour of solutions of a super-linear heat equation, *J. Math. Analysis Applic.* **340** (2008), 401–409.
4. M. FILA AND M. WINKLER, Rate of convergence to a singular steady state of a supercritical parabolic equation, *J. Evol. Eqns* **8** (2008), 673–692.
5. M. FILA, M. WINKLER AND E. YANAGIDA, Grow-up rate of solutions for a supercritical semilinear diffusion equation, *J. Diff. Eqns* **205** (2004), 365–389.
6. M. FILA, M. WINKLER AND E. YANAGIDA, Convergence rate for a parabolic equation with supercritical nonlinearity, *J. Dynam. Diff. Eqns* **17** (2005), 249–269.
7. M. FILA, M. WINKLER AND E. YANAGIDA, Slow convergence to zero for a parabolic equation with supercritical nonlinearity, *Math. Annalen* **340** (2008), 477–496.
8. M. FILA, M. WINKLER AND E. YANAGIDA, Convergence to self-similar solutions for a semilinear parabolic equation, *Discrete Contin. Dynam. Syst.* **21** (2008), 703–716.
9. V. GALAKTIONOV AND J. L. VÁZQUEZ, Continuation of blow-up solutions of nonlinear heat equations in several space dimensions, *Commun. Pure Appl. Math.* **50** (1997), 1–67.
10. C. GUI, W.-M. NI AND X. WANG, On the stability and instability of positive steady states of a semilinear heat equation in  $\mathbb{R}^n$ , *Commun. Pure Appl. Math.* **45** (1992), 1153–1181.
11. C. GUI, W.-M. NI AND X. WANG, Further study on a nonlinear heat equation, *J. Diff. Eqns* **169** (2001), 588–613.
12. M. HOSHINO AND E. YANAGIDA, Sharp estimate of the convergence rate for a semilinear parabolic equation with supercritical nonlinearity, *Nonlin. Analysis TMA* **69** (2008), 3136–3152.

13. D. D. JOSEPH AND T. S. LUNDGREN, Quasilinear Dirichlet problems driven by positive sources, *Arch. Ration. Mech. Analysis* **49** (1973), 241–269.
14. N. MIZOGUCHI, Growup of solutions for a semilinear heat equation with supercritical nonlinearity, *J. Diff. Eqns* **227** (2006), 652–669.
15. P. POLÁČIK AND E. YANAGIDA, On bounded and unbounded global solutions of a supercritical semilinear heat equation, *Math. Annalen* **327** (2003), 745–771.
16. P. POLÁČIK AND E. YANAGIDA, Nonstabilizing solutions and grow-up set for a supercritical semilinear diffusion equation, *Diff. Integ. Eqns* **17** (2004), 535–548.
17. P. QUITTNER AND PH. SOUPLLET, *Superlinear parabolic problems: blow-up, global existence and steady states*, Birkhäuser Advanced Texts (Birkhäuser, Basel, 2007).
18. C. STINNER, Very slow convergence to zero for a supercritical semilinear parabolic equation, *Adv. Diff. Eqns* **14** (2009), 1085–1106.
19. C. STINNER, Very slow convergence rates in a semilinear parabolic equation, *NoDEA. Nonlin. Diff. Eqns Applic.* **17** (2010), 213–227.
20. X. WANG, On the Cauchy problem for reaction-diffusion equations, *Trans. Am. Math. Soc.* **337** (1993), 549–590.