

Results: There were 821 (56%) male and 642 (44%) female mood disorder patients. BPD-NOS diagnosis among all mood disorder patients were 194 (13.2%). Re-hospitalization rate of patients treated with the diagnosis of BPD-NOS was 6.7%. Duration of illness, episode duration, symptom variety and treatment responses were the most common features “making the diagnosis atypical”.

Conclusions: A more systematic and detailed evaluation is needed for appropriate acute or preventive treatment. Studies are needed on patients with BD-NOS with comorbid schizophrenia or delusional disorder as specified in DSM-IV-TR.

P124

Behavioral disorders at adolescent with primary cerebral dysfunction

O. Pilyavskaya, N. Butorina, D. Verbovetskaya. *Department of Child and Adolescent Psychiatry, The Ural State Medical Academy of Additional Education, Chelyabinsk Chelyabinsk Region, Russia*

Introduction: The object was to explore behavioral disorders at adolescent of female sex with a primary cerebral dysfunction.

Methods: 49 adolescent of female gender in age of 11-16 with a primary cerebral dysfunction, behavioral disorders and with intelligence quotient more than 80 on the Wechsler's scale were observed. All adolescent were explored with clinical, neuropsychological and neurological methods.

Results: At all adolescent were early psychical disorders (in 100% of cases), brain trauma in age of 7 (22%) and negative sociological environment (79%). They had symptoms of psychoorganical syndrome with behavioral and emotional disorders. These disorders were presented by such psychopathological variants as apathic (15%), labile (38%), hyperactive (41%) and aggressive (16%). The spectrum of the behavioral disorders was studied in the relation to the age. The standard development of behavioral disorders was noticed in the age of 7-10. The worsening of cerebral dysfunction, psychopathological traits of personality disorders were in the age of 12. And the formation of the psychopathy with behavioral disorders was in the age of 16.

Conclusions: The behavioral disorders at adolescent of female sex with primary cerebral dysfunction were bound with gender and options of psychoorganical syndrome.

P125

Suicidal risk in bipolar disorder patients

M. Pompili^{1,2}, A. Grispiini², P. Girardi², R. Tatarelli².¹ *McLean Hospital, Harvard Medical School, Belmont, MA, USA*
² *Department of Psychiatry, Sant'Andrea Hospital, University of Rome, Rome, Italy*

Background: Suicidal risk is very high among bipolar disorder (BPD) patients. Risks of suicide attempts are not as well quantified but remain the most important factor for predicting risk of completed suicide.

Methods: We retrospectively evaluated 88 patients diagnosed as bipolar I, II, or unipolar depression. Of these, 44 had made at least one suicide attempt, and were matched for age, sex and diagnosis with 44 patients who had never attempted suicide.

Results: In the univariate contrasts, suicidal patients were more likely to be: men, single, younger currently but not at onset, bipolar, substance abusing, and being unemployed. In a logistic regression only older age in the control group and occupation held the significance.

Limitations: Study findings may not generalize to other samples, settings, and treatments.

Conclusions: Our results support previous finding of literature but overall add a consistent emphasis on the role two variables involved in the precipitation of suicide attempts in bipolar and unipolar patients, that is the role of age and that of occupation. The former may be identified as Trait-dependent risk factors (unchangeable) the latter as State-dependent risk factors (which can potentially be modified).

P126

Social networks of patients with bipolar affective disorder.

M. Poradowska-Trzos, D. Dudek, W. Datka, A. Zieba. *Department of Psychiatry, University Hospital, Krakow, Poland*

Background and aims: According to Axer's definition a social network is a certain number of people with whom a person remains contact. These people provide stimuli - both positive and negative. In everyday life and even more during the periods of disease all people need to experience positive aspects of their network - especially support.

The purpose of the research is to analyze the structure of social networks as well as types and sources of support received by patients with bipolar affective disease (BID).

Methods: The study group consisted of euthymic outpatients (40 patients). The control group consisted of healthy volunteers of sex and age corresponding to the case group. Assessment of a structure of a social network and the amount of support received was made according to Bizon's questionnaire and Cohen's ISEL.

Results: It has been shown that social networks of BID patients differ from the ones of healthy people. Patients have social networks characterized by the structure and activity comparable with that of healthy people. But often there is only one person who carries the whole burden of support. When compared to the control group these patients receive less support, and the emotional support is the most deficient.

Conclusions: Information about a patient's social network may be helpful when planning treatment in hospital as well as preparing psychosocial interventions in an outpatients' clinic. The information enables to recognize deficiencies so as to make attempts to reorganize or reconstruct a network.

P127

A perception of social support in the aspect of a cognitive style of patients with affective disorders.

M. Poradowska-Trzos, D. Dudek, W. Datka, A. Zieba. *Department of Psychiatry, University Hospital, Krakow, Poland*

Background: According to Aaron Beck dysfunctional thinking patterns appear also in euthymic patients, after withdrawal of acute disease's symptoms. Patients have disordered, negative image of themselves, of their future and the surrounding world.

It has been shown that a way a man perceives possessed social support has for him a basic meaning.

Aims: The purpose of the research is to analyze the relationship between perceived social support and patient's cognitive style.

Methods: The study group consisted of euthymic outpatients diagnosed with recurrent depressive disorder (40 patients) or bipolar affective disorder (40 patients). The control group consisted of healthy volunteers of sex and age corresponding to the case group.