

physicians alone (0.88%, CI 0.09) ($p < 0.0001$). Return rates were statistically higher for PGY2 (1.67% CI 0.35) and PGY3 (1.66% CI 0.28) residents compared to staff physicians alone ($p < 0.0001$). There was no difference in return visit rates between staff physicians and third year medical students (1.07% CI 0.27), fourth year medical students (1.21% CI 0.37), PGY1 (1.42% CI 0.22), PGY4 (1.23% CI 0.54) or PGY5 (1.33% CI 0.49) residents. **Conclusion:** This study demonstrated that the involvement of learners in ED patient assessments increased the rate of short-term unscheduled return visits. Moreover, return visit rates were highest for PGY2 and PGY3 residents. Further work is needed to understand the factors that contribute to this phenomenon.

Keywords: bounce backs, short-term unscheduled return visits

P042

Pilot study for the inter-arm blood pressure systematic measurement during the diagnosis of transient ischemic attack in the emergency department

P. La Rochelle, MD, S. Lavoie, BN, V. Boucher, BA, M. Émond, MD, MSc, J. Perry, MD, MSc, Université Laval, Québec City, QC

Introduction: Our principal aim was to document the feasibility of the systematic measurement of the inter-arm blood pressure difference (IABPD) during an episode of transient ischemic attack (TIA) or mild stroke diagnosed in the Emergency Department (ED). As secondary goal was to compare the systolic blood pressure (BP) at triage with the systolic BPs measured during the IABPD. **Methods:** This is a single center pilot study. Patients presenting in the ED for a diagnosis of TIA were recruited. Once patient has been triaged and diagnosed of TIA, a research assistant made sure that the patient lay on a stretcher for at least 5 minutes. Two automated sphygmomanometers were applied, on each arm. No specific device or device calibration were required. Three consecutive simultaneous BP readings were performed, inverting cuffs arm to arm between each reading. Only the last two set of readings were used to calculate the mean IABPD. This method enables to minimize the error coming from the potential sphygmomanometers' inaccuracies. **Results:** 32 patients were recruited from June to September 2017 and all had a successful IABPD measurement. Four patients had an IABPD >10 mmHg, varying from 1.5 to 13 mmHg when the left arm was higher and from 1 to 61 mmHg when the right arm was higher. Of the 22 patients where the triage BP arm side selection was recorded, only 11 were congruent with the arm presenting the highest BP during the IABPD measurement. Selecting of the arm with the highest BP value may better reflect cerebrovascular risk exposition. The mean systolic BP at triage was 159.3 mmHg (95%CI: 144.9-173.7) compared to 144.8 mmHg (95%CI: 132.9-156.7) if the arm with the highest value during the IABPD measurement is selected and 142.4 mmHg (95%CI: 130.8-154.0) if the same arm as triage is selected. The p-value for these differences were 0.003 and 0.001 respectively. The patient which presented the IABPD of 61 mmHg, had a stroke 3 days after its ED visit which subsequently led to her death 10 days later. **Conclusion:** Our results show that the systematic IABPD measurement using a pragmatic approach in the ED is feasible and is ready to investigate its use in the context of a new TIA or mild stroke. This information may contribute to a better discrimination of the short-term risk of stroke and may help to diagnose acute aortic dissection, monitor more accurately BP during hyperacute stroke or estimate intracerebral hemorrhage risk if systemic thrombolysis is considered.

Keywords: Interarm blood pressure, pragmatic method, transient ischemic attack

P043

Trauma team leaders in Canada: A national survey

V. Belhumeur, C. Malo, MD, MSc, A. Nadeau, MSc, S. Hegg, PhD, A. Gagné, BA, M. Émond, MD, MSc, Laval, Québec, QC

Introduction: It was demonstrated that the early trauma team activation (TTA) could improve younger trauma patients outcomes and mortality rates. However, the link between older patient prognosis improvement and the activation / effectiveness of the Trauma team (TT) is still unclear. There is also a lack of information about the exact and optimal structure of TTs and their activation criteria, which may differ across centers. The main objective of this study is to provide a description of the current TT available in level 1 and 2 centres across Canada. **Methods:** In 2017, a survey using a modified Dillman technique was sent to 210 health professionals scattered across all Canadian trauma care facilities. The survey included questions regarding 1) the presence and the composition of a TT, 2) the established TT activation criteria, and finally 3) the initial patient care. **Results:** A total of 107 (57%) completed surveys were received. Among them, only 22 (11.7%) were from level 1 or 2 centres and were therefore considered for analyses. Seventeen respondents had a TT in their centre, and they all shared their TT activation criteria (1 to 27 different indications). Most frequently mentioned criteria were: suspected injuries (58.8%), judgment of the emergency physician (41.2%), systolic blood pressure (47.1%), Glasgow Coma score (35.3%) and respiratory rate (28%). In presence of a prehospital care warning trauma, the initial assessment of a severely injured patient is exclusively completed by a member of the TT for only 35.1% of the respondents. For 11.8% of respondents, TT coordinates airway management. For 64.7% of participants, the TT leader is the dedicated care provider to accompany patients until final orientation. **Conclusion:** These results suggest a great variability across Canada regarding the roles assumed by the TT, but also regarding the activation criteria leading them to take action.

Keywords: emergency care, polytrauma, trauma team

P044

Use of a gait tracking device to count steps of older emergency department patient

J. Estrada-Codecido, MD, J. Lee, MD, MSc, University of Toronto, Toronto, ON

Introduction: Delirium is a common complication among older people who need care in the emergency department (ED). Mobility is an evidence-based non-pharmacologic strategy shown to reduce delirium and functional decline among older patients in the acute care setting. However, previous research has shown that compliance with mobility is important to achieve this decreased incidence of delirium. Gait tracking devices have been used in previous studies to accurately measure steps, engagement and intensity of physical activity in older hospitalized patients. The objectives of this study are to compare the feasibility and validate the accuracy of three accelerometer-based gait tracking devices. This is the first step in a program of research to objectively measure mobility among older ED patients as a potential marker of delirium risk. **Methods:** This is a prospective, observational study of patients 65 years of age and older during their ED visit. We excluded those with critical illness, unable to communicate or