

- 12 McWhinney IR, Epstein RM, Freeman TR. Rethinking somatization. *Ann Intern Med* 1997; **126**: 747–50.
- 13 Price J, Farmer G, Harris J, Hope T, Kennedy S, Mayou R. Attitudes of women with chronic pelvic pain to the gynaecological consultation: a qualitative study. *BJOG* 2006; **113**: 446–52.
- 14 Heijmans M, Hartman TCO, van Weel-Baumgarten E, Dowrick C, Lucassen PLBJ, van Weel C. Experts' opinions on the management of medically unexplained symptoms in primary care. A qualitative analysis of narrative reviews and scientific editorials. *Fam Pract* 2011; **28**: 444–55.
- 15 McCabe R, Roder-Wanner UU, Hoffmann K, Priebe S. Therapeutic relationships and quality of life: association of two subjective constructs in schizophrenia patients. *Int J Soc Psychiatry* 1999; **45**: 276–83.
- 16 Highet NJ, McNair BG, Thompson M, Davenport TA, Hickie IB. Experience with treatment services for people with bipolar disorder. *Med J Aust* 2004; **181**: S47–51.
- 17 Page LA, Wessely S. Medically unexplained symptoms: exacerbating factors in the doctor-patient encounter. *J R Soc Med* 2003; **96**: 223–7.
- 18 Ware Jr JE, Davies AR. Behavioral consequences of consumer dissatisfaction with medical care. *Eval Program Plan* 1983; **6**: 291–7.
- 19 Rusch N, Holzer A, Hermann C, Schramm E, Jacob GA, Bohus M, et al. Self-stigma in women with borderline personality disorder and women with social phobia. *J Nerv Ment Dis* 2006; **194**: 766–73.
- 20 Rosendal M, Fink P. Management of medically unexplained symptoms – includes diagnosis, specific treatments, and appropriate communication. *BMJ* 2005; **330**: 4–5.
- 21 Aiarzaguena JM, Grandes G, Gaminde I, Salazar A, Sanchez A, Arino J. A randomized controlled clinical trial of a psychosocial and communication intervention carried out by GPs for patients with medically unexplained symptoms. *Psychol Med* 2007; **37**: 283–94.
- 22 Lequesne ER, Hersh RG. Disclosure of a diagnosis of borderline personality disorder. *J Psychiatr Pract* 2004; **10**: 170–6.
- 23 Sanislow CA, Marcus KL, Reagan EM. Long-term outcomes in borderline psychopathology: old assumptions, current findings, and new directions. *Curr Psychiatry Rep* 2012; **14**: 54–61.
- 24 Woollaston K, Hixenbaugh P. 'Destructive Whirlwind': nurses' perceptions of patients diagnosed with borderline personality disorder. *J Psychiatr Ment Health Nurs* 2008; **15**: 703–9.
- 25 Kealy D, Ogrodniczuk JS. Marginalization of borderline personality disorder. *J Psychiatr Pract* 2010; **16**: 145–54.
- 26 Hersh R. Confronting myths and stereotypes about borderline personality disorder. *Soc Work Ment Health* 2008; **6**: 13–32.



## psychiatry in literature

### Not I: Beckett and psychiatry

Elizabeth Barry, Matthew Broome, Jonathan Heron

On 23 September 1935, Samuel Beckett wrote to his friend Thomas MacGreevy: 'I went down to Bedlam this day week, and went round the wards for the first time, with scarcely any sense of horror, though I saw everything, from mild depression to profound dementia' (*The Letters of Samuel Beckett*, Vol I, p. xx). Beckett was, in this visit, following a long tradition of writers visiting the Bethlem Royal Hospital in search of creative inspiration. A fictitious London asylum (the Magdalen Mercyseat) and its patients (including, in Beckett's words, those 'melancholic . . . hebephrenic . . . hypomanic . . . schizoid') duly appear in Beckett's comic novel, *Murphy* (1938). Beckett's interest in mental disorder was, however, deeper than such early characterisations might suggest. His concerns about his own anxiety, depression and psychosomatic illness prompted him to undertake psychoanalysis with Wilfred Bion at the Tavistock Clinic, and his work offers increasingly sensitive representations of mental distress and speech disorder. His play *Not I* (1972), for instance, features a disembodied (but female) mouth speaking in fractured phrases about a past traumatic event, possibly a rape. Across the stage, a hooded Auditor listens silently, responding only to lift his arms in a gesture of 'helpless compassion'. The character ('Mouth') describes the shift from a state of near mutism ('not as much as goodbye') to one characterised by a pathological pressure of speech, creating acute social anxiety. This tachylalia is enacted by Mouth, although she refuses to identify herself with the subject of her tale ('she found herself in the . . . what? . . . who? . . . no! . . . she!').

These very words are scrawled on the personal script of Billie Whitelaw, the performer directed in the role of 'Mouth' by Anthony Page (and Samuel Beckett) in 1973. Whitelaw's experiences of performing the play were combined with new approaches to staging Beckett in an event, *Beckett and Brain Science*, held at the University of Warwick in collaboration with London (Birkbeck) and Reading in 2012, which brought together humanities scholars, arts practitioners, psychiatrists and trainee doctors. The theatre company Fail Better Productions worked with clinicians to investigate practically how Beckett's characters might inform clinical care.

The trainee doctors found the performance work particularly interesting in relation to the interplay between Mouth and the silent Auditor. We conducted some exploratory activities between these two characters and their analogous relationship to the roles of patient and clinician. Her 'whole body like gone . . . just the mouth . . . like maddened . . . and can't stop . . . no stopping it' became an embodiment of anxiety and pressured speech which the workshop participants could safely explore through both the Auditor's 'helpless compassion' and the experience of speaking the rapid monologue. This workshop, which has now been adapted for NHS Kent, Surrey and Sussex Deanery, and delivered to a group including consultant psychiatrists, specialist nurses and geriatricians, offers medical educators a way to use Beckett in their classroom, and to explore problems relevant to both compassion and care in practice. The play becomes a phenomenological case history and the methods of the workshop offer strategies for interpersonal treatment.

The British Journal of Psychiatry (2014)  
204, 239. doi: 10.1192/bjp.bp.113.130120