

**ON A PROPOSITIONAL CALCULUS
WHOSE DECISION PROBLEM IS RECURSIVELY
UNSOLVABLE¹⁾**

AKIRA NAKAMURA

Dedicated to Professor Katuzi Ono on his 60th birthday

§0. Introduction

The purpose of this paper is to present a propositional calculus whose decision problem is recursively unsolvable. The paper is based on the following ideas:

- (1) Using Löwenheim-Skolem's Theorem and Surányi's Reduction Theorem, we will construct an infinitely many-valued propositional calculus corresponding to the first-order predicate calculus.
- (2) It is well known that the decision problem of the first-order predicate calculus is recursively unsolvable.
- (3) Thus it will be shown that the decision problem of the infinitely many-valued propositional calculus is recursively unsolvable.

In this paper, we consider semantically the problem. That is, we define a validity of wff in our logical system and we will discuss on the problem to decide whether or not an arbitrary wff in our system is valid.²⁾

§1. Logical system L

We consider a logical system L :

- (1) Propositional variables: $F_1, F_2, \dots, G_1, G_2, \dots, P_1, P_2, \dots$.
- (2) Truth-values: Let N be the set of natural numbers and $\Omega = \{0, 1\}$.

We define functions f, g as follows:

Received March 1, 1969

¹⁾ This research was done while the author stayed at Dept. of Information Science, Univ. of North Carolina at Chapel Hill.

²⁾ In the first-order predicate calculus, the semantical decision problem is equivalent to the syntactical one by the completeness theorem.

$$f: N \rightarrow \Omega, \quad f \in 2^N,$$

$$g: N \times N \times N \rightarrow 2^N, \quad g \in (2^N)^{N \times N \times N}.$$

A *truth-value* is defined as a member of $(2^N)^{N \times N \times N}$, i.e. it is such a function g . Let us say here (x, y, z) in $N \times N \times N$ as a *coordinate*, and x, y, z as x -coordinate, y -coordinate, z -coordinate, respectively.

(3) Logical operations¹⁾:

Monadic operations: $X, Y, Z, \exists_x, \exists_y, \exists_z, \diamond, \neg$,

Duadic operation: \vee .

(4) Truth-value functions:

Let us denote as follows:

$$f(\lambda) = *_{\lambda}, \quad \lambda \in N, *_{\lambda} \in \Omega,$$

$$g(x, y, z) = v_{xyz}, \quad x, y, z \in N, v_{xyz} \in 2^N.$$

X : If a truth-value of wff \mathfrak{A} has v_{iii} at (i, i, i) in $N \times N \times N$, the truth-value of $X\mathfrak{A}$ has the same v_{iii} at every (i, y, z) where $y, z = 1, 2, 3, \dots$.

Y, Z : Those are defined by the similar way to X .

\vee, \neg : Those are defined by the usual way.

\exists_x : We consider all elements (x, j, k) in $N \times N \times N$ where j, k are constants. If there exists a such that a $*_{\lambda}$ at (a, j, k) of truth-value of wff \mathfrak{A} is 1, then the truth-value of $\exists_x \mathfrak{A}$ has 1 at $*_{\lambda}$ of every (x, j, k) .

If a truth-value of \mathfrak{A} has 0 at $*_{\lambda}$ of (x, j, k) for every x , then the truth-value of $\exists_x \mathfrak{A}$ has 0 at $*_{\lambda}$ in every (x, j, k) .

\exists_y, \exists_z : Those are defined by the similar way to \exists_x .

\diamond : For every $g(x, y, z)$,

$$\diamond g(x, y, z) = \begin{cases} \text{every } *_{\lambda}(\lambda = 1, 2, 3, \dots) \text{ is 1, if } *_{\lambda} = 1 \text{ for some } \lambda. \\ \text{every } *_{\lambda}(\lambda = 1, 2, 3, \dots) \text{ is 0 otherwise.} \end{cases}$$

The logical system L is considered as a kind of infinitely many-valued propositional logic. In this paper, a truth-value whose $*_{\lambda}$ ($\lambda = 1, 2, 3, \dots$) at every (x, y, z) are all 1 is called the *designated value*. And further a wff \mathfrak{A} is called *valid* if and only if the \mathfrak{A} takes always the designated value independently of truth-values of propositional variables P_1, P_2, \dots, P_n in \mathfrak{A} .

¹⁾ Using those logical operations, we define $P_1 \wedge P_2 \stackrel{D}{=} \neg(\neg P_1 \vee \neg P_2)$, $P_1 \supset P_2 \stackrel{D}{=} \neg P_1 \vee P_2$.

§2. Relations between the first-order predicate calculus K and the system L

We shall give some relations between the first-order predicate calculus K and the logical system L .

According to the Surányi Reduction Theorem, we have the following one:

THEOREM. *For every wff \mathfrak{A} in K , we can construct a wff \mathfrak{B} of the following form:*

$$(I) \quad (\exists x)(\exists y)(\exists z) M_1 \vee (\exists x)(\exists y)(z)M_2$$

where M_1 and M_2 are quantifier-free and contain non but monadic and duadic predicates. And, in this case, \mathfrak{A} is equivalent to \mathfrak{B} in regard to the universal validity.

From now on, we shall denote Surányi Reduction Form (I) of an arbitrary wff \mathfrak{A} in K as \mathfrak{A}^* .

Now, for wff \mathfrak{A}^* and each subformula \mathfrak{S} of \mathfrak{A}^* in K , let $h(\mathfrak{S})$ be a wff in L obtained by using inductively the following (i)-(iii).

(i) If \mathfrak{S} is a monadic predicate $F(x)$, then

$$h(F(x)) \rightarrow \diamond XF.^{1)}$$

where \rightarrow means “correspondence”.

(ii) If \mathfrak{S} is a duadic predicate $G(x, y)$, then

$$h(G(x, y)) \rightarrow \diamond(XG^1 \wedge YG^2)^{1)}.$$

Here, it needs not to consider such a case as $h(H(x, y, z))$ because the form (I) contains only monadic and duadic predicates as shown above.

(iii) If \mathfrak{S} contains logical operations or quantifier, then

$$\begin{aligned} h(\neg \mathfrak{S}_1) &\rightarrow \neg h(\mathfrak{S}_1), \\ h(\mathfrak{S}_1 \vee \mathfrak{S}_2) &\rightarrow h(\mathfrak{S}_1) \vee h(\mathfrak{S}_2), \\ h((\exists x)\mathfrak{S}_1) &\rightarrow \exists_x h(\mathfrak{S}_1), \\ h((\exists y)\mathfrak{S}_1) &\rightarrow \exists_y h(\mathfrak{S}_1), \\ h((\exists z)\mathfrak{S}_1) &\rightarrow \exists_z h(\mathfrak{S}_1). \end{aligned}$$

For example:

$$h((\exists x)(\exists y)(F(x) \& G(x, y))) \rightarrow \exists_x \exists_y (\diamond XF \wedge \diamond(XG^1 \wedge YG^2)).$$

¹⁾ Of course, $h(F(y)) \rightarrow \diamond YF$, $h(F(z)) \rightarrow \diamond ZF$, $h(G(y, z)) \rightarrow \diamond(YG^1 \wedge ZG^2)$, ...

We shall write $h(\mathfrak{A}^*)$ as $\tilde{\mathfrak{A}}^*$.

Then, we shall prove, in §3, the following theorems:

THEOREM 1. *If $\tilde{\mathfrak{A}}^*$ is valid in L , then \mathfrak{A}^* is universally valid in K .*

THEOREM 2. *If \mathfrak{A}^* is universally valid in K , then $\tilde{\mathfrak{A}}^*$ is valid in L .*

Now, assume that the decision problem of validity in L is recursively solvable. Then, we have an effective procedure to decide whether or not an arbitrary wff $\tilde{\mathfrak{A}}^*$ in L is valid. Thus, from Theorem 1 and 2 we have also an effective procedure to decide whether or not \mathfrak{A}^* in K is universally valid. But, \mathfrak{A}^* is Surányi's reduction form of \mathfrak{A} . Therefore it follows that the decision problem of predicate calculus is recursively solvable. This is contradict with (2) in §0. Thus, we know that the decision problem in L is recursively unsolvable.

§3. Proofs of Theorem 1 and 2

Now, we shall give proofs of Theorem 1 and 2.

Theorem 1:

We prove the following Theorem 1' which is equivalent to Theorem 1.

THEOREM 1'. *If \mathfrak{A}^* is not universally valid in K , then $\tilde{\mathfrak{A}}^*$ is not valid in L .*

Proof. To prove this theorem we use Löwenheim-Skolem's Theorem which is expressed as follows: a wff \mathfrak{F} in K is universally valid if \mathfrak{F} is valid in an enumerable infinite domain ω .

Using this theorem and our assumption of Theorem 1', we are able to let a truth-value of \mathfrak{A}^* be F (falsity) in ω by some suitable truth-value assignment. Here let us denote elements in ω as e_1, e_2, e_3, \dots , and assume that the following predicates occur in \mathfrak{A}^* .

$$(II) \quad F_1(x), F_2(x), \dots, F_\alpha(x); \dots; F_1(z), F_2(z), \dots, F_\alpha(z), \\ G_1(x, x), \dots, G_\beta(x, x); G_1(x, y), \dots; G_1(z, z), \dots, G_\beta(z, z).$$

Here, it is possible to assume that those predicates actually occur in \mathfrak{A}^* . For if $F_1(x)$ occurs neither in $(\exists x)(\exists y)(\exists z)M_1$ nor in $(\exists x)(\exists y)(z)M_2$, then it is sufficient to consider a formula $(\exists x)(\exists y)(\exists z)(M_1 \& F_1(x) \vee \neg F_1(x))$ which is equivalent to $(\exists x)(\exists y)(\exists z)M_1$.

Now, according to our assumption a truth-value of \mathfrak{A}^* is F in ω by a truth-value assignment for predicates (II). Say that the truth-value assignment is as follows:

such that $\diamond(G_i^1 \wedge G_i^2)$ obtained from $G_i(e_k, e_l)$ whose value is \mathbf{F} in (III) does not take the value $(1, 1, 1, 1, \dots)$.

By repeated applications of this process, we give values to $G_1^1, G_2^1, \dots, G_\beta^1; G_1^2, G_2^2, \dots, G_\beta^2$ at $(1, 1, 1), (2, 2, 2), (3, 3, 3), \dots$ in $N \times N \times N$ and in this case values at (ν_1, ν_2, ν_3) where at least two of ν_1, ν_2 and ν_3 are different are arbitrary.

This process is always possible too. Because since our $(*_1, *_2, *_3, \dots)$ is an infinite sequence of 0,1, it is possible by the definition of truth-value function of \diamond .

That is, for example: let us assume that

$$\begin{array}{rcccl}
 & \textcircled{1} & G_i(e_1, e_1): \mathbf{T}, & & \textcircled{2} & G_i(e_2, e_1): \mathbf{F}, & & \textcircled{4} & G_i(e_3, e_1): \mathbf{F}, & \dots \\
 1) & \textcircled{3} & G_i(e_1, e_2): \mathbf{T}, & \swarrow & \textcircled{5} & G_i(e_2, e_2): \mathbf{T}, & \swarrow & \textcircled{8} & G_i(e_3, e_2): \mathbf{T}, & \dots \\
 & \textcircled{6} & G_i(e_1, e_3): \mathbf{F}, & \swarrow & \textcircled{9} & G_i(e_2, e_3): \mathbf{T}, & \swarrow & & G_i(e_3, e_3): \mathbf{F}, & \dots
 \end{array}$$

Then, first we enumerate those predicates as the above-mentioned $\textcircled{1}, \textcircled{2}, \textcircled{3}, \dots$ and we give an assignment as follows:

$$\begin{array}{rcc}
 & G_i^1(e_1): (1, 0, 1, *_4, *_5, \dots) & G_i^2(e_1): (1, 0, 0, \dots) \\
 2) & G_i^1(e_2): (0, 0, 0, 0, 1, \dots) & G_i^2(e_2): (0, 0, 1, 0, 1, 0, 0, 1, \dots) \\
 & G_i^1(e_3): (0, 0, 0, 0, 0, 0, 0, 1, \dots) & G_i^2(e_3): (0, 0, 0, 0, 0, 0, 0, 0, 1, \dots) \\
 & \vdots & \vdots
 \end{array}$$

where $G_i^j(e_1)$ ($j = 1, 2$) means a value of G_i^j at $(1, 1, 1)$ in $N \times N \times N$ and $G_i^j(e_2)$ means a value of G_i^j at $(2, 2, 2)$ in $N \times N \times N$, etc..

2) is constructed such that the first 1, the third 1 from the left in $(1, 0, 1, *_4, *_5, \dots)$ of $G_i^1(e_1)$ correspond to \mathbf{T} of the enumeration $\textcircled{1}, \textcircled{3}$ in 1).

Now, notice that in an enumerable infinite domain N the operation $(\exists x)((\exists y), (\exists z))$ can be interpreted as an infinite disjunction on x -coordinate (y -coordinate, z -coordinate). For example: $(\exists x)\mathfrak{A}(x, y, z)$ is interpreted as

$$\mathfrak{A}(1, y, z) \vee \mathfrak{A}(2, y, z) \vee \dots$$

And also we notice that

- (1) if $F(e_1, e_2)$ is a truth-value of $F(x, y)$, it is considered by the definition of X, Y, Z as a value at $(1, 2, z)$ in $N \times N \times N$ where $z = 1, 2, 3, \dots$.
- (2) if $F(e_1, e_2)$ is a truth-value of $F(y, x)$, it is considered as a value at $(2, 1, z)$ in $N \times N \times N$ where $z = 1, 2, 3, \dots$.

- (3) if $F(e_2, e_2)$ is a truth-value of $F(x, x)$, it is considered as a value at $(2, y, z)$ in $N \times N \times N$ where $y, z = 1, 2, 3, \dots$,
- (4) and so on.

Thus, from the above-mentioned truth-value assignment, the construction of $\tilde{\mathfrak{U}}^*$ and the interpretation of existential quantifier in the domain N , we are able to let $\tilde{\mathfrak{U}}^*$ be not valid in L . Therefore, we get Theorem 1'.

Throem 2:

We shall prove the following Theorem 2' which is equivalent to Theorem 2.

THEOREM 2'. *If $\tilde{\mathfrak{U}}^*$ is not valid in L , then \mathfrak{A}^* is not universally valid in K .*

Proof. Let us notice that $\tilde{\mathfrak{U}}^*$ is of a form $\exists_x \exists_y \exists_z \tilde{M}_1^* \vee \exists_x \exists_y \neg \exists_z \neg \tilde{M}_2^*$ where \tilde{M}_1^* and \tilde{M}_2^* correspond to M_1 and M_2 respectively. From this matter and the assumption of this theorem, we can give truth-values for propositions $F_1, F_2, \dots, F_\alpha; G_1^1, G_2^1, \dots, G_\beta^1; G_1^2, G_2^2, \dots, G_\beta^2$ in $\tilde{\mathfrak{U}}^*$ by which $\tilde{\mathfrak{U}}^*$ takes $(0, 0, 0, 0, 0, \dots)$ at every (x, y, z) in $N \times N \times N$.

Here we make a correspondence of $(0, 0, 0, 0, 0, \dots), (1, 1, 1, 1, 1, \dots)$ to F, T as mentioned above. Then, we consider only values at (i, i, i) in $N \times N \times N$ in the assignment where $i = 1, 2, 3, \dots$.

Now, let us say that the truth-value assignment is as follows:

$$\begin{array}{ll}
 F_1(e_1): (t_{111}, t_{112}, \dots) & F_2(e_1): (t_{211}, t_{212}, \dots) \dots \\
 F_1(e_2): (t_{121}, t_{122}, \dots) & F_2(e_2): (t_{221}, t_{222}, \dots) \\
 \vdots & \\
 G_1^1(e_1): (\tau_{111}^1, \tau_{112}^1, \dots) & G_1^2(e_1): (\tau_{111}^2, \tau_{112}^2, \dots) \dots \\
 G_1^1(e_2): (\tau_{121}^1, \tau_{122}^1, \dots) & G_1^2(e_2): (\tau_{121}^2, \tau_{122}^2, \dots) \\
 \vdots & \vdots
 \end{array}$$

where $F_1(e_1), F_1(e_2), \dots$ mean values of F_1 at $(1, 1, 1), (2, 2, 2), \dots$ in $N \times N \times N$ as before.

Then, we take $(1, 1, 1), (2, 2, 2), \dots$ as an infinite domain ω . Further we take a value of $\diamond XF_i$ obtained from $F_i(e_i)$ as a truth-value of predicate $F_i(x)$ for $x = (i, i, i)$ and also a value of $\diamond (XG_i^1 \wedge YG_i^2)$ obtained from $G_i^1(e_i),$

$G_i^2(e_i)$ as a truth-value of predicate $G_i(x, y)$ for $x = (k, k, k)$, $y = (l, l, l)$ and so on. From the definitions of X, Y, Z and the $h: h(F_i(x)) \rightarrow \diamond XF_i$, $h(G_i(x, y)) \rightarrow \diamond (XG_i^1 \wedge YG_i^2)$, \dots , we know that \mathfrak{A}^* takes F in the enumerable infinite domain ω .

Thus, we get Theorem 2'.

*Faculty of Engineering
Hiroshima University*