

closely with the figure for institutional cases. In the direct line there was a neuropathic taint in 68%. In the direct line psychoses and nervous disorders were more frequent in extra-mural cases; alcoholism, psychopathic personality and suicide were less frequent. The author considers that the very mild and brief cases of manic-depressive depression, frequently not recognized as such, are in their nature identical with the classic institutional cases.

G. W. T. H. FLEMING.

*Mental and Emotional Phenomena of Some Psychoses in their Relation to Blood-pressure; Diagnostic and Prognostic Significance of the Latter.* (*Journ. of Nerv. and Ment. Dis.*, October, 1930.) Gordon, A.

The author investigated 37 cases of psychosis. All gave a lowering of blood-pressure half an hour after the injection of 200 grm. of milk. There was no alteration in a control group of 15 normal individuals and 7 obsessional cases. In the manic cases the difference between the terminal ascent and the original blood-pressure was smaller than in the depressives. There appeared to be a relationship between these differences and the duration of each phase of a manic-depressive psychosis; the smaller the difference the longer the manic phase lasted, and the greater the difference the longer the depressive phase lasted. In the confusional cases there was a striking similarity to the depressive phases of the manic-depressive psychosis. In the dementia præcox group, during the gradual descent of blood-pressure there was a considerable abatement in restlessness, especially in the automatic and stereotyped movements. In the cases of anxiety neurosis the lowering of the blood-pressure was accompanied by an amelioration of the main symptoms.

G. W. T. H. FLEMING.

*Post-Encephalitic Parkinsonism with Psychosis.* (*Journ. of Nerv. and Ment. Dis.*, September, 1930.) Alpers, B. A.

The author thinks that the number of cases of post-encephalitic Parkinsonism which develop a psychosis is small. He describes three cases in his own practice which had similar mental features. All presented hallucinations, chiefly auditory but sometimes visual. There were few other mental disturbances. Hallucinations have not often been reported in the psychosis complicating Parkinsonism.

G. W. T. H. FLEMING.

*Personality Factors in Alcoholism.* (*Arch. of Neur. and Psychiat.*, July, 1930.) Hart, H. H.

The author examined 30 cases at the Blythwood Sanatorium. Most of the patients were typically American; there were no Jews. In all cases there was present a constitutional instability, with parental discord or alcoholism, and lack of discipline and sensible direction in the environment of childhood. Alcohol provides its own Nemesis. Alcoholism is an emotional and moral problem. The treatment, so far as it is possible, resolves itself into building