



Fibrations in semitoric and generalized complex geometry

Gil R. Cavalcanti, Ralph L. Klaasse, and Aldo Witte

Abstract. This paper studies a class of singular fibrations, called self-crossing boundary fibrations, which play an important role in semitoric and generalized complex geometry. These singular fibrations can be conveniently described using the language of Lie algebroids. We will show how these fibrations arise from (nonfree) torus actions, and how to use them to construct and better understand self-crossing stable generalized complex four-manifolds. We moreover show that these fibrations are compatible with taking connected sums, and use this to prove a singularity trade result between two types of singularities occurring in these types of fibrations (a so-called nodal trade).

1 Introduction

1.1 Fibrations in geometry

A general theme in the world of geometric structures is the interplay between specific geometric structures and particular types of maps. The simplest example arises in the context of fibrations, where one studies whether the presence of a geometric structure on the base and the fiber implies its existence on the total space. However, to obtain interesting spaces and geometric structures, one should allow maps to have more singularities.

A concrete case that illustrates this comes from symplectic geometry: symplectic fibrations play an important role, but they are quite rare on general symplectic manifolds. Instead, if one allows the fibration to have Lefschetz singularities, one obtains enough flexibility to establish broad existence results [13, 16, 17].

In this paper, we use the point of view of decoupling the differential properties of maps from the desired underlying geometric structure. This decoupling has proved useful and allowed for several extensions of the results mentioned above, including those in [1–6, 9, 10, 14, 15, 23]. Here, allowing for maps to have Lefschetz and other similar singularities leads to existence results for several different types of geometric structures.

Received by the editors August 2, 2021; revised January 5, 2022; accepted March 15, 2022.

Published online on Cambridge Core March 29, 2022.

R.L.K. was supported by ERC consolidator grant 646649 “SymplecticEinstein,” and by the FWO and FNRS under EOS Project No. G0H4518N. A.W. was supported by the NWO through the UGC Graduate Program.

AMS subject classification: 53D18, 53D17, 53D05.

Keywords: Generalized complex structure, elliptic symplectic structure, Lie algebroid, boundary Lefschetz fibration, toric geometry, semi-toric systems.

Another way in which singular fibrations arise is from proper group actions. In this setting, the “fibration” condition translates to the group action being free, which is restrictive. The quotient map of a generic action will have singularities at points with nontrivial isotropy. Such singular fibrations are particularly well studied for torus actions, $T^n \times M^{2n} \rightarrow M^{2n}$, where, even if the actions considered are not free, they are well behaved enough to ensure that the quotient space admits the structure of a manifold with corners. The coupling of torus actions with geometric structures leads to many fruitful concepts, one of the highlights being toric geometry.

The Lefschetz and the toric pictures come together in semitoric geometry [28], where maps are allowed to have both types of singularities. In this setting, the maps can have three types of singularities: elliptic, elliptic–elliptic, and focus–focus, with the latter being equivalent to Lefschetz singularities. However, in semitoric geometry, the decoupling of maps and geometric structures has happened only partially, because local singular behavior (Lefschetz and toric) and more global properties (integral affine structures) are mixed, leading to topological results [24].

In this paper, we introduce the differential objects hinted at by semitoric geometry: these are called *self-crossing boundary (Lefschetz) fibrations* (cf. Definitions 3.11 and 3.12). These types of maps incorporate local phenomena from both symplectic fibrations and quotient maps of semitoric manifolds, but do not require global structures to be present, such as group actions or integral affine structures. In this paper, we use these boundary fibrations to study an a priori seemingly unrelated geometric structure, namely a generalized complex structure.

1.2 Generalized complex structures

Generalized complex structures [20, 22] are a simultaneous generalization of symplectic and complex structures. Infinitesimally, these structures induce the product of a symplectic and complex vector space on each tangent space. However, the number of complex and symplectic directions, called the *type*, can vary from point to point, leading to the notion of type change. These type-changing generalized complex structures are among the most interesting to study. Within the type-changing generalized complex structures, one class was put forward in [8, 11] for being geometrically very rich and having well-controlled singular behavior: *self-crossing stable generalized complex structures*.

It is shown in [11] that self-crossing stable generalized complex structures on a manifold M are in one-to-one correspondence with certain Lie algebroid symplectic structures, so that this paper makes extensive use of Lie algebroids. The singularities at the type-change locus D induce a Lie algebroid $\mathcal{A}_{|D} \rightarrow M$ called the *self-crossing elliptic tangent bundle*, and the generalized complex structure makes it into a symplectic Lie algebroid, carrying an *elliptic symplectic structure*. An elliptic symplectic structure corresponds to a self-crossing stable generalized complex structure if the locus D is co-orientable and its so-called *index* is 1.

Given a self-crossing boundary Lefschetz fibration $f: (M, D) \rightarrow (N, Z)$ where Z is a hypersurface of N , there is another relevant Lie algebroid, namely the *self-crossing log-tangent bundle* $\mathcal{A}_Z \rightarrow N$. The map f has singularities precisely such that it induces

a Lie algebroid morphism

$$(\varphi, f): (\mathcal{A}_{|D|}, M) \rightarrow (\mathcal{A}_Z, N),$$

where φ is now a Lie algebroid version of a Lefschetz fibration. The relevant geometric structure on the base of this fibration is a symplectic structure on \mathcal{A}_Z , also known as a *self-crossing log-symplectic structure*. These have also appeared in [21, 26]. They are compatible with the elliptic symplectic structure in the following sense.

Definition 3.20 A self-crossing boundary fibration $f: (M, D) \rightarrow (N, Z)$ is *compatible* with the elliptic symplectic structure on its total space if $\ker \varphi \subseteq \mathcal{A}_{|D|}$ consists of symplectic subspaces.

In turn, a stable generalized complex structure is *compatible* with a boundary Lefschetz fibration if its induced elliptic symplectic structure is.

1.3 Results

In this section, we describe the main results obtained in this paper. In the interest of brevity, more precise versions of the results below can be found in the main body of the text.

1.3.1 Existence

Following the strategy of constructing symplectic structures out of fibrations, we prove a Gompf–Thurston theorem for self-crossing stable generalized complex structures. This result is the generalization of a similar result for stable generalized complex structures with embedded type-change locus appearing in [9], but requires several adaptations of the argument.

Definition 3.19 A boundary Lefschetz fibration, $f: (M^4, D) \rightarrow (N^2, Z)$, is *homologically essential* if the homology class $[F]$ of the fiber of $f: M \setminus D \rightarrow N \setminus Z$ is nontrivial in $H_2(M \setminus D; \mathbb{R})$.

Theorem 3.23 Let $f: (M^4, D^2) \rightarrow (N^2, \partial N)$ be a homologically essential self-crossing boundary Lefschetz fibration. Then M^4 admits an elliptic symplectic structure compatible with f , which induces a self-crossing stable generalized complex structure compatible with f if the locus D is co-orientable and its index is equal to 1.

1.3.2 Construction

Having established that boundary Lefschetz fibrations supply self-crossing stable generalized complex structures, we decouple the map from the geometric structure and study them separately. These types of maps are flexible enough to admit connected sums.

Theorem 4.6 Let $f_i: (M_i^4, D_i^2) \rightarrow (N_i^2, \partial N_i)$, for $i = 1, 2$, be boundary Lefschetz fibrations, and let $p_i \in M_i$ be such that $q_i = f_i(p_i)$ are corners of the manifolds N_i . Then there exists a boundary Lefschetz fibration on their connected sum,

$$f_1 \# f_2: (M_1 \#_{p_1, p_2} M_2, D_1 \# D_2) \rightarrow (N_1 \#_{q_1, q_2} N_2, \partial(N_1 \# N_2)),$$

which is compatible with the inclusion $M_i \setminus \{p_i\} \hookrightarrow M_1 \# M_2$. Moreover, the map $f_1 \# f_2$ is homologically essential if and only if f_1 and f_2 are.

This result is in sheer contrast with the situation in toric geometry. There is no symplectic connected sum procedure, and most of the manifolds obtained using the above proposition will have no toric structure. This difference in rigidity between the generalized complex and toric worlds is already apparent on the base of these fibrations. Namely, for generalized complex structures, the base carries a self-crossing log-symplectic structure, which is quite flexible. On the other hand, in toric geometry, the base carries an integral affine structure, which is very rigid. In other words, although toric manifolds do not behave well with respect to connected sums, the underlying torus actions and abstract quotient maps do.

1.3.3 Singularity trades

The nodal-trade procedure in semitoric geometry exchanges elliptic–elliptic singularities of the moment map for focus–focus singularities [32, 33] and vice-versa [24]. These procedures rely heavily on the existence of a singular integral affine structure on the base. Following our general strategy, decoupling the geometric structure from the maps allows us to prove an abstract statement for boundary Lefschetz fibrations.

Theorem 5.3 *Let $f: (M^4, D^2) \rightarrow (N^2, \partial N)$ be a boundary Lefschetz fibration, and let $p \in M$ be an elliptic–elliptic singularity. Then there exists a boundary Lefschetz fibration*

$$\tilde{f}: (M, \tilde{D}) \rightarrow (\tilde{N}, \partial \tilde{N})$$

agreeing with f outside a neighborhood of p , and such that the elliptic–elliptic singularity is traded for a Lefschetz singularity. The map \tilde{f} is homologically essential if and only if f is.

The proof of this result, and its converse, Theorem 5.4, relies on the connected sum procedure and the existence of a particular boundary Lefschetz fibration on S^4 . Figure 1 illustrates the singularity trade from Theorem 5.3 in $\mathbb{C}P^2$.

1.3.4 Examples

Using simple manifolds as building blocks, the connected sum procedure allows us to construct many examples of boundary Lefschetz fibrations (and consequently of self-crossing stable generalized complex structures) on the following manifolds.

Theorem 6.12 *The manifolds in the following two families:*

- $X_{n,\ell} := \#n(S^2 \times S^2) \# \ell(S^1 \times S^3)$, with $n, \ell \in \mathbb{N}$,
- $Y_{n,m,\ell} := \#n\mathbb{C}P^2 \# m\overline{\mathbb{C}P}^2 \# \ell(S^1 \times S^3)$, with $n, m, \ell \in \mathbb{N}$,

admit homologically essential boundary Lefschetz fibrations whenever their Euler characteristic is nonnegative. Therefore, each of these manifolds admits a compatible elliptic symplectic structure, which induces a self-crossing stable generalized complex structure if $1 - b_1 + b_2^+$ is even.

Combining this result with Theorem 5.3, we conclude that the above manifolds moreover admit stable generalized complex structures with embedded degeneracy




Figure 1: The picture on the left presents $\mathbb{C}P^2$ using the usual moment map, which is the prototypical example of a self-crossing boundary fibration. Theorem 5.3 tells us that we can slightly modify this fibration to obtain a boundary Lefschetz fibration, with three Lefschetz singularities.

loci. These examples have already appeared in the literature [7, 18, 30, 31], and the authors obtained them as well in [11, Theorem 7.5]. However, the above result shows that the structures in these examples can be made compatible with boundary Lefschetz fibrations. This gives us much more control about stable generalized complex structures in concrete applications. For instance, we believe it will help in the development of a Fukaya category for stable generalized complex structures just like ordinary Lefschetz fibrations can be used to understand the Fukaya category of symplectic manifolds [29].

1.4 Organization of the paper

This paper is organized as follows. In Section 2, we recall from [11] the notions of self-crossing divisors, and their associated Lie algebroids and symplectic structures. We also recall the definition of self-crossing stable generalized complex structures and that they are in one-to-one correspondence with particular self-crossing elliptic symplectic structures. In Section 3, we extend the notion of boundary (Lefschetz) fibration from [9] to allow for self-crossing of the degeneracy locus. We moreover prove the Gompf–Thurston result, Theorem 3.23. In Section 4, we show that boundary Lefschetz fibrations allow for taking connected sums and prove Theorem 4.6. In Section 5, we will prove the singularity trade results, namely Theorems 5.3 and 5.4. Finally, in Section 6, we show that torus actions give rise to boundary fibrations, and exhibit several examples, including Theorem 6.12 and Proposition 6.9.

2 Divisors, Lie algebroids, and symplectic structures

In this section, we study geometric structures with specific singularities. To work with these singularities, we will recall the concept of a divisor, and the particular cases of log and elliptic divisors which we will mainly use in this paper. Using these divisors, we will recall the associated Lie algebroids and their Lie algebroid symplectic structures. Then we will introduce the objects which we want to construct in this paper, namely stable generalized complex structures. We will show that these structures correspond to certain Lie algebroid symplectic structures, which is how we will treat them in the rest of the paper.

2.1 Divisors

We will use an adaptation of the notion of a divisor from algebraic geometry to smooth manifolds in order to describe the singularities of geometric structures. We will only briefly go over the main concepts we need and refer to [8, 9, 11] for more information.

Definition 2.1 A real/complex *divisor* on M is a locally principal ideal I of $C^\infty(M; \mathbb{R})$, respectively, $C^\infty(M; \mathbb{C})$, which is locally generated by functions with nowhere dense zero set.

Equivalently, divisors can be described as follows.

Proposition 2.2 Let I be a real/complex divisor on M . Then there exists a real/complex line bundle $L \rightarrow M$ with section $\sigma \in \Gamma(L)$ such that $\sigma(\Gamma(L^*)) = I$.

Note that the line bundle L is uniquely determined up to vector bundle isomorphism (covering the identity), and that the section σ is unique up to multiplication by a smooth function. Given a pair (L, σ) , we denote the associated divisor by I_σ .

Definition 2.3 Let (M, I_M) and (N, I_N) be manifolds with divisors. A smooth map $\varphi: M \rightarrow N$ is a *morphism of divisors* if $\varphi^* I_N = I_M$, where the left-hand side denotes the ideal generated by all pullbacks. It is called a *diffeomorphism of divisors* if φ is a diffeomorphism.

Definition 2.4 A *smooth real/complex log divisor* is a real/complex divisor I locally generated by transverse vanishing functions.

The *vanishing locus* of a real log divisor has codimension 1 and is denoted by Z . The vanishing locus of a complex log divisor has codimension 2 and is denoted by D . By locally demanding a divisor to be a product of log divisors, we obtain the following.

Definition 2.5 A *self-crossing real/complex log divisor* on a manifold M is a divisor I , such that for every point $p \in M$, there exists a neighborhood U of p such that

$$I(U) = I_1 \cdot \dots \cdot I_j,$$

where I_1, \dots, I_j are real/complex log ideals with transversely intersecting vanishing loci.

Example 2.6 There is a standard example for each of the divisor types described above.

The *standard real log divisor* on $\mathbb{R}^j \times \mathbb{R}^m$ is defined using the coordinates (x_1, \dots, x_j, y_i) by the ideal $I_Z := \langle x_1 \cdot \dots \cdot x_j \rangle$;

The *standard complex log divisor* on $\mathbb{C}^j \times \mathbb{R}^m$ is defined using the coordinates (z_1, \dots, z_j, y_i) by the ideal $I_D := \langle z_1 \cdot \dots \cdot z_j \rangle$.

Each of these examples provides the local normal form for their associated divisor type.

A self-crossing real log divisor is determined by its vanishing locus Z , as its ideal equals the ideal of functions vanishing on Z . In contrast, for a self-crossing complex log divisor, the subspace D does not determine the divisor.

Definition 2.7 [8] A smooth elliptic divisor is a real divisor $I_{|D|}$ locally generated by definite Morse–Bott functions with codimension-2 critical set.

In fact, asking the existence of local generating function of $I_{|D|}$ as above implies that there exists a global generator $f \in I_{|D|}$.

Again by taking appropriate products, we obtain the following.

Definition 2.8 [11] A self-crossing elliptic divisor is a divisor $I_{|D|}$ on a manifold M , such that for every point $p \in M$, there exists a neighborhood U of p such that

$$I_{|D|}(U) = I_1 \cdot \dots \cdot I_j,$$

where the I_1, \dots, I_j are smooth elliptic divisors with transversely intersecting vanishing loci.

Example 2.9 The standard elliptic divisor on $\mathbb{R}^{2j} \times \mathbb{R}^m$ is defined using the coordinates $(x_1, y_1, \dots, x_j, y_j, u_i)$ by the ideal $I_{|D|} := \left((x_1^2 + y_1^2) \cdot \dots \cdot (x_j^2 + y_j^2) \right)$. Lemma 2.23 in [11] ensures that any self-crossing elliptic divisor is of this form.

As for smooth elliptic divisors, the ideal $I_{|D|}$ is generated by a single global function $f: M \rightarrow \mathbb{R}_+$, which by Example 2.9 is locally of the form

$$f(x_1, y_1, \dots, x_j, y_j, x_{k+1}, \dots, x_n) = (x_1^2 + y_1^2) \dots (x_j^2 + y_j^2).$$

We mostly deal with self-crossing divisors in this paper, and we will often omit the prefix “self-crossing.” Whenever we mean a smooth log or elliptic divisor, we will explicitly stress this.

The vanishing loci of both log and elliptic divisors are not embedded, but are stratified.

Definition 2.10 Let I be a real/complex log or elliptic divisor on M with vanishing locus D . The multiplicity of a point $p \in M$ is the minimum of the integers j from Definitions 2.5 or 2.8 over all neighborhoods U of p . If I has points of multiplicity at most n , the sets $D(j)$ of points of multiplicity at least j induce a filtration of M :

$$M = D(0) \supset D(1) = D \supset \dots \supset D(n),$$

with induced stratification with strata $D[j]$ of points with multiplicity precisely j .

That this is a stratification follows readily from the normal forms of the divisors in Examples 2.6 and 2.9. Also, note that if a divisor I has multiplicity n and $i \leq n$ is given, then the restriction $I|_{M \setminus D(i+1)}$ defines a divisor with multiplicity i .

Example 2.11 Another important example for this paper is a manifold with corners $(M, \partial M)$. The boundary of a manifold with corners naturally defines a real log divisor.

Given a self-crossing complex log divisor I_D , its associated (self-crossing) elliptic divisor is the real divisor $I_{|D|}$ defined by $I_{|D|} \otimes \mathbb{C} = I_D \otimes \overline{I_D}$.

2.2 Lie algebroids and residue maps

Each of the divisors introduced in the previous section gives rise to a corresponding Lie algebroid via the Serre–Swan theorem and the local normal forms contained in Examples 2.6 and 2.9.

Definition 2.12 Let I_Z be a real log divisor, I_D a complex log divisor, and $I_{|D|}$ be an elliptic divisor. The vector fields preserving each of these ideals define Lie algebroids:

- $\mathcal{A}_Z \rightarrow TM$, the *real log-tangent bundle* associated to I_Z ;
- $\mathcal{A}_D \rightarrow T_{\mathbb{C}}M$, the *complex log-tangent bundle* associated to I_D ; and
- $\mathcal{A}_{|D|} \rightarrow TM$, the *elliptic tangent bundle* associated to $I_{|D|}$.

Remark 2.13 The above Lie algebroids can be described in the local coordinates of in Examples 2.6 and 2.9. Indeed, around a point of multiplicity j , we have:

- $\Gamma(\mathcal{A}_Z) = \langle x_1\partial_{x_1}, \dots, x_j\partial_{x_j}, \partial_{y_i} \rangle$;
- $\Gamma(\mathcal{A}_D) = \langle z_1\partial_{z_1}, \partial_{\bar{z}_1}, \dots, z_j\partial_{z_j}, \partial_{\bar{z}_j}, \partial_{y_i} \rangle$; and
- $\Gamma(\mathcal{A}_{|D|}) = \langle r_1\partial_{r_1}, \partial_{\theta_1}, \dots, r_j\partial_{r_j}, \partial_{\theta_j}, \partial_{u_i} \rangle$.

In the latter case, we have $r_i\partial_{r_i} := x_i\partial_{x_i} + y_i\partial_{y_i}$ and $\partial_{\theta_i} := x_i\partial_{y_i} - y_i\partial_{x_i}$.

When I_D is a complex log divisor on M and $I_{|D|}$ is its associated elliptic divisor, there is a fiber product relation between the corresponding Lie algebroids as follows:

$$\mathcal{A}_D \times_{T_{\mathbb{C}}M} \mathcal{A}_{\overline{D}} \cong \mathcal{A}_{|D|} \otimes \mathbb{C}.$$

This isomorphism provides an inclusion $\iota^*: \Omega^\bullet(\mathcal{A}_D) \rightarrow \Omega_{\mathbb{C}}^\bullet(\mathcal{A}_{|D|})$.

We now turn to describing several of the residue maps carried by these Lie algebroids. Let $(I_{|D|}, \sigma)$ be a smooth elliptic divisor, together with a co-orientation of D . As explained in [8], the elliptic tangent bundle has an *elliptic* and *radial residue* map. These are maps of cochain complexes, and they extract the coefficients of the singular generators. In the coordinates of Remark 2.13, these are given by

$$(2.1) \quad \begin{aligned} \text{Res}_q: \Omega^\bullet(\mathcal{A}_{|D|}) &\rightarrow \Omega^{\bullet-2}(D), & \text{Res}_q(\alpha) &= \iota_D^*(\iota_{r\partial_r \wedge \partial_\theta} \alpha), \\ \text{Res}_r: \Omega^\bullet(\mathcal{A}_{|D|}) &\rightarrow \Omega^{\bullet-1}(S^1ND), & \text{Res}_r(\alpha) &= \iota_D^*(\iota_{r\partial_r} \alpha), \end{aligned}$$

where $S^1ND \rightarrow D$ is the S^1 -bundle associated to the co-orientation σ of D .

These residue maps can be extended to self-crossing elliptic divisors if we restrict our attention to the stratum $D[1]$. We say that a self-crossing elliptic divisor is *co-oriented* if the normal bundle $ND[1] \rightarrow D[1]$ is oriented.

Definition 2.14 Let $(I_{|D|}, \sigma)$ be a self-crossing elliptic divisor together with a co-orientation σ of $D[1]$. The *elliptic* and *radial residues* of $\alpha \in \Omega^\bullet(\mathcal{A}_{|D|})$ are $\text{Res}_q(\alpha) := \text{Res}_q(\iota_{D[1]}^* \alpha)$ and $\text{Res}_r(\alpha) := \text{Res}_r(\iota_{D[1]}^* \alpha)$.

In later constructions, the cohomology of the complex of forms with vanishing radial residue will play a role.

Lemma 2.15 Let $I_{|D|}$ be a self-crossing elliptic divisor on a manifold M , and let $\Omega_{0,0}^\bullet(\mathcal{A}_{|D|}) \subset \Omega^\bullet(\mathcal{A}_{|D|})$ be the subcomplex defined as the kernel of the map Res_r . Then the inclusion map $\iota: M \setminus D \rightarrow M$ of the complement of D induces a quasi-isomorphism $\iota^*: \Omega_{0,0}^\bullet(\mathcal{A}_{|D|}) \rightarrow \Omega^\bullet(M \setminus D)$.

Proof The argument uses the observation from [19, Theorem 1.2] (and the fact that $\mathcal{A}_{|D|}$ defines a soft sheaf) that it suffices to show that ι^* induces an isomorphism on the level of sheaf cohomology. Below we will implicitly identify the sheaf $\Omega^\bullet(M \setminus D)$ with its push-forward $\iota_*(\Omega^\bullet(M \setminus D))$. For all points $p \in M \setminus D$, there exists a contractible

open neighborhood U of p disjoint from D . On this open $\mathcal{A}_{|D|} = TM$ and $\text{Res}_r \equiv 0$, and therefore ι_* is simply the identity. Let j be any integer less than or equal to the point of highest multiplicity of D , take $p \in D[j]$, and let U be a contractible open around p as in Remark 2.13. In those coordinates, $H_{0,0}^\bullet(U, \mathcal{A}_{|D|})$ is the free algebra generated by $\{1, d\theta_1, \dots, d\theta_j\}$. By an elementary argument, $U \setminus D$ is homotopic to \mathbb{T}^j , and using this homotopy ι^* takes the generators of $H_{0,0}^\bullet(\mathcal{A}_{|D|})$ to the generators of $H^\bullet(U \setminus D)$. Therefore, we conclude that ι^* is a local isomorphism, and consequently a global isomorphism. ■

We will need a few more residue maps for self-crossing elliptic divisors.

Definition 2.16 [11] Let $(I_{|D|}, \mathfrak{o})$ be a co-oriented elliptic divisor, and let $\omega \in \Omega^2(\mathcal{A}_{|D|})$. Let $p \in D(k)$ with $k \geq 2$ and consider oriented coordinates as in Remark 2.13. We define

$$\begin{aligned} \text{Res}_{r_i r_j} \omega(p) &:= \omega_p(r_i \partial_{r_i}, r_j \partial_{r_j}), \\ \text{Res}_{r_i \theta_j} \omega(p) &:= \omega_p(r_i \partial_{r_i}, \partial_{\theta_j}), \quad \text{Res}_{\theta_i \theta_j} \omega(p) := \omega_p(\partial_{\theta_i}, \partial_{\theta_j}). \end{aligned}$$

These pointwise expressions depend on \mathfrak{o} and the ordering of coordinates, but only up to sign.

2.3 Lie algebroid symplectic structures

We will use the language of symplectic Lie algebroids to translate certain Poisson and generalized complex structures into simpler Lie algebroid objects. Given a Lie algebroid two-form $\omega \in \Omega^2(\mathcal{A})$, we say it is nondegenerate if $\omega^b: \mathcal{A} \rightarrow \mathcal{A}^*$ is an isomorphism.

Definition 2.17 Let I_Z and $I_{|D|}$ be real log and elliptic divisors on a given manifold M . Then:

- A form $\omega \in \Omega^2(\mathcal{A}_Z)$ is *log-symplectic* if $d\omega = 0$ and it is nondegenerate.
- A form $\omega \in \Omega^2(\mathcal{A}_{|D|})$ is *elliptic symplectic* if $d\omega = 0$ and it is nondegenerate.

One can prove Darboux-type normal form theorems for symplectic Lie algebroids using a thorough understanding of their Lie algebroid cohomology, by a straightforward adaptation of the Moser lemma. However, in the above cases, this cohomology is generally locally nontrivial, so that there is no unique local model. In dimension 2, we have the following (which is the real analogue of Proposition 5.2 in [11]).

Lemma 2.18 Let I_Z be a real log divisor on Σ^2 , and let $\omega \in \Omega^2(\mathcal{A}_Z)$ be a log-symplectic form. For each point $p \in Z[2]$, there are coordinates (x_1, x_2) centered at p and $\lambda \in \mathbb{R}$ such that

$$\omega = \lambda d \log x_1 \wedge d \log x_2.$$

Since in two dimensions every nowhere zero two-form is closed and nondegenerate, we have the following source of examples of log-symplectic manifolds.

Lemma 2.19 Let Σ^2 be a compact oriented surface with corners. Then $(\Sigma, I_{\partial\Sigma})$ admits a log-symplectic structure.

Proof The ideal $I_{\partial\Sigma}$ defines a real log divisor. Because Σ is oriented, let $h \in C^\infty(M)$ be a defining function for ∂M , so that $I_{\partial\Sigma} = \langle h \rangle$, and let $\omega \in \Omega^2(\Sigma)$ be a volume form. Then $h^{-1}\omega \in \Omega^2(\mathcal{A}_{\partial M})$ is a nondegenerate log two-form that is closed for dimensional reasons. ■

2.4 Self-crossing stable generalized complex structures

In this section, we recall the notion of a self-crossing stable generalized complex structure [11]. This is a well-behaved class of generalized complex structures [20], i.e., complex structures on the bundle $\mathbb{T}M := TM \oplus T^*M$. We furthermore recall that they are equivalent to certain elliptic symplectic structures.

Definition 2.20 A *generalized complex structure* on a manifold M is a pair (\mathbb{J}, H) where $H \in \Omega^3(M)$ is a closed three-form and \mathbb{J} is a skew-symmetric endomorphism of $\mathbb{T}M$ for which $\mathbb{J}^2 = -\text{Id}$ and the $+i$ -eigenbundle, $L \subset (\mathbb{T}M) \otimes \mathbb{C}$, is involutive with respect to the Dorfman bracket:

$$[[X + \xi, Y + \eta]]_H := [X, Y] + \mathcal{L}_X \eta - \iota_Y d\xi + \iota_X \iota_Y H, \quad X + \xi, Y + \eta \in \Gamma(\mathbb{T}M).$$

Given a generalized complex structure (\mathbb{J}, H) on M , one can decompose it in a two-by-two block matrix, using the decomposition $\mathbb{T}M = TM \oplus T^*M$. The skew-symmetry of \mathbb{J} ensures that it is of the form $\mathbb{J} = \begin{pmatrix} J & \pi_{\mathbb{J}}^\sharp \\ \sigma^b & -J^* \end{pmatrix}$ with $J \in \mathcal{J}(TM)$, $\pi_{\mathbb{J}}^\sharp : T^*M \rightarrow TM$ corresponding to a bivector $\pi_{\mathbb{J}} \in \mathfrak{X}^2(M)$ and $\sigma^b : TM \rightarrow T^*M$ corresponding to a two-form $\sigma \in \Omega^2(M)$.

Two generalized complex structures (\mathbb{J}, H) and (\mathbb{J}', H') are *gauge equivalent* if there exists $B \in \Omega^2(M)$ such that $H' = H + dB$ and, using the associated map $B^\flat : TM \rightarrow T^*M$, we have

$$\mathbb{J}' = \begin{pmatrix} 1 & 0 \\ B^\flat & 1 \end{pmatrix} \mathbb{J} \begin{pmatrix} 1 & 0 \\ -B^\flat & 1 \end{pmatrix}.$$

Lemma 2.21 [12] Let $\mathbb{J} = \begin{pmatrix} J & \pi_{\mathbb{J}}^\sharp \\ \sigma^b & -J^* \end{pmatrix}$ be a generalized complex structure on M . Then $\pi_{\mathbb{J}} \in \mathfrak{X}^2(M)$ is a Poisson structure on M . Moreover, if \mathbb{J} and \mathbb{J}' are gauge equivalent, then $\pi_{\mathbb{J}} = \pi_{\mathbb{J}'}$.

Given an element $X + \xi \in \mathbb{T}_{\mathbb{C}}M$, let $(X + \xi) \cdot \rho := \iota_X \rho + \xi \wedge \rho$ denote the Clifford action of $\mathbb{T}M$ on elements $\rho \in \wedge^\bullet T_{\mathbb{C}}^*M$. A generalized complex structure \mathbb{J} is alternatively characterized by its *canonical bundle* $K \subset \wedge^\bullet T_{\mathbb{C}}^*M$ defined by the relation

$$L = \{u \in \mathbb{T}_{\mathbb{C}}M : u \cdot K = 0\}.$$

Its dual carries a natural section $s \in \Gamma(K^*)$, given by the map which sends $\rho \in \Gamma(K)$ to its degree zero part, and is called the *anticanonical section* of \mathbb{J} . The pair (K^*, s) , called the *anticanonical divisor*, can be used to define a specific class of generalized complex structures:

Definition 2.22 [11] Let M be a manifold and $H \in \Omega^3(M)$ closed. A generalized complex structure \mathbb{J} on (M, H) is *stable with self-crossings* if (K^*, s) defines a self-crossing complex log divisor.

As before, we will often simply call these structures “stable,” and when their divisor is in fact smooth, we will explicitly stress this.

If \mathbb{J} is a stable generalized complex structure on M , one can show that $\pi_{\mathbb{J}}$ admits a nondegenerate lift to $\mathcal{A}_{|D|}$, the elliptic tangent bundle with respect to the elliptic divisor induced by (K^*, s) . Inverting this nondegenerate lift results in an elliptic symplectic form $\omega \in \Omega^2(\mathcal{A}_{|D|})$. Under certain conditions, this procedure can be reversed.

Theorem 2.23 [11] *Let M be a manifold. There is a one-to-one correspondence between gauge equivalence classes of stable generalized complex structures on M and co-oriented elliptic divisors together with an elliptic symplectic form $\omega \in \Omega^2(\mathcal{A}_{|D|})$, satisfying:*

- $\text{Res}_q(\omega) = 0$.
- $\text{Res}_{\theta_i r_j}(\omega) = \text{Res}_{r_i \theta_j}(\omega)$.
- $\text{Res}_{r_i r_j}(\omega) = -\text{Res}_{\theta_i \theta_j}(\omega)$.

Explicitly, this map is given by

$$\left\{ \begin{array}{l} (\mathbb{J}, H): \\ \mathbb{J} \text{ is a stable GCS} \end{array} \right\} \rightarrow \left\{ \begin{array}{l} (I_{|D|}, \mathfrak{o}, \pi_{\mathbb{J}}^{-1}): \\ (I_{|D|}, \mathfrak{o}) \text{ is a co-oriented elliptic divisor and} \\ \pi_{\mathbb{J}}^{-1} \text{ is an elliptic symplectic form satisfying the above relations} \end{array} \right\}.$$

Here, $(I_{|D|}, \mathfrak{o})$ is the co-oriented elliptic divisor induced by the anti-canonical divisor.

In the above, the co-orientation \mathfrak{o} is defined using the fact that the normal derivative of the anti-canonical section s induces an isomorphism $d^v s: ND|_{D[1]} \simeq K^*|_{D[1]}$.

Example 2.24 Consider the generalized complex structure on \mathbb{C}^2 with trivial canonical bundle determined by the form

$$\rho = z_1 z_2 + \tau dz_1 \wedge dz_2, \quad \tau \in \mathbb{C}.$$

In terms of the dual section of $\rho^* \in \Gamma(K^*)$, the anticanonical section is given by $s = z_1 z_2 \rho^* \in \Gamma(K^*)$, and therefore ρ defines a stable structure with elliptic ideal $|z_1|^2 |z_2|^2$. The elliptic symplectic form induced by ρ is

$$\omega = \text{Im}(\tau)(d \log r_1 \wedge d \log r_2 - d\theta_1 \wedge d\theta_2) + \text{Re}(\tau)(d \log r_1 \wedge d\theta_2 + d\theta_1 \wedge d \log r_2).$$

This structure provides the normal form for a four-dimensional stable generalized complex structure around a point in $D[2]$.

In this paper, we will predominantly consider examples of stable generalized complex manifolds for which the local normal form has parameter $\tau = i\lambda$ for a real number λ ; thus, it is worth recalling the following definition.

Definition 2.25 Let M^4 be a four-dimensional manifold endowed with an elliptic divisor. We say $\omega \in \Omega^2(\mathcal{A}_{|D|})$ with zero elliptic residue has *imaginary parameter* at a point $p \in D[2]$ if:

- $|\text{Res}_{r_1 r_2}(\omega)(p)| = |\text{Res}_{\theta_1 \theta_2}(\omega)(p)|$.
- $\text{Res}_{r_1 \theta_2} \omega(p) = 0$.
- $\text{Res}_{r_2 \theta_1} \omega(p) = 0$.

We say that ω has imaginary parameter if it has imaginary parameter at all points $p \in D[2]$.

Recall that these residues are only well defined up to sign, so that their absolute values are well defined. Although elliptic symplectic forms with imaginary parameter seem very close to being induced by generalized complex structures, and in fact locally they are, due to possible orientation issues, they might not globally correspond to a stable generalized complex structure. To see when this is the case, we need the following definition.

Definition 2.26 Let M^4 be an oriented manifold with a co-oriented elliptic divisor $(I_{|D|}, \mathfrak{o})$. Given $p \in D[2]$, let (D_1, D_2) be two local embedded submanifolds for which $D = D_1 \cup D_2$ around p . The *intersection index* of p is

$$\varepsilon_p = \begin{cases} +1, & \text{if the isomorphism } N_p D_1 \oplus N_p D_2 \simeq T_p M \text{ is orientation-preserving,} \\ -1, & \text{otherwise.} \end{cases}$$

The *parity* of a connected component D' of D is given by the product $\varepsilon_{D'} = \prod_{p \in D'[2]} \varepsilon_p$.

The parity of a connected component D' of D does not depend on the choice of co-orientation of D , and if $D'[2]$ has n points, a change of orientation of M changes the parity of D' by $(-1)^n$. We extend the definition of *parity* to a smooth connected component D' of D by declaring its parity $\varepsilon_{D'}$ to be $+1$ if D' is co-orientable, and -1 if it is not.

An elliptic symplectic form ω defines an orientation because ω^n is nonzero outside a codimension-2 subset. Using this orientation, we can determine when an elliptic symplectic form induces a stable generalized complex structure.

Corollary 2.27 [11] Let M^4 be a manifold endowed with a co-orientable elliptic divisor $I_{|D|}$, and let $\omega \in \Omega^2(\mathcal{A}_{|D|})$ be elliptic symplectic with zero elliptic residue and imaginary parameter. If the parity of all connected components of D with respect to the orientation determined by ω is 1, then there exists a co-orientation \mathfrak{o} for $I_{|D|}$ such that $(I_{|D|}, \mathfrak{o}, \omega)$ induces an equivalence class of stable generalized complex structures.

This corollary gives us the following strategy: first, construct elliptic symplectic structures with imaginary parameter, and then compute the parity of the connected components of the divisor. This is more convenient than using Theorem 2.23 directly, as it separates the construction of the symplectic structure from the existence of a particular co-orientation of the divisor.

3 Boundary maps and Lefschetz fibrations

The game we play next is to single out a class of maps that admits enough singularities to make them interesting, while also giving us enough control on the singular behavior so that we can use these maps to perform geometric constructions. The main point of this section is to extend the notion of boundary Lefschetz fibration defined in [9] for manifolds with smooth divisors to manifolds with self-crossing divisors. This extension allows for maps to have one extra type of singularity: elliptic–elliptic type. This

change allows us to get a much better grasp on many generalized complex manifolds as those can be easily described as fibrations with elliptic–elliptic singularities.

3.1 Boundary maps

Our first step is to single out a very general class of maps which is compatible with the Lie algebroids introduced in Section 2.2. These are the *boundary maps* which already illustrate how singular behavior of maps can be coupled with Lie algebroids.

We start with some basic terminology. A *pair*, (M, D) , is a manifold M together with a (possibly) immersed submanifold $D \subseteq M$. A *map of pairs* $f: (M, D) \rightarrow (N, Z)$ is a map $f: M \rightarrow N$ such that $f(D) \subseteq Z$. A *strong map of pairs* furthermore satisfies $f^{-1}(Z) = D$. Finally, (N, Z) is a *log pair* if the vanishing ideal I_Z is a log divisor ideal on N .

Definition 3.1 Let $f: (M, D) \rightarrow (N, Z)$ be a strong map of pairs onto a real log pair. Then f is a *boundary map* if $I_{|D|} := f^* I_Z$ defines an elliptic divisor ideal.

Example 3.2 The basic example to have in mind for boundary map is

$$f_1: (\mathbb{C}^2, D) \rightarrow (\mathbb{R}^2, Z), \quad f_1(z_1, z_2) = (|z_1|^2, |z_2|^2),$$

where $D \subset \mathbb{C}^2$ and $Z \subset \mathbb{R}^2$ are the two coordinate axes.

There are other examples of boundary maps that we will eventually exclude by imposing further requirements, but which are also interesting to keep in mind for now:

$$f_2: (\mathbb{C}^2, D) \rightarrow (\mathbb{R}, \{0\}), \quad f_2(z_1, z_2) = |z_1|^2 |z_2|^2,$$

where $D \subset \mathbb{C}^2$ is again the two coordinate axes and

$$f_3: (S^2, \{p_N, p_S\}) \rightarrow (S^1, \{-1\}), \quad f_3(x, y, z) = \exp(\pi i z),$$

where p_N, p_S are the north and south poles of the unit sphere and we regard S^1 as the complex numbers of length 1.

Notice that in the first two examples above, the image of the maps considered are manifolds with corners, and for all intents and purposes, we could have considered them as maps into their image with the divisor being determined by the boundary. This is in line with the original idea behind log geometry (also known as *b-geometry*) developed by Mazzeo and Melrose [25]. The third map shows that sometimes the image may be a genuine manifold (without boundary). Note that f_3 factors through the height map $\tilde{f}_3: (S^2, \{p_N, p_S\}) \rightarrow (I, \partial I)$, $\tilde{f}_3(x, y, z) = z$,

$$\begin{array}{ccc} & & (I, \partial I) \\ & \nearrow \tilde{f}_3 & \downarrow \exp(\pi i \cdot) \\ (S^2, \{p_N, p_S\}) & \xrightarrow{f_3} & (S^1, \{-1\}), \end{array}$$




Figure 2: Boundaryfication of \mathbb{R}^2 with two coordinate axes.

which has image a manifold with boundary, and boundary as divisor. This is a specific example of a more general construction, namely that we can “cut N open along Z .” Next, we will describe this procedure, which justifies the name *boundary map*.

Lemma 3.3 *Let (N, Z) be a real log pair with N a manifold without boundary. Then there is a manifold with corners \tilde{N} and a map $p: (\tilde{N}, \partial\tilde{N}) \rightarrow (N, Z)$ such that:*

- p is a map of divisors.
- $p: \tilde{N} \setminus \partial\tilde{N} \rightarrow N \setminus Z$ is a bijection.
- Every point $x \in \tilde{N}$ has a neighborhood U such that $p: U \rightarrow p(U)$ is a diffeomorphism.

Furthermore, if $p': (N', \partial N') \rightarrow (N, Z)$ is another manifold and map satisfying the properties above, then there is a unique diffeomorphism $\Psi: (N', \partial N') \rightarrow (\tilde{N}, \partial\tilde{N})$ for which $p' = p \circ \Psi$.

Proof We start with a local construction. Denoting by \mathbb{R}_ℓ^n the manifold \mathbb{R}^n with divisor given by the hyperplanes determined by the equation $x_1 \cdots x_\ell = 0$, we let $\tilde{\mathbb{R}}_\ell^n$ be given by the disjoint union

$$\tilde{\mathbb{R}}_\ell^n = \bigcup_{K \in \{-1, 1\}^\ell} \{(x_1, \dots, x_n) \in \mathbb{R}^n : k_i x_i \geq 0, \text{ where } K = (k_1, \dots, k_\ell)\},$$

and we let $p: \tilde{\mathbb{R}}_\ell^n \rightarrow \mathbb{R}_\ell^n$ be the natural inclusion: $p(x) = x$. Figure 2 shows this construction for \mathbb{R}^2 with the two coordinate axes as its real log divisor. We call each connected component of $\tilde{\mathbb{R}}_\ell^n$ defined above a *quadrant*.

Notice that $p: \tilde{\mathbb{R}}_\ell^n \rightarrow \mathbb{R}_\ell^n$ is a map of divisors, and if a smooth map $f: M \rightarrow \mathbb{R}^n$ has its image in a quadrant, then it admits a smooth lift to $\tilde{\mathbb{R}}_\ell^n$. Furthermore, if M is connected and the image of f has points which are not in the hyperplanes determined by $x_1 \cdots x_\ell = 0$, then this lift is unique.

For the global construction, we observe that charts in N provide a way to glue the local construction above to produce a manifold with corners. Indeed, given two charts that render the divisor in standard form, in their overlap, the change of coordinates gives a diffeomorphism $\Phi: \mathbb{R}_\ell^n \rightarrow \mathbb{R}_\ell^n$, for some ℓ . Since the charts are adapted to the divisor, Φ also induces a diffeomorphism of quadrants, that is, it lifts to a diffeomorphism $\tilde{\Phi}: (\tilde{\mathbb{R}}_\ell^n, \partial\tilde{\mathbb{R}}_\ell^n) \rightarrow (\tilde{\mathbb{R}}_\ell^n, \partial\tilde{\mathbb{R}}_\ell^n)$.

Since the changes of coordinates arising from an atlas for N give rise to a Čech cocycle of diffeomorphisms, the same holds for their lifts, so the procedure can be used to produce a manifold with corners \tilde{N} . Furthermore, the natural local maps, “ p ,”

defined in coordinate charts above patch together to give a global map of divisors $p: (\tilde{N}, \partial\tilde{N}) \rightarrow (N, Z)$. By construction, $p: \tilde{N} \setminus \partial\tilde{N} \rightarrow N \setminus Z$ is a bijection away from the divisors and a local diffeomorphism onto its local image.

Finally, if $p': (N', \partial N') \rightarrow (N, Z)$ is a map of divisors with the two properties above, then we show that p' has a unique lift $\Psi: (N', \partial N') \rightarrow (\tilde{N}, \partial\tilde{N})$:

$$\begin{array}{ccc}
 & & (\tilde{N}, \partial\tilde{N}) \\
 & \nearrow \Psi & \downarrow p \\
 (N', \partial N') & \xrightarrow{p'} & (N, Z)
 \end{array}$$

Indeed, in this case, $p^{-1} \circ p': N' \setminus \partial N' \rightarrow \tilde{N} \setminus \partial\tilde{N}$ is a diffeomorphism and, by the third property, any point $x \in \partial N'$ has a connected neighborhood $U \subset N'$ that maps diffeomorphically onto its image. Hence, taking U small enough, since U is connected, $p'(U)$ lies in a quadrant for a coordinate chart in N and hence p' has a unique (local) lift to \tilde{N} . Patching these local lifts together, we obtain the map Ψ . Since Ψ is a diffeomorphism in the interior of N' and by construction also a local diffeomorphism for points in the boundary of N' , it is a global diffeomorphism. ■

Definition 3.4 The *boundarification* of a manifold without boundary together with a real divisor, (N, Z) , is a manifold with corners $(\tilde{N}, \partial\tilde{N})$ together with a map $p: (\tilde{N}, \partial\tilde{N}) \rightarrow (N, Z)$ satisfying the properties of Lemma 3.3.

Example 3.5 If we take N to be the two-dimensional torus and Z to be an embedded circle which represents a primitive homology class, the boundarification of N is a cylinder and the map p identifies the two ends of the cylinder. If we take Z to be a pair of embedded circles intersecting transversely and which represent a basis for the homology of the torus, then the boundarification is a rectangle and the quotient map identifies opposite sides in the usual fashion.

Proposition 3.6 Let $f: (M, D) \rightarrow (N, Z)$ be a boundary map onto a manifold without boundary equipped with a real log divisor. Then there exists a unique boundary map $\tilde{f}: (M, D) \rightarrow (\tilde{N}, \partial\tilde{N})$ to its boundarification that is a lift of f , i.e., which satisfies $f = p \circ \tilde{f}$ for $p: (\tilde{N}, \partial\tilde{N}) \rightarrow (N, Z)$.

Proof All we need to prove is that every point $x \in M$ has a neighborhood U such that $f|_U: U \rightarrow N$ admits a unique lift $\tilde{f}|_U: U \rightarrow \tilde{N}$. Indeed, if this is the case, then any two such local lifts will agree in their overlap by uniqueness and hence the local lifts patch together to give a unique global map.

Since D has codimension 2 in M , every point $x \in M$ has a neighborhood U such that $U \setminus D$ is connected. It follows that $f(U \setminus D)$ lies in a connected component of $N \setminus Z$. By taking U small enough, we have that $f(U \setminus D)$ lies in a connected component of the complement of Z in a coordinate patch $V \subset N$, that is, $f(U \setminus D)$ lies in a quadrant and, by continuity, so does $f(U)$. As such, there is a unique lift to a map $\tilde{f}: U \rightarrow \tilde{N}$. ■

We intend to use boundary maps to construct geometric structures on their total space. Thus, we can, without loss of generality, assume that the target of a boundary

map is $(N, \partial N)$, a manifold with corners whose real log divisor is determined by its boundary. This also explains the terminology “boundary map.”

3.2 Boundary Lefschetz fibrations

The notion of a boundary map f is still too general to give us enough information about the singularities of the map. To get a good grasp on f , we need to ensure that its singularities are well controlled and this is what we do next. There are two ways to constrain the singularities of f : we can either impose that they display a specific behavior with respect to the ideals (and Lie algebroids) present, or we can impose that singularities disjoint from the vanishing loci of those ideals acquire a specific normal form. We will follow both routes here.

Note that a boundary map is by definition a map of pairs, so that it satisfies $f(D) \subseteq Z$. The first restriction we impose is that the map moreover respects the stratifications present on both D and Z .

Definition 3.7 A *fibrating boundary map* is a boundary map $f: (M, D) \rightarrow (N, Z)$ such that, for each $k \geq 1$, we have that:

- $f: (M, D[k]) \rightarrow (N, Z[k])$ is a strong map of pairs.
- Each restriction $f|_{D[k]}: D[k] \rightarrow Z[k]$ is a submersion.

In Example 3.2, f_1 and f_3 are fibrating boundary maps, whereas f_2 is not as it does not satisfy the first condition.

For a fibrating boundary map, f , we can use the ideals on M and N to control the singular behavior of f in a neighborhood of their corresponding divisors. Concretely, we have a pointwise normal form for the map.

Lemma 3.8 Let $f: (M^n, D^{n-2}) \rightarrow (N^m, Z^{m-1})$ be a fibrating boundary map, and let $x \in D[k]$. Then there exist coordinates (x_1, \dots, x_n) around x and (z_1, \dots, z_k, y_i) around $f(x)$ such that:

- Z is the standard log divisor with intersection number k on $\mathbb{R}^k \times \mathbb{R}^{m-k}$.
- D is the standard elliptic divisor with intersection number k on $\mathbb{R}^{2k} \times \mathbb{R}^{n-2k}$.
- In these coordinates, the map f takes the form

$$f(x_1, \dots, x_n) = (x_1^2 + x_2^2, \dots, x_{2k-1}^2 + x_{2k}^2, x_{n-m+k}, \dots, x_n).$$

Conversely, if for every point in D the map f is given in standard coordinates for the divisors by the expression above, then it is a fibrating boundary map.

Proof Choose a tubular neighbourhood \mathcal{V} of $Z[k]$ and denote by $\text{pr}_{Z[k]}: N Z[k] \rightarrow Z[k]$ the projection. Let $V' \subset \mathcal{V}$ be an open neighborhood of $f(x)$ on which $Z[k]$ is the standard log divisor and write $V' \cap Z[k] = \{z_1 \cdot \dots \cdot z_k = 0\}$. Choose a coordinate system (y_{k+1}, \dots, y_m) on $V' \cap Z[k]$, so that the set $\{z_1, \dots, z_k, \text{pr}_{Z[k]}^* y_{k+1}, \dots, \text{pr}_{Z[k]}^* y_m\}$ forms a coordinate system on V' which is possible because $\text{pr}_{Z[k]}$ is a submersion. Because f is a morphism of divisors, $f^*(z_1 \cdot \dots \cdot z_k)$ generates an elliptic divisor ideal on $U := f^{-1}(V') \subseteq M$. Using that f is fibrating, after possibly shrinking U around x , let (x_1, \dots, x_n) be coordinates on U in which this is the standard elliptic divisor, and such that $f^*(z_j) = x_{2j-1}^2 + x_{2j}^2$.

Because the restriction $f|_{Z[k]}$ is a submersion, we see that

$$\{x_1, \dots, x_{2k}, f^* \text{pr}_{Z[k]}^* \gamma_{k+1}, \dots, f^* \text{pr}_{Z[k]}^* \gamma_m\}$$

forms a functionally independent set. We can complete this to a coordinate system on M and relabel these as (x_1, \dots, x_n) . If we use the coordinate system $(z_1, \dots, z_k, \text{pr}_{Z[k]}^* \gamma_{k+1}, \dots, \text{pr}_{Z[k]}^* \gamma_m)$ on N and the above coordinates on M , then f takes the required form.

The converse follows immediately from the local expression for f . ■

Remark 3.9 Even if M and N are oriented manifolds and we require the use of coordinate charts compatible with orientations, we can still arrange that the local expression for f is given by the expression in Lemma 3.8. Indeed, using complex conjugation on the domain and permutation of the coordinates on both domain and codomain, we can change a coordinate chart which is not compatible with the given orientations into one that is.

In four dimensions, if $D[2]$ is nonempty, Lemma 3.8 implies that N is two-dimensional. Moreover, in a neighborhood of a point $p \in D[2]$, orientations of M and N in fact dictate which one is “the first” strand of D arriving at p and which one is “the second,” as this information is determined by the orientation of N .

Lemma 3.10 *Let $f: (M, D) \rightarrow (N, Z = \partial N)$ be a fibrating boundary map whose fibers near D are connected. Then the fibers of $f|_{D[k]}: D[k] \rightarrow Z[k]$ are connected for all $k \geq 1$.*

Proof The proof goes by induction over the strata. Note that $Z[k + 1]$ is a hypersurface in $Z[k]$, and therefore

$$f|_{D[k]}: (D[k], D[k + 1]) \rightarrow (Z[k], Z[k + 1])$$

is a fibrating boundary map for all $k \geq 0$. Applying [9, Proposition 5.25] to $f|_{M \setminus D(2)}$ tells us that the fibers of $f|_{D[1]}$ are connected. Thus, we can apply the same result to $f|_{D[1]}$ to conclude that the fibers of $f|_{D[2]}$ are connected. Continuing inductively, we arrive at the desired result. ■

The conditions imposed on the maps have, up to this point, been on behavior near D . Next, we impose the conditions away from D :

Definition 3.11 A *boundary fibration* is a fibrating boundary map $f: (M, D) \rightarrow (N, Z)$ such that $f|_{M \setminus D}: M \setminus D \rightarrow N \setminus Z$ is a surjective submersion.

Definition 3.12 A *boundary Lefschetz fibration* is a fibrating boundary map $f: (M^{2n}, D) \rightarrow (\Sigma^2, Z)$ between oriented manifolds such that $f|_{M \setminus D}: M \setminus D \rightarrow \Sigma \setminus Z$ is a Lefschetz fibration. That is, the map $f: M \rightarrow N$ is proper, $f|_{M \setminus D}$ is injective on critical points, and for each critical point $p \in M \setminus D$, there exist orientation-preserving complex coordinate charts centered at p and $f(p)$ in which f takes the form

$$f: \mathbb{C}^n \rightarrow \mathbb{C}, \quad f(z_1, \dots, z_n) = z_1^2 + \dots + z_n^2.$$

If M is four-dimensional, the definition above allows for three different types of singularities. These singularities have the following names (see also Section 6.4 for the relation with semitoric geometry).

Definition 3.13 Let $f: M^4 \rightarrow \Sigma^2$ be a smooth map.

- An *elliptic singularity* of f is a point p for which f has the local expression

$$f(x_1, x_2, x_3, x_4) = (x_1^2 + x_2^2, x_4), \quad x_i \in \mathbb{R}.$$

- An *elliptic–elliptic singularity* of f is a point p for which f has the local expression

$$f(x_1, x_2, x_3, x_4) = (x_1^2 + x_2^2, x_3^2 + x_4^2), \quad x_i \in \mathbb{R}.$$

- A *Lefschetz singularity* of f is a point p for which f has the local expression

$$f(z_1, z_2) = z_1^2 + z_2^2, \quad z_i \in \mathbb{C}.$$

The level sets associated to these singularities are, respectively, an *elliptic*, *elliptic–elliptic*, and *Lefschetz fiber*.

For a boundary Lefschetz fibration $f: (M^4, D) \rightarrow (\Sigma^2, Z)$, the first two singularities above happen at the different strata of the divisor, whereas the Lefschetz singularities do not interact with the divisor. In dimension 4, the geometry of these fibrations can be understood.

Proposition 3.14 Let $f: (M^4, D^2) \rightarrow (\Sigma^2, Z)$ be a boundary Lefschetz fibration with connected fibers, and let D' be a connected component of D . Then:

- The regular fibers of f near D are tori.
- When $D'[2]$ has $k \geq 1$ points, then D' is a union of k pairwise transversely intersecting spheres.

In particular, if $D'[2] \neq \emptyset$, then D' is co-orientable.

Proof The first point follows immediately from [9, Corollary 5.18].

For the second, assume that $D'[2]$ has at least one point. Then $f|_{D[1]}: D[1] \rightarrow Z[1]$ is a surjective submersion by assumption, which by Lemma 3.10 has connected fibers. The corresponding locus $Z'[1]$ is a disjoint union of k open intervals, and as the fibers of $f|_{D[1]}$ are connected, they must be circles. Therefore, $D'[1]$ has to be a disjoint union of cylinders. The immersed submanifold D' is obtained from $D'[1]$ by replacing the boundary circles by points and pairwise glueing these points, which implies it is as described above.

Finally, because each component of D' is an immersed sphere and thus automatically co-orientable, each component of $D'[1]$ is also co-orientable. ■

Therefore, to construct stable generalized complex structures using Corollary 2.27, the condition of co-orientability of D is satisfied as long as $D'[2]$ is nonempty for every component D' of D . For smooth components D' of D , however, i.e., when $D'[2] = \emptyset$, co-orientability is not guaranteed.

3.3 Boundary maps and Lie algebroids

Given that the ideals I_Z and $I_{|D|}$ of a boundary map determine Lie algebroids, one should expect that boundary maps (and their further specializations) are compatible with them. This is indeed the case.

Lemma 3.15 Let $f: (M, D) \rightarrow (N, Z)$ be a boundary map. Then there is a Lie algebroid morphism $(\varphi, f): \mathcal{A}_{|D|} \rightarrow \mathcal{A}_Z$ such that $\varphi \equiv df$ on sections.

Proof To prove that df induces a Lie algebroid morphism φ , by [9, Proposition 3.14], it suffices to show that f^* extends to an algebra morphism $\varphi^*: \Omega^\bullet(\mathcal{A}_Z) \rightarrow \Omega^\bullet(\mathcal{A}_{|D|})$. This can be done locally, so given $p \in D$ and $f(p) \in Z$, consider coordinates adapted to the divisors as in Examples 2.6 and 2.9:

$(X_1, Y_1, \dots, X_s, Y_s, X_{2s+1}, \dots, X_n)$ around p , $(x_1, \dots, x_j, y_{j+1}, \dots, y_m)$ around $f(p)$.

In these coordinates, we have

$$\begin{aligned} \Omega^\bullet(\mathcal{A}_{|D|}) &= \langle d \log r_1, d\theta_1, \dots, d \log r_s, d\theta_s, dX_{2s+1}, \dots, dX_n \rangle, \\ \Omega^\bullet(\mathcal{A}_Z) &= \langle d \log x_1, \dots, d \log x_j, dy_{j+1}, \dots, dy_m \rangle. \end{aligned}$$

We must verify that $f^*(d \log x_i)$ defines an elliptic form. Because f is a morphism of divisors and the ideals are locally principal, it sends generators to generators; thus, there must exist a nowhere-vanishing function g such that $f^*(x_1 \cdots x_j) = gr_1^2 \cdots r_s^2$. Consequently, by functional indivisibility of the r_i^2 , we conclude that $f^*(x_i) = hr_{i_1}^2 \cdots r_{i_\ell}^2$ for some nowhere vanishing function h and (possibly empty) subset $\{i_1, \dots, i_\ell\} \subseteq \{1, \dots, s\}$. We find that

$$f^*(d \log x_i) = d \log f^*(x_i) = d \log h + 2d \log r_{i_1} + \cdots + 2d \log r_{i_\ell},$$

which is an elliptic form as desired, so that φ is a Lie algebroid morphism.

The conditions imposed on boundary maps have a direct counterpart in Lie algebroid language. Given a Lie algebroid $\rho_{\mathcal{A}}: \mathcal{A} \rightarrow M$, let $M_{\mathcal{A}}$ be the open subset where the anchor map is an isomorphism.

Definition 3.16 [9] A Lie algebroid morphism $(\varphi, f): (\mathcal{A}, M) \rightarrow (\mathcal{A}', N)$ is said to be a:

- *Lie algebroid fibration* if the induced morphism $\varphi^1: \mathcal{A} \rightarrow f^*\mathcal{A}'$ is surjective and
- *Lie algebroid Lefschetz fibration* if $M_{\mathcal{A}}$ is dense, $f^{-1}(N_{\mathcal{A}'}) = M_{\mathcal{A}}$, and there exists a discrete set $\Delta \subset M_{\mathcal{A}}$ such that:
 - $f|_{M_{\mathcal{A}}}: M_{\mathcal{A}} \rightarrow N_{\mathcal{A}'}$ is a Lefschetz fibration with $\text{Crit}(f|_{M_{\mathcal{A}}}) = \Delta$ and
 - $(\varphi, f): (\mathcal{A}, M \setminus f^{-1}(f(\Delta))) \rightarrow (\mathcal{A}', N \setminus f(\Delta))$ is a Lie algebroid fibration.
 Note that the Lefschetz condition forces that $\text{rank}(\mathcal{A}) = 2n$ and $\text{rank}(\mathcal{A}') = 2$.

The following lemmas follow immediately from the definition, combined with Lemma 3.15.

Lemma 3.17 Let $f: (M, D) \rightarrow (N, Z)$ be a boundary fibration. Then there is a Lie algebroid fibration $(\varphi, f): (\mathcal{A}_{|D|}, M) \rightarrow (\mathcal{A}_Z, N)$ such that $\varphi \equiv df$ on sections.

Lemma 3.18 Let $f: (M^4, D^2) \rightarrow (N^2, Z^1)$ be a boundary Lefschetz fibration. Then there is a Lie algebroid Lefschetz fibration $(\varphi, f): (\mathcal{A}_{|D|}, M^4) \rightarrow (\mathcal{A}_Z, N^2)$ such that $\varphi \equiv df$ on sections.

We summarize these statements and the relationship between the different concepts in the table below:

Boundary (Lefschetz) fibration	\Rightarrow	Lie algebroid (Lefschetz) fibration
\Downarrow		\Downarrow
Fibrating boundary map	\Rightarrow	Lie algebroid map <i>submersive</i> over the singular locus
\Downarrow		\Downarrow
Boundary map	\Rightarrow	Lie algebroid map

3.4 Construction of self-crossing stable generalized complex structures

With the desired notion of boundary Lefschetz fibration in hand, we are set to prove our first result relating them to stable generalized complex structures.

From now on, we will adopt the following convention: given a boundary Lefschetz fibration $f: (M, D) \rightarrow (N, Z)$, we will orient the fibers of $f: M \setminus D \rightarrow N \setminus Z$ by declaring that the orientation of the fiber together with the orientation of the base yield the orientation of M , so that each fiber determines a homology class on $M \setminus D$. With this convention, integration over the fiber is a well-defined operation which induces the natural pairing between homology and cohomology.

Definition 3.19 A surjective boundary Lefschetz fibration, $f: (M^4, D) \rightarrow (N^2, \partial N)$, defined on a connected manifold M is *homologically essential* if the homology class $[F]$ of a (and hence any) regular fiber of $f: M \setminus D \rightarrow N \setminus Z$ is nontrivial in $H_2(M \setminus D; \mathbb{R})$ or, equivalently, if there is a class $c \in H^2(M \setminus D; \mathbb{R})$ such that $\langle c, [F] \rangle \neq 0$.

Definition 3.20 A boundary Lefschetz fibration, $f: (M^4, D) \rightarrow (N^2, Z)$, and an elliptic symplectic form $\omega \in \Omega^2(\mathcal{A}_{|D|})$ are *compatible* if $\ker \varphi \subseteq \mathcal{A}_{|D|}$ consists of symplectic vector spaces, where $\varphi: \mathcal{A}_{|D|} \rightarrow \mathcal{A}_Z$ is the induced map of Lie algebroids.

In what follows, we will have two ongoing simplifying assumptions:

- (1) We will assume that the target manifold is $(N, \partial N)$. This is not a restriction since by Proposition 3.6 we can lift f to a boundary Lefschetz fibration over the boundaryfication of (N, Z) .
- (2) We will assume that the level sets of f are connected. This also is not restriction since by [9, Proposition 5.24] we may assume that the generic fibers of f are connected and Lemma 3.10 then implies that the level sets over $Z[1]$ and $Z[2]$ are connected as well.

Before we continue, it is worth to stop and take stock of where we stand and place our quest into context. The case when the elliptic divisor is smooth was already treated in [9]. Even though there the authors only dealt with the compact case, the following is an immediate generalization for a proper map.

Theorem 3.21 [9, Theorem 7.1] *Let $(M^4, I_{|D|})$ be an oriented manifold with a smooth elliptic divisor, and let $f: (M, D) \rightarrow (N^2, Z, \omega_N)$ be a homologically essential, proper, Lefschetz fibration with connected fibers over a possibly open log-symplectic surface. Denote by $\varphi: \mathcal{A}_{|D|} \rightarrow \mathcal{A}_Z$ the induced map of Lie algebroids.*

Let $c \in H^2(M \setminus D) = H^2_{0,0}(M, \mathcal{A}_{|D|})$ be a cohomology class such that $\langle c, [F] \rangle > 0$, where F is a regular fiber of f . Then there exists a closed two-form $\eta \in \Omega^2_{0,0}(M, \mathcal{A}_{|D|})$ with $[\eta] = c$ and a positive function $\rho_0 \in C^\infty(N)$ such that:

- η is fiberwise nondegenerate, that is, for every $p \in M$, η is nondegenerate in $\ker(\varphi_p)$.
- The form $\omega = \eta + f^*(\rho\omega_\Sigma)$ is symplectic with zero elliptic residue on $\mathcal{A}_{|D|}$ for every $\rho \in C^\infty(N)$ as long as $\rho \geq \rho_0$.

Apart from the theorem above, [9] also includes a general Gompf–Thurston result for Lie algebroid Lefschetz fibrations: under similar conditions on a Lie algebroid Lefschetz fibration, one can construct a Lie algebroid symplectic form on the domain by adding a form which is symplectic on the fibers to a large multiple of the pull back of a symplectic form on the base.

Neither result can be directly applied to our case: Theorem 3.21 does not work because our divisor is not smooth, while the failure of the general result on Lie algebroid fibrations to yield stable generalized complex structures can already be seen in the simplest example.

Example 3.22 Consider the boundary fibration:

$$f_1: (\mathbb{C}^2, D) \rightarrow (\mathbb{R}^2, Z), \quad f_1(z_1, z_2) = (|z_1|^2, |z_2|^2),$$

where D and Z are the coordinate axes on \mathbb{C}^2 and \mathbb{R}^2 , respectively, as in Example 3.2.

We can endow \mathbb{R}^2 with the log-symplectic structure $d \log x_1 \wedge d \log x_2$, and consider on \mathbb{C}^2 the closed elliptic form

$$\eta = -d\theta_1 \wedge d\theta_2 + d \log r_1 \wedge d\theta_2 + d \log r_2 \wedge d\theta_1,$$

which is nondegenerate on the fibers of f_1 . The Gompf–Thurston theorem then provides us with a one-parameter family of forms

$$\omega_t = \eta + t f_1^*(d \log x_1 \wedge d \log x_2),$$

which is elliptic symplectic for $t > 1$. This poses a problem: although this defines a legitimate elliptic symplectic form, there is no value of t for which it corresponds to a stable generalized complex structure, since $|\text{Res}_{r_1 r_2} \omega_t| \neq |\text{Res}_{\theta_1 \theta_2} \omega_t|$ for $t > 1$. We conclude that the process of scaling up the log-symplectic structure on the base to achieve nondegeneracy is incompatible with the residue conditions.

What we do next is to adapt Theorem 3.21 for the self-crossing case.

Theorem 3.23 Let $f: (M^4, D^2) \rightarrow (N^2, Z = \partial N)$ be a homologically essential boundary Lefschetz fibration with connected fibers between compact connected oriented manifolds. Denote by $\varphi: \mathcal{A}_{|D|} \rightarrow \mathcal{A}_Z$ the induced map of Lie algebroids. Then $(M, I_{|D|})$ admits an elliptic symplectic structure with zero elliptic residue and imaginary parameter which is compatible with f .

If D is co-orientable and the index of each connected component of D is 1, this elliptic symplectic structure induces a stable generalized complex structure.

Proof Fix a log-symplectic structure $\omega_N \in \Omega^2(N, \mathcal{A}_Z)$. First, we consider $f: M \setminus D[2] \rightarrow N \setminus Z[2]$. This is a homologically essential, proper, boundary Lefschetz fibration with smooth elliptic divisor; hence, by Theorem 3.21, there is a form

$\eta \in \Omega_{0,0}^2(M \setminus D[2]; \log |D \setminus D[2]|)$ and a function $\rho_0 \in \Omega^0(N \setminus Z[2])$ (recall Definition 2.14) such that $\omega = \eta + f^*(\rho\omega_N)$ is a zero elliptic residue symplectic form for any function $\rho \in \Omega^0(N \setminus Z[2])$ with $\rho \geq \rho_0$.

Now, we show how to change this construction, so that the form it yields extends over $D[2]$, is elliptic symplectic with zero elliptic residue, and has imaginary parameter.

For each point $p \in D[2]$, fix open neighborhoods $U_1 \Subset U_2 \Subset U_3$ and oriented coordinates charts defined on U_3 and $f(U_3)$ in which f has the form

$$f(z_1, z_2) = (|z_1|^2, |z_2|^2).$$

As usual, we express the complex coordinates in U_3 in polar form, $z_i = r_i e^{i\theta_i}$, and denote by (x_1, x_2) the coordinates on the base, so $f^*x_i = r_i^2$.

The strategy will be to change the symplectic form ω described above in a very precise way:

- In the complement of U_3 , ω remains unchanged except for a further constant scaling of the symplectic form ω_N .
- In $U_3 \setminus U_2$, we change η into a multiple of $d\theta_2 \wedge d\theta_1$ and we preserve nondegeneracy by rescaling the symplectic form ω_N by a constant.
- In $U_2 \setminus U_1$, we interpolate the possibly large $f^*\omega_N$ to $f^*(d \log x_1 \wedge d \log x_2)$ and observe that this interpolation does not spoil the symplectic condition.
- In U_1 , we extend the symplectic form as $d\theta_2 \wedge d\theta_1 + d \log r_1 \wedge d \log r_2$, which clearly has the desired properties at p .

Now, we carry out his plan explicitly. Fix $\rho \geq \rho_0$. On U_3 , we have by Lemma 2.15 that

$$[\eta] \in H_{0,0}^2(U_3 \setminus D[2], \mathcal{A}_{|D \setminus D[2]|}) \cong H^2(U_3 \setminus D) = \mathbb{R},$$

and the generator of this cohomology pairs nonzero with the torus given in coordinates by $F = f^{-1}(r_1, r_2)$, where r_1 and r_2 are any two small positive numbers. Let $\lambda = \int_F \eta$ where integration is with respect to the fiber orientation of F , and hence $\lambda > 0$. On U_3 , consider the elliptic form $\tilde{\eta} = \frac{\lambda}{4\pi^2} d\theta_2 \wedge d\theta_1$. Then $\tilde{\eta}$ is closed in U_3 and also integrates to λ over F . Therefore, $[\eta] = [\tilde{\eta}] \in H^2(U_3 \setminus \{p\}, \mathcal{A}_{D \setminus \{p\}})$, and there is a one-form $\alpha \in \Omega^1(U_3 \setminus \{p\}, \mathcal{A}_{|D \setminus \{p\}|})$ such that $\tilde{\eta} = \eta + d\alpha$.

Let $k \geq 1$, and let ψ_1 and ψ_2 be positive functions on $f(U_3)$ such that ψ_1 is equal to 1 in neighborhood of $f(U_1)$ and has support in $f(U_2)$ and ψ_2 is equal to 1 in neighborhood of $f(U_2)$ and has support in $f(U_3)$. Then consider the form

$$\tilde{\omega} := \begin{cases} \frac{\lambda}{4\pi^2} (d\theta_2 \wedge d\theta_1 + d \log r_1 \wedge d \log r_2) & \text{in } U_1, \\ \tilde{\eta} + f^*((1 - \psi_1)k\rho\omega_N + \psi_1 \frac{\lambda}{4\pi^2} d \log x_1 \wedge d \log x_2) & \text{in } U_2 \setminus U_1, \\ \eta + d((f^*\psi_2)\alpha) + f^*(k\rho\omega_N) & \text{in } U_3 \setminus U_2, \\ \eta + f^*(k\rho\omega_N) & \text{in } M \setminus U_3. \end{cases}$$

Because of our choice of bump functions, this form is smooth. Moreover, it is clearly closed. Since $k \geq 1$, we have $k\rho \geq \rho \geq \rho_0$, and hence $\tilde{\omega}$ is symplectic in $M \setminus U_3$ for all possible values of k .

On $\overline{U_3 \setminus U_2}$, we observe that the form $\eta + d((f^* \psi)\alpha)$ is fiberwise symplectic. Indeed, its restriction to each fiber is given by

$$\eta + (f^* \psi_2)d\alpha = (f^* \psi_2)(\eta + d\alpha) + (1 - (f^* \psi_2))\eta = (f^* \psi_2)\tilde{\eta} + (1 - (f^* \psi_2))\eta;$$

hence, it is a convex combination of η and $\tilde{\eta}$, and these are both symplectic and determine the same orientation on each fiber. Since $\eta + d((f^* \psi)\alpha)$ is fiberwise symplectic and $\rho\omega_N$ is symplectic on N , the combination $\eta + d((f^* \psi)\alpha) + f^*(k\rho\omega_N)$ is symplectic on the compact set $\overline{U_3 \setminus U_2}$ as long as k is large enough.

On $U_2 \setminus U_1$, the form $\tilde{\eta}$ is given by $\frac{\lambda}{4\pi^2}d\theta_2 \wedge d\theta_1$, while the summand $f^*((1 - \psi_1)k\rho\omega_N + \psi_1 d \log x_1 \wedge d \log x_2)$ is a convex combination of two log-symplectic structures on N which determine the same orientation, that is,

$$f^*((1 - \psi_1)k\rho\omega_N + \psi_1 \frac{\lambda}{4\pi^2}d \log x_1 \wedge d \log x_2) = f^*(\kappa d \log x_1 \wedge d \log x_2),$$

for some positive function κ , and hence, on $U_2 \setminus U_1$,

$$\tilde{\omega} = \frac{\lambda}{4\pi^2}d\theta_2 \wedge d\theta_1 + (f^* \kappa)d \log r_1 \wedge d \log r_2,$$

which is clearly (zero residue) elliptic symplectic.

Finally, on U_1 , we have $\omega = \text{Im}(i \frac{\lambda}{4\pi^2}d \log z_1 \wedge d \log z_2)$, showing that it has the desired properties. ■

4 Connected sums of boundary Lefschetz fibrations

In this section, we describe a connected sum procedure for boundary Lefschetz fibrations along zero-dimensional strata of their elliptic divisors. This procedure will allow us to construct elaborate examples out of basic ones. For simplicity, we immediately restrict ourselves to dimension 4, but we note that since the connected sum takes place at points of the divisor, this procedure can also be carried out for boundary fibrations in higher dimensions.

Before we start taking connected sums of boundary Lefschetz fibrations, first recall from [11, Lemma 6.1] that we can take connected sums of elliptic divisors.

Lemma 4.1 *Let M_1^4, M_2^4 be oriented manifolds endowed with elliptic divisors $I_{|D_1|}, I_{|D_2|}$, respectively, and let $p_i \in D_i[2]$. Then $M_1 \#_{p_1, p_2} M_2$ admits an elliptic divisor $I_{|\overline{D}|}$ for which the natural inclusions $(M \setminus \{p_i\}, I_{|D_i|}) \rightarrow (M_1 \#_{p_1, p_2} M_2, I_{|\overline{D}|})$ are morphisms of divisors.*

Similarly, recall from [11, Lemma 6.1] that we can perform a self-connected sum, which when M is connected corresponds to attaching a 1-handle, and hence the diffeomorphism type of the resulting space is $M \#(S^1 \times S^3)$.

Lemma 4.2 *Let M^4 be an oriented connected manifold endowed with an elliptic divisor $I_{|D|}$, and let $p_1, p_2 \in D[2]$ be distinct points. Then $M \#(S^1 \times S^3)$ admits an elliptic divisor $I_{|\overline{D}|}$ for which the natural inclusion $(M \setminus \{p_1, p_2\}, I_{|D|}) \rightarrow (M \#(S^1 \times S^3), I_{|\overline{D}|})$ is a morphism of divisors.*

In this connected sum procedure, the map $\Phi(z_1, z_2) = \frac{1}{|z_1|^2 + |z_2|^2}(z_2, \bar{z}_1)$ is used to identify annuli. Here, (z_1, z_2) are local complex coordinates compatible with the

orientation on the manifolds. There is some freedom in the constructions above. Given a choice of local coordinates (z_1, z_2) around p_1 and p_2 , we can compose the map Φ by a permutation of the coordinates. This does not change the topology of $M_1 \#_{p_1, p_2} M_2$, but it could change the topology of the zero locus of the divisor. In dimension 4, because of this freedom in the ordering, there are potentially two different topological types of the zero locus of the divisor. Notice, however, that our notation does not reflect this ambiguity.

Remark 4.3 (Connected components) Although there is some freedom in the choices, we can still distinguish the number of connected components on the divisor on the connected sum:

- (1) When p_1 and p_2 lie in different connected components of the divisor, be that either in the connected sum of two manifolds or in a self-connected sum, the connected components containing p_1 and p_2 will combine into a single connected component of \tilde{D} .
- (2) When p_1 and p_2 are in the same connected component, D , a case that can only happen in a self-connected sum, the resulting divisor, $\tilde{D} \subset M \# (S^1 \times S^3)$, may have one or two connected components originating from D .

Next, we show that the connected sum operation is also compatible with boundary (Lefschetz) fibrations. To describe how the connected sum procedure interacts with the base of the fibration, we first consider what happens in the local model.

Lemma 4.4 Let $\Delta_r \subseteq \mathbb{R}^2$ be the triangle bounded by the axes and the line $x + y = r$, and let (x, y) be oriented coordinates on Δ_r and (z_1, z_2) be complex coordinates on \mathbb{D}_r^4 , the disk of radius r . Consider the following maps:

- $p: (\mathbb{D}_2^4 \setminus \mathbb{D}_{1/2}^4) \rightarrow (\Delta_2 \setminus \Delta_{1/2})$, given by $(z_1, z_2) \mapsto (|z_1|^2, |z_2|^2)$.
- $\Phi: (\mathbb{D}_2^4 \setminus \mathbb{D}_{1/2}^4) \rightarrow (\mathbb{D}_2^4 \setminus \mathbb{D}_{1/2}^4)$, given by $(z_1, z_2) \mapsto \frac{1}{|z_1|^2 + |z_2|^2} (z_2, \bar{z}_1)$.
- $\Psi: (\Delta_2 \setminus \Delta_{1/2}) \rightarrow (\Delta_2 \setminus \Delta_{1/2})$ given by $(x, y) \mapsto \frac{(y, x)}{(x+y)^2}$.

Then the following diagram commutes:

$$\begin{array}{ccc}
 (\mathbb{D}_2^4 \setminus \mathbb{D}_{1/2}^4) & \xrightarrow{\Phi} & (\mathbb{D}_2^4 \setminus \mathbb{D}_{1/2}^4) \\
 \downarrow p & & \downarrow p \\
 (\Delta_2 \setminus \Delta_{1/2}) & \xrightarrow{\Psi} & (\Delta_2 \setminus \Delta_{1/2}).
 \end{array}$$

The proof of this lemma is a simple verification. Just as we used the map Φ to perform a connected sum compatible with elliptic divisors, we want to use the map Ψ to define a sort of connected sum operation of the base.

Definition 4.5 Let Σ_1, Σ_2 be oriented surfaces with corners, and let q_1, q_2 be corners of Σ_1, Σ_2 , respectively. The *oriented corner connected sum* of Σ_1 and Σ_2 is defined by identifying a trapezoid neighborhood of q_1 to a trapezoid neighborhood of q_2 via Ψ . The oriented corner connected sum is an oriented surface with corners denoted by $\Sigma_1 \#_{q_1, q_2} \Sigma_2$ (see Figure 3).




Figure 3: Local oriented and orientation-reversing corner connected sums.

The oriented corner connected sum is naturally oriented and does not depend on the neighborhoods chosen. Together with the local normal form for fibrating boundary maps, we can now prove the following.

Theorem 4.6 *Let $f_i: (M_i^4, D_i) \rightarrow (N_i^2, \partial N_i)$ be boundary (Lefschetz) fibrations with connected fibers between oriented manifolds for $i = 1, 2$, let $p_i \in D_i[2]$ and $q_i = f(p_i)$. Then there exists a boundary (Lefschetz) fibration on one of the two possible connected sums $M_1 \#_{p_1, p_2} M_2$ whose base is the oriented corner sum $N_1 \#_{q_1, q_2} N_2$:*

$$(f_1 \# f_2): (M_1 \#_{p_1, p_2} M_2, \tilde{D}) \rightarrow (N_1 \#_{f(p_1), f(p_2)} N_2, \partial N_1 \#_{q_1, q_2} \partial N_2),$$

which is compatible with the (orientation-preserving) inclusions $M_i \setminus \{p_i\} \hookrightarrow M_1 \# M_2$.

Furthermore, let D'_1, D'_2 denote the connected components of the zero locus of the divisor D containing p_1, p_2 , respectively. Then the parities satisfy:

$$\varepsilon_{\tilde{D}} = -\varepsilon_{D'_1} \varepsilon_{D'_2}.$$

Finally, $f_1 \# f_2$ is homologically essential if and only if f_1 and f_2 are.

Proof By Lemma 3.8, there exist neighborhoods U_1, U_2 of $f(p_1), f(p_2)$, respectively, which provide coordinates as in the setting of Lemma 4.4. We perform the connected sum procedure using the maps described there. Because these maps are compatible with the fibrations on M_1 and M_2 , we conclude that $M_1 \#_{p_1, p_2} M_2$ admits a boundary fibration. The computation of the parity is given in [11, Theorem 6.7]. ■

Recall that there are a priori two possible topological types for the elliptic divisor, depending on the ordering of the local coordinates. However, when we are presented with fibrations between oriented manifolds $f: (M_i, D_i) \rightarrow (N_i, \partial N_i)$, the orientation

on the base determines an order for the strands of D for every point $p_i \in D_i[2]$ (cf. Remark 3.9), and the gluing of fibrations which is compatible with orientations on M_i and N_i is the one that flips the first and second strands arriving the points where the sum is performed. In particular, from the possible divisors discussed in Remark 4.3(2), only the one with two connected components occurs.

Remark 4.7 (Nonorientable case) If we were to allow the map Ψ used in the corner sum to be orientation-reversing, we would still be able to define a corner connected sum and obtain a boundary fibration. When taking the connected sum of two manifolds, this does not cause a qualitative change in the outcome. However, if we use the orientation-reversing corner sum on the base for a self-connected sum, we see that the resulting base manifold is not orientable as a Möbius band appears.

Now that we understand precisely what happens to the connected components of the divisor on the self-connected sum, we can state the following.

Corollary 4.8 Let $f: (M^4, D^2) \rightarrow (N^2, \partial N)$ be a boundary (Lefschetz) fibration with connected fibers between oriented manifolds, and let $p_1, p_2 \in D[2]$ be distinct. Then $M\#(S^1 \times S^3)$ admits a boundary (Lefschetz) fibration \tilde{f} which is compatible with the inclusion $M \setminus \{p_1, p_2\} \hookrightarrow M\#_{p_1, p_2}(S^1 \times S^3)$, and for which $\tilde{D}[2] = D[2] \setminus \{p_1, p_2\}$.

Moreover, let D'_{p_1}, D'_{p_2} denote the connected components of D containing p_1, p_2 , respectively.

- If $p_i \in D'_{p_i}[2]$ and $D'_{p_1} \neq D'_{p_2}$, then the corresponding connected component \tilde{D}' of \tilde{D} satisfies:

$$\varepsilon_{\tilde{D}'} = -\varepsilon_{D'_1} \varepsilon_{D'_2}.$$

- If $p_i \in D'_{p_i}[2]$ and $D'_{p_1} = D'_{p_2}$, then the corresponding connected components $\tilde{D}'_1, \tilde{D}'_2$ of \tilde{D} satisfy:

$$\varepsilon_{\tilde{D}'_1} \varepsilon_{\tilde{D}'_2} = -\varepsilon_{D'_{p_1}} \varepsilon_{D'_{p_2}}.$$

Finally, \tilde{f} is homologically essential if and only if f is.

5 Singularity trades

The goal of this section is to prove two theorems which allow one to trade Lefschetz for elliptic–elliptic singularities and vice versa. To formulate these results, we need to recall the notion of vanishing cycle for both Lefschetz and elliptic singularities.

Given a boundary Lefschetz fibration $f: (M^4, D^2) \rightarrow (N^2, \partial N)$ and an elliptic or a Lefschetz singularity $p_1 \in M$, let $q_1 = f(p_1)$ be the corresponding singular value. We fix $q \in N$, a reference regular point of f and $\gamma: [0, 1] \rightarrow N$, a simple path connecting q to q_1 which goes through no critical values of f except for q_1 at time 1. We can consider $F_q = f^{-1}(q)$, $F_\gamma = f^{-1}(\gamma([0, 1]))$ and the natural inclusion $\iota: F_q \rightarrow F_\gamma$. Then F_q is a two-torus and $H_1(F_\gamma)$ is one-dimensional:

- In the case of a Lefschetz singularity, the inclusion $H_1(F_q) \rightarrow H_1(F_\gamma)$ has kernel given by the Lefschetz vanishing cycle, which corresponds to the boundary of a Lefschetz thimble emanating from the singularity.

- In the case of an elliptic singularity, F_γ is the product of circle and a solid torus with F_q as boundary, and hence $\iota_*: H_1(F_q) \rightarrow H_1(F_\gamma)$ also has one-dimensional kernel given by the cycle in F_q which becomes a boundary in F_γ .

In both cases, the kernel of ι_* is generated by one primitive element in $H_1(F_q; \mathbb{Z})$ which depends only on the homotopy class of γ in $N \setminus \text{Crif}f$.

Definition 5.1 In the situation above, the *vanishing cycle* associated to the singular value q_1 and the homotopy class of the path γ is either of the primitive elements in $H_1(F_q; \mathbb{Z})$ which generates the kernel of $H_1(F_q; \mathbb{Z}) \rightarrow H_1(F_\gamma; \mathbb{Z})$.

Definition 5.2 Let $f: (M^4, D^2) \rightarrow (N^2, \partial N)$ be a boundary Lefschetz fibration, and let F_{q_0} and F_{q_1} be Lefschetz or elliptic fibers. We say that the vanishing cycles at F_{q_0} and F_{q_1} are a *dual pair* if there is a simple path $\gamma: [0, 1] \rightarrow N$ such that:

- $\gamma(0) = q_0$ and $\gamma(1) = q_1$.
- $\gamma((0, 1))$ only contains regular values of f .
- The vanishing cycles on both ends of γ together generate the integral homology of the regular torus fiber, say $F_{\gamma(1/2)}$.

With these notions at hand, we can give the precise statements of our singularity trade theorems.

Theorem 5.3 (Elliptic–elliptic trade) *Let $f: (M^4, D) \rightarrow (N^2, \partial N)$ be a boundary Lefschetz fibration with connected fibers, and let $p \in D[2]$. Then M admits a boundary Lefschetz fibration $\tilde{f}: (M^4, \tilde{D}) \rightarrow (\tilde{N}^2, \partial \tilde{N})$ such that:*

- \tilde{N} is obtained from N by smoothing out the corner $f(p)$.
- \tilde{f} and \tilde{D} agree with f and D outside a small ball centered at p .
- $\tilde{D}[2] = D[2] \setminus \{p\}$, i.e., \tilde{f} has one elliptic–elliptic singularity less than f .
- \tilde{D} and D have the same parity.
- \tilde{f} has one Lefschetz singularity more than f .
- \tilde{f} has an elliptic singularity whose vanishing cycle forms a dual pair with the new Lefschetz vanishing cycle.
- \tilde{f} is homologically essential if and only if f is.

By induction, any manifold which admits a boundary Lefschetz fibration admits one with a smooth embedded divisor.

The converse trade is given by the next theorem.

Theorem 5.4 (Lefschetz trade) *Let $\tilde{f}: (\tilde{M}^4, \tilde{D}) \rightarrow (\tilde{N}^2, \partial \tilde{N})$ be a boundary Lefschetz fibration with connected fibers, and assume that the vanishing cycles at a Lefschetz fiber, F_{q_0} , and at an elliptic fiber, F_{q_1} , form a dual pair. Then there is a boundary Lefschetz fibration, $f: (M^4, D) \rightarrow (N^2, \partial N)$, such that:*

- N is obtained from \tilde{N} by adding a corner at q_1 .
- f and D agree with \tilde{f} and \tilde{D} outside $\tilde{f}^{-1}(V_2)$, where V_2 is a neighborhood of the path that expresses a vanishing cycles as a dual pair.
- $D[2] = \tilde{D}[2] \cup \{p\}$ and hence f has one elliptic–elliptic singularity more than \tilde{f} .
- D and \tilde{D} have the same parity.
- f has one Lefschetz singularity less than \tilde{f} .




Figure 4: The base of the boundary Lefschetz fibration on S^4 together with a path expressing the Lefschetz and elliptic singularities as a dual pair.

- f is homologically essential if and only if \tilde{f} is.

The proofs of these theorems rely on the existence of specific boundary Lefschetz fibrations on S^4 and on the open disk \mathbb{D}^4 .

Lemma 5.5 *There exists a homologically essential boundary Lefschetz fibration with connected fibers, $f_{S^4}: (S^4, D^2) \rightarrow (N, \partial N)$, with the following properties (see Figure 4):*

- $D[2]$ has only one point, which has index -1 .
- N is the disk with one corner.
- f_{S^4} has only one Lefschetz singularity.
- The vanishing cycles of the Lefschetz fiber and any elliptic fiber form a dual pair.

The proof of this lemma is somewhat long, so we will postpone it to this end of this section.

Lemma 5.6 *Let (\mathbb{D}^4, D) be the open disk in \mathbb{C}^2 with divisor I_{z_1} , and let $\mathbb{D}_+^2 \subset \mathbb{R}^2$ be the open half-disk with boundary in the real axis*

$$\mathbb{D}_+^2 = \{(x, y) \in \mathbb{R}^2: x^2 + y^2 < 1 \text{ and } x \geq 0\}.$$

Then there is a proper boundary Lefschetz fibration with connected fibers, $f_{\mathbb{D}^4}: (\mathbb{D}^4, D) \rightarrow (\mathbb{D}_+^2, \partial\mathbb{D}_+^2)$, such that:

- $f_{\mathbb{D}^4}$ has a single Lefschetz fiber.
- The vanishing cycles of the Lefschetz fiber and the elliptic fiber form a dual pair.

Furthermore, if $f: (M, D) \rightarrow (\mathbb{D}_+^2, \partial\mathbb{D}_+^2)$ is a proper boundary Lefschetz fibration with connected fibers with the two properties above, then f is equivalent to $f_{\mathbb{D}^4}$, that is, there is a commutative diagram:

$$\begin{array}{ccc} \mathbb{D}^4 & \longrightarrow & M \\ \downarrow f_{\mathbb{D}^4} & & \downarrow f \\ (\mathbb{D}_+^2, \partial\mathbb{D}_+^2) & \longrightarrow & (\mathbb{D}_+^2, \partial\mathbb{D}_+^2), \end{array}$$




Figure 5: The base of the boundary Lefschetz fibration on S^4 split in two halves, each half being a fibration of \mathbb{D}^4 .

where the horizontal maps are diffeomorphisms.

Proof The existence of the fibration $f_{\mathbb{D}^4}$ follows from Lemma 5.5. Indeed, we split the base of f_{S^4} in two parts, V_1 , a neighborhood of the vertex, and V_2 , the rest of the base plus a small overlap with V_1 , as indicated in Figure 5. Then, due to Lemma 3.8, on $f^{-1}(V_1)$, in appropriate coordinates, we have

$$V_1 = \{(x, y) \in \mathbb{R}^2: x + y < 1, x \geq 0, y \geq 0\},$$

and the fibration is given by

$$f_{S^4}(z_1, z_2) = (|z_1|^2, |z_2|^2).$$

Hence, $f_{S^4}^{-1}(V_1)$ is a disk and its complement $f_{S^4}^{-1}(V_2)$ is also a disk. However,

$$f_{S^4}|_{V_2}: V_2 \rightarrow f_{S^4}(V_2)$$

has all the properties required in the lemma after we choose a diffeomorphism between V_2 and \mathbb{D}_+^2 . Therefore, we have existence.

To prove the uniqueness part, we study all possible ways such a fibration may arise. Let $f: M \rightarrow (\mathbb{D}_+^2, \partial\mathbb{D}_+^2)$ be a boundary Lefschetz fibration satisfying the assumptions of the lemma. Without loss of generality, we assume that the image of the Lefschetz singularity is $(2/3, 0)$, and we split \mathbb{D}_+^2 in two parts:

$$U_1 = \{(x, y) \in \mathbb{D}_+^2: x \leq 1/2\},$$

$$U_2 = \{(x, y) \in \mathbb{D}_+^2: x \geq 1/2\}.$$

The set $f^{-1}(U_2)$ is a neighborhood of the Lefschetz fiber, and hence its differentiable type as a fibration is fully determined [17]. Similarly, the set $f^{-1}(U_1)$ is a neighborhood of an elliptic fiber, and hence its differentiable type as a fibration is also fully determined:

$$f^{-1}(U_1) = \mathbb{D}^2 \times S^1 \times (-1, 1), \quad f(re^{i\theta}, \psi, t) = (r^2, t).$$

Therefore, all possible different fibrations with the desired properties are determined by the different ways these two pieces can be glued together modulo the action of the isomorphism group of each half of the fibration.

Since the gluing takes place over a regular fibration over an interval, the isotopy class of the gluing map is determined by the isotopy class of the map it induces at a single fiber. Since the fibers are tori, this is in turn determined by the corresponding map in homology. Since the vanishing cycles form a dual pair, there is, modulo the action of the isomorphism group of the fibration over V_1 , a unique way to glue these together. ■

Next, we show how to use Lemmas 5.5 and 5.6 to prove both singularity trade theorems.

Proof Applying Theorem 4.6 to the boundary Lefschetz fibration on M and on S^4 gives rise to a boundary Lefschetz fibration on $M\#S^4 \simeq M$, for which the inclusion $M \setminus \{p\} \hookrightarrow M$ preserves fibrations, in particular, we see that the new fibration on M only changes in the small ball around p used for the connected sum procedure. Since the divisor in S^4 has only one point in the top stratum, the new divisor satisfies $\tilde{D}[2] = D[2] \setminus \{p\}$, and \tilde{D} and D have the same index. Given the way the fibrations are glued, we see that the effect on the base is to smooth out the corner corresponding to $f(p)$. Continuing inductively gives rise to a boundary Lefschetz fibration with embedded divisor. ■

Proof Under the conditions of the theorem, γ has a neighborhood, V_2 , diffeomorphic to \mathbb{D}_+^2 in which the fibration has only one Lefschetz singularity whose vanishing cycle forms a dual pair with the elliptic singularity. Hence, by Lemma 5.6, $f^{-1}(V_2)$ is diffeomorphic to \mathbb{D}^4 , and f is equivalent to the fibration of Lemma 5.6. Since the fibration on S^4 splits as two disks, one fibering over \mathbb{D}_+^2 and the other fibering over a neighborhood, V_1 , of the origin in $(\mathbb{R}_+)^2$ (see Figure 5), we can realize $M\#S^4$ as follows: remove the disk $f^{-1}(V_2)$ and glue back, by the natural identification of the boundary, $f_{S^4}^{-1}(V_1)$.

Since this procedure corresponds to performing connected sum with S^4 , the final manifold is still diffeomorphic to M and the fibration only changes in the part that has been surged in, which includes the removal of the Lefschetz singularity from $f^{-1}(V_2)$ and the inclusion of the elliptic–elliptic singularity of $f_{S^4}^{-1}(V_1)$. Finally, notice that the process of filling the boundary of $f^{-1}(V_2)$ with $f_{S^4}^{-1}(V_1)$ is not compatible with the given orientations of these spaces, since they both appear at opposite sides of a boundary in S^4 . That is, the orientation of M is compatible with the opposite orientation of $f_{S^4}^{-1}(V_1)$. Since the elliptic–elliptic singularity for the fibration in S^4 had index -1 and the orientation of S^4 was reversed in the connected sum process, the intersection index of the new elliptic–elliptic singularity on M is $+1$, and hence the overall parity of the divisor is unchanged. ■

To finish the proof of the trade theorems, we must establish Lemma 5.5, which we do next.

Proof The proof is done in two steps. In the first step, we show that if M is the total space of a boundary Lefschetz fibration whose singularities are as stated in Lemma 5.5, then $M = S^4$. In the second step, we show that such a fibration exists.

Step 1. We observe once again that M is made of two fibrations glued together, as illustrated in Figure 5: one fibration with an elliptic–elliptic singularity over V_1 and one with a Lefschetz singularity over V_2 . The fibration over V_1 is a copy of \mathbb{D}^4 added




Figure 6: Kirby diagram for the total space of the fibration described in Lemma 5.6.

along its S^3 boundary, that is, $M = f^{-1}(V_2) \cup 4\text{-handle}$. The space $f^{-1}(V_2)$ itself can be readily described as a handlebody: we start with a neighborhood of a regular fiber, then add a -1 -framed 2-handle along the vanishing cycle of the Lefschetz singularity to obtain a neighborhood of the Lefschetz singular fiber and a 0 -framed 2-handle along the vanishing cycle of the elliptic singularity. Therefore, the Kirby diagram of M is the one depicted in Figure 6a. We can then slide the 2-handle that goes around both 1-handles to obtain Figure 6b and see that the resulting 2-handle separates as a 0 -framed 2-handle from the rest of the diagram and hence cancels with the 3-handle. The remaining pairs of 1- and 2-handles clearly cancel each other (Figure 6c), leaving us with the empty diagram, which corresponds to S^4 .

Step 2. To construct the desired fibration, we will use a plumbing construction applied to the disk bundle of $\mathcal{O}(2) \rightarrow \mathbb{C}P^1$ in a way that is compatible with the natural torus fibration of that space. Throughout, we will use fixed parameterizations $\varphi_1, \varphi_2: \mathbb{C}^2 \rightarrow \mathcal{O}(2)$ for which the change of coordinates is given by

$$\varphi_2^{-1} \circ \varphi_1(z, w) = (z^{-1}, z^{-2}w).$$

We will refer to φ_1 as parameterizing a trivialization of $\mathcal{O}(2)$ with the south pole removed and similarly φ_2 does not cover the fiber over the north pole.

Rotation on both coordinates in the parameterization φ_2 give rise to a torus action on $\mathcal{O}(2)$ which, in the parameterizations above, is given by

$$(5.1) \quad \begin{aligned} (e^{i\theta_1}, e^{i\theta_2}) \cdot \varphi_1(z, w) &= \varphi_1(e^{-i\theta_1}z, e^{i(-2\theta_1+\theta_2)}w), \\ (e^{i\theta_1}, e^{i\theta_2}) \cdot \varphi_2(z, w) &= \varphi_2(e^{i\theta_1}z, e^{i\theta_2}w). \end{aligned}$$

To describe the quotient of $\mathcal{O}(2)$ by this torus action, we will also want to consider $[-1, 1] \times \mathbb{R}_+$. Of course, this space can be parameterized by a single, rather obvious, chart, but it will be convenient to parameterize it by two charts instead. We consider the parameterizations

$$\psi_1: \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow (-1, 1] \times \mathbb{R}_+ \subset [-1, 1] \times \mathbb{R}_+ \quad \psi_1(x_1, y_1) = \left(\frac{1 - x_1}{1 + x_1}, \frac{y_1}{(1 + x_1)^2} \right),$$

$$\psi_2: \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow [-1, 1) \times \mathbb{R}_+ \subset [-1, 1] \times \mathbb{R}_+ \quad \psi_2(x_2, y_2) = \left(-\frac{1 - x_2}{1 + x_2}, \frac{y_2}{(1 + x_2)^2} \right),$$

and keep in mind that these parameterizations induce opposite orientations, with ψ_2 agreeing with the natural orientation of $[-1, 1] \times \mathbb{R}_+$.

Lemma 5.7 *If we let $h: S^2 \rightarrow \mathbb{R}$ be the height function and $g: \text{Sym}^2 \mathcal{O}(2) \rightarrow \mathbb{R}$ be the Fubini–Study metric, then*

$$\begin{aligned} f: \mathcal{O}(2) &\rightarrow [-1, 1] \times \mathbb{R}_+ \\ (z, w) &\mapsto (h(z), g_z(w, w)) \end{aligned}$$

defines a quotient map for the torus action on $\mathcal{O}(2)$. Furthermore, f is a proper boundary fibration with elliptic divisor induced by the holomorphic log divisor consisting of the zero section and fibers over the north and south poles. ■

Proof In the parameterizations φ_i , the height and distance functions take the form:

$$\begin{aligned} h \circ \varphi_1(z, w) &= \frac{1 - |z|^2}{1 + |z|^2}, & g \circ \varphi_1(z, w) &= \frac{|w|^2}{(1 + |z|^2)^2}, \\ h \circ \varphi_2(z, w) &= -\frac{1 - |z|^2}{1 + |z|^2}, & g \circ \varphi_2(z, w) &= \frac{|w|^2}{(1 + |z|^2)^2}, \end{aligned}$$

which are clearly invariant under the T^2 -action in equation (5.1). Furthermore, for $i = 1, 2$, the image of $f \circ \varphi_i$ lands in the image of the parameterization ψ_i , and we can compute the expression for f in these parameterizations:

$$(5.2) \quad f_i(z, w) := \psi_i^{-1} \circ f \circ \varphi_i(z, w) = (|z|^2, |w|^2),$$

which shows clearly that f is not only the quotient map, but also a boundary fibration. ■

Now, we perform a plumbing on $\mathcal{O}(2)$.

Definition 5.8 Let $\pi: M^{2n} \rightarrow N^n$ be a \mathbb{D}^n -bundle, and let $\mathbb{D}_1, \mathbb{D}_2$ be disjoint disks in N over which π is trivializable. A *self-plumbing* of π at \mathbb{D}_1 and \mathbb{D}_2 is obtained by identifying $\pi^{-1}(\mathbb{D}_1) \simeq \mathbb{D}_1 \times \mathbb{D}^n$ and $\pi^{-1}(\mathbb{D}_2) \simeq \mathbb{D}_2 \times \mathbb{D}^n$ using a map which preserves the product structure but reverses the factors.

For the case at hand, let $\mathbb{D}^2 \mathcal{O}(2)$ be the open ε -disk bundle with respect to the Fubini–Study metric. By restricting f to $\mathbb{D}^2 \mathcal{O}(2)$, we obtain a proper boundary fibration $f: \mathbb{D}^2 \mathcal{O}(2) \rightarrow [-1, 1] \times [0, \varepsilon)$.

Furthermore, we observe that φ_1 and φ_2 provide trivializations of $\mathbb{D}^2\mathcal{O}(2)$, and hence we can use them to perform a self-plumbing of $\mathbb{D}^2\mathcal{O}(2)$ at the north and south poles. Let M be defined as the self-plumbing of $\mathbb{D}^2\mathcal{O}(2)$ via the trivializations φ_i and the map

$$\Phi: \mathbb{C}^2 \rightarrow \mathbb{C}^2 : (z, w) \mapsto (\bar{w}, \bar{z}),$$

that is, $\varphi_1(z, w)$ is identified with $\varphi_2(\bar{w}, \bar{z})$.

Since the map used for the plumbing preserves elliptic ideals and identifies the north and south poles, M is endowed with an elliptic divisor with a single point in $D[2]$. Since the map Φ does not match co-orientations, the elliptic divisor in M has intersection index -1 .

To endow M with a boundary fibration, we only need to take a quotient of the base, $[-1, 1] \times [0, \varepsilon)$, by the equivalence relation that makes the following diagram commute:

$$\begin{array}{ccc} \mathbb{D}^2\mathcal{O}(2) & \xrightarrow{\sim\Phi} & \mathbb{D}^2\mathcal{O}(2) \\ \downarrow f & & \downarrow f \\ [-1, 1] \times [0, \varepsilon) & \xrightarrow{\sim\Psi} & [-1, 1] \times [0, \varepsilon). \end{array}$$

Since f is surjective, there is a unique identification, \sim_Ψ , that gives rise to such a diagram. In fact, we can easily compute it in the parameterizations ψ_i , where it is induced by the map $\Psi(x, y) = (1 - y, x + 1)$. That is, the point $\psi_1(x, y)$ is identified with the point $\psi_2(y, x)$. Since ψ_1 and ψ_2 induce opposite orientations, this identification preserves the natural orientation of $[-1, 1] \times [0, \varepsilon)$ and the quotient is an oriented half-open cylinder with one corner (see Figure 7).

Lemma 5.9 *The map $f: \mathbb{D}^2\mathcal{O}(2) \rightarrow [-1, 1] \times [0, \varepsilon)$ descends to a boundary fibration $\hat{f}: M \rightarrow N$.*

Next, we compute its monodromy along a generator of $\pi_1(N)$.

Lemma 5.10 *Let $\hat{f}: M \rightarrow N$ be the boundary fibration from Lemma 5.9. Then the monodromy of \hat{f} around a loop around the hole is a positive Dehn twist.*

Proof This is a direct computation using the given change of coordinates and the plumbing map Φ . Indeed, all we need to do is to track what happens with the torus action as we move along from the chart covered by φ_2 to the chart covered by φ_1 and then back to φ_2 via Ψ :

$$\begin{aligned} (e^{i\theta_1}, e^{i\theta_2}) \cdot \varphi_2(z, w) &= \varphi_2(e^{i\theta_1}z, e^{i\theta_2}w) = \varphi_1(e^{-i\theta_1}z^{-1}, e^{i(-2\theta_1+\theta_2)}z^{-2}w) \\ &\sim_\Phi \varphi_2(e^{i(2\theta_1-\theta_2)}\bar{z}^{-2}\bar{w}, e^{i\theta_1}\bar{z}^{-1}) = (e^{i(2\theta_1-\theta_2)}, e^{i\theta_1})\varphi_2(\bar{z}^{-2}\bar{w}, \bar{z}^{-1}). \end{aligned}$$

Therefore, we see that, in the basis $\{e_{\theta_1}, e_{\theta_2}\}$ for $H^1(F)$ corresponding to the generators of the action, the monodromy transformation is given by the matrix

$$\begin{pmatrix} 2 & -1 \\ 1 & 0 \end{pmatrix}.$$




Figure 7: The base of the boundary fibration constructed in Lemma 5.9.

Notice that using the complex orientation of $\mathcal{O}(2)$ and the standard orientation of \mathbb{R}^2 , $\{e_{\theta_1}, e_{\theta_2}\}$ is a negative basis for the homology of the fiber. Using this, we see that the transformation above is a positive Dehn twist on the cycle $e_{\theta_1} + e_{\theta_2}$. ■

Now, we can complete M to a closed manifold by glueing a neighborhood of a single Lefschetz fiber with vanishing cycle $e_{\theta_1} + e_{\theta_2}$ in the hole of the annulus. Finally, we observe that this vanishing cycle forms a dual pair with either of the two vanishing cycles of the elliptic singularity, which in the parameterization φ_2 are given by either of the cycles e_{θ_1} or e_{θ_2} .

Remark 5.11 Simply drawing a base diagram for a boundary Lefschetz fibration does not guarantee the existence of a fibration that realizes it. For example, there is no manifold whose base diagram is that of Figure 4, but for which the elliptic–elliptic singularity has intersection index 1. In the construction above, this would manifest itself in the fact that, without using complex conjugation, the monodromy of the plumbing would be a negative Dehn twist. This highlights that the long second step in the proof above is indeed necessary.

6 Examples

In this section, we give several concrete examples of boundary fibrations. We will first show that they arise naturally as the quotient maps of effective torus actions, and that our framework fits particularly well with the theory of integrable systems. This connection provides us immediately with a wealth of examples of both boundary fibrations and stable generalized complex structures. We will further illustrate our constructions by showing how starting with simple examples (of manifolds with torus actions) we can use the connected sum procedure to obtain many more examples of boundary fibrations.

6.1 Torus actions

We show that quotient maps of torus actions provide boundary fibrations.

Proposition 6.1 *Let T^n act effectively on a smooth manifold M^{2n} , with connected isotropy groups. Then:*

- $N := M^{2n}/T^n$ is a manifold with corners.
- The quotient map defines a boundary fibration $f: (M, D) \rightarrow (N, \partial N)$ with connected fibers.
- The intersection stratification of the elliptic ideal coincides with the stratification by orbit types on M .
- $ND[1]$ is co-orientable.
- If M is oriented, then so is N .
- If M is four-dimensional and the action is not free, f is homologically essential.

Proof Let $p \in M$, let G_p denote the isotropy group of p , and let \mathcal{O}_p denote the orbit of p . By assumption, G_p is connected and therefore isomorphic to T^ℓ for some $\ell \leq n$. By the slice theorem, there exists a neighborhood of \mathcal{O}_p which is equivariantly diffeomorphic to a neighborhood of the zero section in

$$G \times_{G_p} N_p \mathcal{O}_p,$$

where G_p acts linearly on $N_p \mathcal{O}_p$ by the differentiated action. Because all groups in consideration are abelian and connected, this implies that there is a neighborhood U around p of the form

$$U = T^{n-\ell} \times (\mathbb{R}^{n-\ell} \times \mathbb{C}^\ell).$$

The $T^n = (T^{n-\ell} \times T^\ell)$ -action of U decomposes as $T^{n-\ell}$ acting by multiplication on $T^{n-\ell}$ and T^ℓ acting linearly on \mathbb{C}^ℓ . Since the irreducible representations of T^ℓ are one-dimensional, we may without loss of generality assume that each coordinate line in \mathbb{C}^ℓ is preserved by the action. Therefore, if we let \mathfrak{t} denote the Lie algebra of T^ℓ , let \mathfrak{l} denote the kernel of $\exp: \mathfrak{t} \rightarrow T^\ell$, with minimal generating set $\{\xi_1, \dots, \xi_\ell\}$ and choose $\{\alpha_1, \dots, \alpha_\ell\} \in \mathfrak{l}^*$ the dual basis for the dual lattice, then the action on each irreducible representation has the form

$$\exp(\Theta) \cdot z_j = e^{2\pi i \langle \Theta, n_j \alpha_j \rangle} z_j, \quad \Theta \in \mathfrak{t}.$$

Since the action is effective, we have that $n_j \neq 0$, and because the isotropy groups are connected, we must furthermore have $n_j = \pm 1$. Hence, after appropriately changing the signs of some of the α_j , the T^ℓ -action is given by

$$(\exp(\theta_1 \xi_1 \dots \theta_\ell \xi_\ell)) \cdot (z_1, \dots, z_\ell) = (e^{2\pi \theta_1 i} z_1, \dots, e^{2\pi \theta_\ell i} z_\ell).$$

This normal form for the action has the following consequences:

- The quotient manifold is endowed with charts of the form $\mathbb{R}^{n-\ell} \times (\mathbb{C}^\ell)/T^\ell \simeq \mathbb{R}_\ell^n$, and is therefore a manifold with corners.
- The quotient map $f: M \rightarrow N$ in the above local coordinates is given by

$$f: T^{n-\ell} \times (\mathbb{R}^{n-\ell} \times \mathbb{C}^\ell) \rightarrow \mathbb{R}_\ell^n \\ (q, x, z_1, \dots, z_\ell) \mapsto (x, |z_1|^2, \dots, |z_\ell|^2).$$

By Lemma 3.8, we see that f is a boundary fibration with respect to the log divisor ∂N .

- Because the vanishing locus of the induced elliptic ideal is given by $f^{-1}(\partial N)$, it follows that the intersection stratification coincides with the orbit-type stratification.

- At points $p \in D[1]$, the isotropy group is given by S^1 , and therefore $N_p \mathcal{O}_p$ inherits an S^1 -action and consequently admits an orientation. We conclude that $D[1]$ is co-orientable.
- When M is oriented, a choice of orientation for T^n gives rise to an orientation for N by observing that $M \setminus D \rightarrow N \setminus \partial N$ is a principal T^n -bundle.
- When M is four-dimensional and the action is not free, it is shown in [27] that f admits a section. As a generic fiber and the image of this section intersect only once, it follows that the intersection pairing of the fiber with the image of this section is nonzero, and therefore f is homologically essential. ■

The group actions underlying toric manifolds satisfy the conditions of this proposition, leading to the following result.

Corollary 6.2 *Let (M^{2n}, ω) be a toric manifold, and let $f: M^{2n} \rightarrow \Delta^n$ denote the quotient map. Then f is a boundary fibration.*

In four dimensions, Proposition 6.1 provides us with fibrations that satisfy nearly all the assumptions required to apply Theorem 3.23. However, the torus action does not guarantee that the parity of the elliptic divisor is 1. To proceed, we must add hypotheses to ensure that this is the case.

Proposition 6.3 *Let $f: (M^4, \omega) \rightarrow \mathbb{R}^2$ be a toric manifold. Then the parity of the elliptic divisor obtained from Proposition 6.1 is 1, and therefore M admits a stable generalized complex structure compatible with f .*

Proof By Proposition 6.1, we have that f is a boundary fibration, and therefore by Theorem 3.23 the manifold M admits an elliptic symplectic structure. As each of the preimages of the faces of the moment polytope is a symplectic submanifold of (M, ω) , the symplectic structure provides each component of the elliptic divisor with a natural co-orientation for which the intersections have positive index. It follows that the parity of the elliptic divisor is 1. ■

6.2 Simple examples

We give examples of boundary fibrations obtained from torus actions which will serve as the building blocks for the connected sum procedure.

Example 6.4 ($\mathbb{C}P^2$) Consider the standard toric structure on $\mathbb{C}P^2$. Proposition 6.3 implies that f is a homologically essential boundary fibration and that $\mathbb{C}P^2$ admits an elliptic divisor with parity 1 (three lines intersecting at different points). Therefore, $\mathbb{C}P^2$ admits a stable generalized complex structure compatible with its moment map.

Example 6.5 ($\overline{\mathbb{C}P}^2$) We consider $\overline{\mathbb{C}P}^2$, i.e., $\mathbb{C}P^2$ with the orientation opposite to the standard complex structure. As an oriented manifold, this is not a toric manifold, but there is still a T^2 -action with connected isotropies present. Therefore, Proposition 6.1 implies that the quotient map is a homologically essential boundary fibration. Consequently, by Theorem 3.23, there exists a compatible elliptic symplectic structure with imaginary parameter on $\overline{\mathbb{C}P}^2$. The parity of the elliptic divisor is -1 , so this symplectic structure does not induce a stable generalized complex structure. As $\overline{\mathbb{C}P}^2$ is

not almost complex, it cannot have a stable generalized complex structure, and hence this problem cannot be remedied.

Example 6.6 ($S^2 \times S^2$) Let $(S^2 \times S^2)$ be given its standard toric structure, i.e., the symplectic form is the product of the standard area forms and T^2 acts on rotation by S^1 one each of the factors. Proposition 6.3 implies that the quotient map is a homologically essential boundary fibration and that $S^2 \times S^2$ admits a compatible stable generalized complex structure.

Example 6.7 (S^4) Consider $S^4 \subset \mathbb{C}^2 \times \mathbb{R}$, and let T^2 act in the standard way on \mathbb{C}^2 . This provides an effective T^2 -action on S^4 with connected isotropies. Therefore, by Proposition 6.1, we find that the quotient map is a homologically essential boundary fibration. Consequently, Theorem 3.23 implies the existence of a compatible elliptic symplectic structure with imaginary parameter on S^4 . The parity of the divisor is -1 . Just as $\overline{CP^2}$, S^4 is not almost-complex, so the index cannot be fixed by making different choices of divisor or orientations.

The following example of a boundary fibration appears also in [9].

Example 6.8 ($S^3 \times S^1$) There are two interesting T^2 -actions on $S^3 \times S^1$. First, consider $S^3 \subset \mathbb{C}^2$ as the unit sphere and restrict the natural T^2 -action on \mathbb{C}^2 to S^3 . This provides an effective T^2 -action on S^3 with S^1 isotropy at all points in the intersection with the coordinate hyperplanes. Extending the T^2 -action trivially to the S^1 -factor provides an effective T^2 -action on $S^3 \times S^1$ with only S^1 isotropy groups. The quotient map

$$f_1: (S^3 \times S^1, D_1) \rightarrow (I \times S^1, \{0, 1\} \times S^1)$$

then becomes a homologically essential boundary fibration by Proposition 6.1. Note that D_1 is given by the union of two disjoint tori.

Another T^2 -action on $S^3 \times S^1$ is obtained by letting one S^1 act by rotation on one of the coordinates of $S^3 \subset \mathbb{C}^2$, and let the other act by multiplication on S^1 . The quotient map

$$f_2: (S^3 \times S^1, D_2) \rightarrow (\mathbb{D}^2, \partial\mathbb{D}^2)$$

then again becomes a homologically essential boundary fibration by Proposition 6.1. In this case, D_2 is a single torus. In both cases, Theorem 3.23 implies the existence of a compatible elliptic symplectic structure with zero elliptic residue. Moreover, as the vanishing locus of the elliptic divisor is smooth and co-orientable, we obtain two stable generalized complex structures on $S^3 \times S^1$.

The example we consider next is more elaborate than the previous ones. The existence of stable generalized complex structures on these spaces is a consequence of the more general Theorem 2 from [31].

Example 6.9 ($(\#nS^1 \times S^2) \times S^1$) In [27], it is shown that for $2g + h > 1$, the manifold $M = (\#(2g + h - 1)S^1 \times S^2) \times S^1$ admits an effective T^2 -action with connected isotropy groups over a base, B , which is a surface of genus g with h small open disks removed. In fact, part of the action is just rotation of the last S^1 -factor, so this action has no fixed points (a fact that also follows from the Euler characteristic of M being 0).

By Proposition 6.1, we conclude that there exists a homologically essential boundary fibration

$$f: ((\#(2g + h - 1)S^1 \times S^2) \times S^1, D) \rightarrow (B, \partial B).$$

The degeneracy locus consists of h disjoint tori—precisely, the number of boundary components of B , and is in particular co-orientable. Consequently, by Theorem 3.23, there exists a compatible (smooth) stable generalized complex structure on M whose type change locus has h connected components.

To illustrate the elliptic-elliptic trade theorem we give some examples:

Example 6.10 ($\mathbb{C}P^2$) Applying Theorem 5.3 to Example 6.4 yields several boundary Lefschetz fibrations $f: (\mathbb{C}P^2, D) \rightarrow (N, \partial N)$. The number of elliptic–elliptic and Lefschetz singularities adds up to 3, but any combination is possible. See also Figure 1 and Remark 6.13.

Example 6.11 (S^4) Applying Theorem 5.3 to Example 6.7 yields a boundary Lefschetz fibration $f: (S^4, \tilde{D}) \rightarrow (\mathbb{D}^2, \partial\mathbb{D}^2)$ with two Lefschetz singularities. Because the parity of the original divisor on S^4 is -1 , the new divisor \tilde{D} will be non-co-orientable. Therefore, it is nonorientable, and as it admits an S^1 -fibration, it must then be a Klein bottle.

6.3 Main class of examples

Using the above examples as building blocks, we can now construct many more examples.

Theorem 6.12 *The manifolds in the following two families admit homologically essential boundary fibrations:*

- $X_{n,\ell} := \#n(S^2 \times S^2) \# \ell(S^1 \times S^3)$, with $n, \ell \in \mathbb{N}$,
- $Y_{n,m,\ell} := \#n\mathbb{C}P^2 \# m\overline{\mathbb{C}P}^2 \# \ell(S^1 \times S^3)$, with $n, m, \ell \in \mathbb{N}$,

whenever their Euler characteristic is nonnegative. Therefore, each of these manifolds admits a compatible elliptic symplectic structure, which induces a stable generalized complex structure if $1 - b_1 + b_2^+$ is even.

Proof In the previous section, we exhibited boundary fibrations on $\mathbb{C}P^2, \overline{\mathbb{C}P}^2$ and $S^2 \times S^2$ with 3, 3, 4 points in $D[2]$, respectively. Therefore, we may apply Theorem 4.6 inductively to obtain homologically essential boundary fibrations on $X_{n,0}$ and $Y_{n,m,0}$ for all possible values of n and m , including $n = m = 0$ by Example 6.7. The number of points in $D[2]$ for these manifolds is $2n + 2$ and $n + m + 2$, respectively. Therefore, we can apply Corollary 4.8, respectively, $n + 1$ and $\lfloor \frac{n+m+2}{2} \rfloor$ -times to obtain homologically essential boundary fibrations on $X_{n,\ell}$ and $Y_{n,m,\ell}$, for $\ell \leq n + 1, \lfloor \frac{n+m+2}{2} \rfloor$, respectively. A simple computation of the Euler characteristic of these manifolds shows that this is precisely when their Euler characteristic is nonnegative. The parity of the divisor in $\mathbb{C}P^2, \overline{\mathbb{C}P}^2$, and $S^2 \times S^2$ is 1, $-1, 1$, respectively. Therefore, Theorem 4.6 gives us that the parity of $X_{n,0}$ and $Y_{n,m,0}$ is $(-1)^{n-1}$. Corollary 4.8 gives us that the parity of the divisor in $X_{n,\ell}$ and $Y_{n,m,\ell}$ is $(-1)^{n-1+\ell}$. By Theorem 3.23, these manifolds admit

compatible elliptic symplectic structures. These induce stable generalized complex structures when $(-1)^{n-1+\ell} = 1$, which is to say that $1 - b_1 + b_2^+$ is even. ■

The following remarks elaborate on the assumptions on the n , m , and ℓ in the above theorem.

Remark 6.13 (Euler characteristic) The condition on the Euler characteristic is necessary. Indeed, a simple application of Mayer–Vietoris shows that if $f: M^4 \rightarrow \Sigma^2$ is a boundary Lefschetz fibration over a surface with k corners, and ℓ singular fibers, then $\chi(M) = k + \ell$. In particular, we find that the Euler characteristic of a manifold admitting a boundary Lefschetz fibration is necessarily nonnegative. Therefore, we conclude that we found all members of the families appearing in Theorem 6.12 that admit boundary fibrations.

Remark 6.14 (Betti numbers) The existence of a generalized complex structure on a manifold implies the existence of an almost-complex structure. Such a structure cannot exist when $1 - b_1 + b_2^+$ is odd, which explains that we found all members of the families appearing in Theorem 6.12 that admit stable generalized complex structures arising from boundary fibrations.

Remark 6.15 (Torus actions) Torus actions persist under taking connected sums of disjoint manifolds at fixed points [27]. In fact, [27] provides a classification of simply connected four-manifolds with effective torus actions and connected isotropy groups. The manifolds admitting such actions are precisely the manifolds $X_{n,0}$, $Y_{n,m,0}$, and S^4 appearing in Theorem 6.12. Whenever such a T^2 -action is present, it is possible to ensure that the elliptic symplectic structure arising from Theorem 6.12 is T^2 -invariant, and hence we obtained all such simply connected four-manifolds admitting T^2 -invariant stable generalized complex structures.

In the nonsimply connected case, [27] also provides a classification of effective nonfree torus actions with only S^1 -isotropy groups on compact oriented connected four-manifolds. It is proved that any of these manifolds is of the form as described in the Example 6.9, and hence we have also obtained all manifolds with such actions and T^2 -invariant stable generalized complex structures.

6.4 Relation to semitoric geometry

We finish by relating our results to semitoric geometry. Recall that a *focus–focus* singularity of a completely integrable system (M, ω, f) is a point $p \in M$ where there are Darboux coordinates (x_1, y_1, x_2, y_2) for ω in which f takes the form

$$(x_1, y_1, x_2, y_2) \xrightarrow{f} (x_1 y_2 - x_2 y_1, x_1 x_2 + y_1 y_2).$$

Semitoric manifolds [28] are generalizations of four-dimensional toric manifolds where the moment map, besides elliptic and elliptic–elliptic singularities, may also have focus–focus singularities. If we use the above Darboux coordinates to define complex coordinates

$$(w_1, w_2) = \frac{1}{4}(x_1 + y_2 + i(x_1 - y_2), x_1 - y_2 + i(x_1 + y_2)),$$

we see that the point p becomes a Lefschetz singularity of the moment map f .

Proposition 6.16 *Moment maps of semitoric manifolds are boundary Lefschetz fibrations. Consequently, semitoric manifolds admit compatible stable generalized complex structures, for which the elliptic divisor is the preimage of the boundary of the moment map image.*

Proof In light of Theorem 3.23 and the toric case (Proposition 6.3), we need only argue that the map is homologically essential. This follows because the homotopy type of $M \setminus D$ is obtained from a regular fiber by adding 2-cells along the vanishing cycles corresponding to each Lefschetz singularity. ■

Remark 6.17 Theorem 5.3 trades an elliptic–elliptic singularity for a Lefschetz singularity in the context of a fibration without further geometric structures. This is reminiscent of the nodal-trade/Hamiltonian Hopf bifurcation from semitoric geometry [32, 33] in the context of Lagrangian fibrations. In the Hamiltonian Hopf bifurcation, elliptic–elliptic singularities are traded for focus–focus singularities, which by the above are equivalent to Lefschetz singularities. However, these maps interact differently with the underlying geometric structure. Notably, in the semitoric version, the base of the fibration has a singular integral affine structure which helps with the extension of the fibration beyond a neighborhood of the singularities involved.

The converse trade for semitoric geometry, similar to our Theorem 5.4, appeared in [24]. There, the authors also make use of the singular integral affine structure present in such integrable systems.

Acknowledgment The authors would like to thank Conan Leung, Eduard Looijenga, Maarten Mol, and Álvaro Pelayo for useful discussions.

References

- [1] S. Akbulut and Ç. Karakurt, *Every 4-manifold is BLF*. J. Gökova Geom. Topol. 2(2008), 83–106.
- [2] D. Auroux, S. K. Donaldson, and L. Katzarkov, *Singular Lefschetz pencils*. Geom. Topol. 9(2005), 1043–1114.
- [3] R. İ. Baykur, *Kähler decomposition of 4-manifolds*. Algebr. Geom. Topol. 6(2006), 1239–1265.
- [4] R. İ. Baykur, *Existence of broken Lefschetz fibrations*. Int. Math. Res. Not. IMRN. (2008), no. rnn 101, 15 pp.
- [5] R. İ. Baykur, *Topology of broken Lefschetz fibrations and near-symplectic four-manifolds*. Pacific J. Math. 240(2009), no. 2, 201–230.
- [6] G. R. Cavalcanti, *Examples and counter-examples of log-symplectic manifolds*. J. Topol. 10(2017), no. 1, 1–21.
- [7] G. R. Cavalcanti and M. Gualtieri, *Blow-up of generalized complex 4-manifolds*. J. Topol. 2(2009), no. 4, 840–864.
- [8] G. R. Cavalcanti and M. Gualtieri, *Stable generalized complex structures*. Proc. Lond. Math. Soc. 116(2018), no. 5, 1075–1111.
- [9] G. R. Cavalcanti and R. L. Klaasse, *Fibrations and stable generalized complex structures*. Proc. Lond. Math. Soc. 117(2018), no. 6, 1242–1280.
- [10] G. R. Cavalcanti and R. L. Klaasse, *Fibrations and log-symplectic structures*. J. Symplectic Geom. 17(2019), no. 3, 603–638.
- [11] G. R. Cavalcanti, R. L. Klaasse, and A. Witte, *Self-crossing stable generalized complex structures*. J. Symplectic Geom. (2020), to appear. [arXiv:2004.07559](https://arxiv.org/abs/2004.07559).
- [12] M. Crainic, *Generalized complex structures and Lie brackets*. Bull. Braz. Math. Soc. (N.S.) 42(2011), no. 4, 559–578.

- [13] S. K. Donaldson, *Lefschetz pencils on symplectic manifolds*. J. Differential Geom. 53(1999), no. 2, 205–236.
- [14] J. B. Etnyre and T. Fuller, *Realizing 4-manifolds as achiral Lefschetz fibrations*. Int. Math. Res. Not. IMRN. (2006), no. 70272, 21 pp.
- [15] D. T. Gay and R. Kirby, *Constructing Lefschetz-type fibrations on four-manifolds*. Geom. Topol. 11(2007), 2075–2115.
- [16] R. E. Gompf, *Toward a topological characterization of symplectic manifolds*. J. Symplectic Geom. 2(2004), no. 2, 177–206.
- [17] R. E. Gompf and A. I. Stipsicz, *4-manifolds and Kirby calculus*, Graduate Studies in Mathematics, 20, American Mathematical Society, Providence, RI, 1999.
- [18] R. Goto and K. Hayano, *C^∞ -logarithmic transformations and generalized complex structures*. J. Symplectic Geom. 14(2016), no. 2, 341–357.
- [19] A. Grothendieck, *On the de Rham cohomology of algebraic varieties*. Publ. Math. Inst. Hautes Études Sci. 29(1966), no. 1, 95–103.
- [20] M. Gualtieri, *Generalized complex geometry*. Ann. of Math. (2) 174(2011), no. 1, 75–123.
- [21] M. Gualtieri, S. Li, A. Pelayo, and T. S. Ratiu, *The tropical momentum map: a classification of toric log symplectic manifolds*. Math. Ann. 367(2017), nos. 3–4, 1217–1258.
- [22] N. Hitchin, *Generalized Calabi–Yau manifolds*. Q. J. Math. 54(2003), no. 3, 281–308.
- [23] Y. Lekili, *Wrinkled fibrations on near-symplectic manifolds*. Geom. Topol. 13 (2009), no. 1, 277–318. Appendix B by R. İnanç Baykur.
- [24] N. C. Leung and M. Symington, *Almost toric symplectic four-manifolds*. J. Symplectic Geom. 8(2010), no. 2, 143–187.
- [25] R. B. Melrose, *The Atiyah–Patodi–Singer index theorem*, Research Notes in Mathematics, 4, A K Peters, Wellesley, MA, 1993.
- [26] E. Miranda and G. Scott, *The geometry of E-manifolds*. Preprint, 2018. [arXiv:1802.02959](https://arxiv.org/abs/1802.02959) [math.SG]
- [27] P. Orlik and F. Raymond, *Actions of the Torus on 4-Manifolds. I*. Trans. Amer. Math. Soc. 152(1970), no. 2, 531–559.
- [28] A. Pelayo and S. Vũ Ngọc, *Semitoric integrable systems on symplectic 4-manifolds*. Invent. Math. 177(2009), no. 3, 571–597.
- [29] P. Seidel, *Fukaya A_∞ -structures associated to Lefschetz fibrations. I*. J. Symplectic Geom. 10(2012), no. 3, 325–388.
- [30] R. Torres, *Constructions of generalized complex structures in dimension four*. Comm. Math. Phys. 314(2012), no. 2, 351–371.
- [31] R. Torres and J. Yazinski, *On the number of type change loci of a generalized complex structure*. Lett. Math. Phys. 104(2014), no. 4, 451–464.
- [32] J.-C. van der Meer, *The Hamiltonian Hopf bifurcation*, Lecture Notes in Mathematics, 1160, Springer, Berlin, 1985.
- [33] N. T. Zung, *Symplectic topology of integrable Hamiltonian systems, II: topological classification*. Compos. Math. 138(2003), no. 2, 125–156.

Department of Mathematics, Utrecht University, 3508 TA Utrecht, The Netherlands
 e-mail: g.r.cavalcanti@uu.nl aldo.witte@kuleuven.be

Department of Mathematics, KU Leuven, Celestijnenlaan 200B, BE-3001 Leuven, Belgium
 e-mail: r.l.klaasse@gmail.com