INSTRUCTION TO CONTRIBUTORS

The Journal of Helminthology publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance.

Manuscripts, which must be in English or French (with an English summary), should be addressed to:

The Editor, Journal of Helminthology, London School of Hygiene and Tropical Medicine, Winches Farm Field Station, 395 Hatfield Road, St Albans, Herts AL4 0XQ, England.

Two copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief Introduction; Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Research Notes should also be preceded by a brief abstract. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration may be printed is 12.0×18.5 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space, should be indicated. Some indication of scale should normally be given on the figure. Photocopies of illustrations should be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size 12.0×18.5 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

DUKE, B.O. L. (1971) The ecology of onchocerciasis in man and animals. In: *Ecology* and *Physiology of parasites* (editor, A. M. Fallis) pp. 213–222. Adam Hilger Ltd.: London.

JAMES, C. & WEBBE, G. (1973) A comparison of Egyptian and East African strains of Schistosoma haematobium. Journal of Helminthology, 47, 49-59.

25 offprints are provided free of charge; additional copies may be ordered at the proof stage.

Contents

	Pages
Some quantitative characters used in the identification of <i>Taenia hydatigena</i> , <i>T. ovis</i> , <i>T. pisiformis</i> and <i>T. multiceps</i> adult worms, and <i>T. multiceps</i> metacestodes. G. T. EDWARDS and I. V. HERBERT	1–7
Schistosoma bovis, S. intercalatum and S. haematobium: methods for in vivo radiolabelling of cercariae with radioselenium. N. Ø. CHRISTENSEN	9–12
Studies on Onchocerca tarsicola Bain and Schulz-Key, 1974 in reindeer (Rangifer tarandus) in northern Finland. G. BYLUND, HP. FAGERHOLM, C. KROGELL and S. NIKANDER	13–20
A biochemical study of adult and cystic stages of <i>Echinococcus granulosus</i> of human and animal origin from Kenya. D. P. McMANUS	21–27
Studies on amphistomiasis in Andhra Pradesh (India) VI. Immunization of lambs and kids with gamma irradiated metacercariae of <i>Cercariae indicae</i> XXVI. MD. HAFEEZ and B. V. RAO	29—32
Schistosoma mansoni: changes in the rate of tyrosin uptake by unisexual females after stimulation by males and male extracts. I. POPIEL and D. A. ERASMUS	33–37
Six platyhelminths from Malaysian reptiles including <i>Paradistomoidella cerperi</i> n.g., n.sp. (Digenea: Dicrocoeliidae). D. R. BROOKS and J. R. PALMIERI	39-43
Enhancement of Schistosoma mansoni infection induced by the methanol-extraction residue (MER) fraction of BCG. (Research Note). J. HAMBURGER and S. SAVION	45-48
A simplified radioimmunoassay technique for hydatid disease and human trichinosis. R. M. MATOSSIAN	49–57
Quantification of cryopreservation technique for <i>Onchocerca</i> microfilariae in skin snips. (Research Note). P. J. HAM and A. E. BIANCO	59–61
Resistance to Fasciola hepatica in calves harbouring primary patent Schistosoma bovis infections. S. B. SIRAG, N. Ø. CHRISTENSEN, P. NANSEN, J. MONRAD and F. FRANDSEN	63–70
Karyotypes and chromosome morphologies of <i>Megalodiscus temperatus</i> and <i>Philoph-thalmus gralli</i> . A. I. GROSSMAN and G. D. CAIN	71–78