

Research Paper

Cite this article: Dobbratz M, Gutknecht J, Wyse D, Sheaffer CC, Jungers JM (2022). Inconsistent effects of species diversity and N fertilization on soil microbes and carbon storage in perennial bioenergy cropping systems. *Renewable Agriculture and Food Systems* **37**, 60–70. <https://doi.org/10.1017/S1742170521000302>

Received: 21 March 2021
Revised: 18 June 2021
Accepted: 1 July 2021
First published online: 4 August 2021

Keywords:

Biofuels; carbon sequestration; extracellular enzyme activity; plant diversity; soil microbial carbon

Author for correspondence:

Jacob M. Jungers, E-mail: junge037@umn.edu

Inconsistent effects of species diversity and N fertilization on soil microbes and carbon storage in perennial bioenergy cropping systems

Michelle Dobbratz¹, Jessica Gutknecht², Donald Wyse¹, Craig C. Sheaffer¹
and Jacob M. Jungers¹ 

¹Department of Agronomy and Plant Genetics, University of Minnesota-Twin Cities, Saint Paul, MN, USA and

²Department of Soil, Water, and Climate, University of Minnesota-Twin Cities, Saint Paul, MN, USA

Abstract

Positive relationships between plant species diversity, soil microbial function and nutrient cycling have been well documented in natural systems, and these relationships have the potential to improve the production and sustainability of agroecosystems. Our objectives were to study the long-term effects of planted species composition and nitrogen (N) fertilization on soil microbial biomass C, extracellular enzyme activity, changes in total soil C, soil fertility and aboveground biomass yield in mixtures of native prairie species managed with and without N fertilizer for bioenergy production at four sites in Minnesota (MN), USA. Species were sown into mixture treatments and composition was not maintained (i.e., no weeding) throughout the duration of the study. Species mixture treatments at establishment included a switchgrass (*Panicum virgatum* L.) monoculture (SG), a four-species grass mixture (GM), an eight-species legume/grass mixture (LG) and a 24-species high diversity forb/legume/grass mixture (HD). Species diversity and aboveground productivity were similar for most mixture treatments at final sampling after 11 or 12 years of succession. Despite this homogenization of productivity and diversity throughout the study, the effects of planted species diversity and a decade of succession resulted in some differences in soil variables across species mixture treatments. On a peat soil in Roseau, MN, soil enzyme activities including β -glucosidase (BG), cellobiohydrolase (CBH) and phosphatase (PHOS) were highest in HD compared to GM treatments. On a sandy soil at Becker, MN, total soil C increased in all treatment combinations at the 0–15 and 15–30 cm depth intervals, with SG showing greater increases than HD at the 15–30 cm depth. Final soil pH also varied by species mixture at the Becker and Roseau sites, but differences in treatment comparisons varied by location. Nitrogen fertilization did not affect any response variable alone, but interacted with species mixture treatment to influence PHOS and total soil C at Becker. The inconsistent effects of species mixture and N fertilization on soil biological and chemical properties observed across sites highlight the importance of local soil and climate conditions on bioenergy and ecosystem service provisioning of perennial bioenergy cropping systems.

Introduction

Perennial cropping systems are increasingly studied for their capacity to balance provisioning services—such as biofuel, forage and food products—with regulating and supporting services—such as carbon sequestration and enhanced nutrient cycling (Werling *et al.*, 2014; Crews and Rumsey, 2017). Currently, less than 20% of global cropland is occupied by perennial crops (Cox *et al.*, 2006). The annual crops that dominate cropland are only grown for a portion of the year, exposing soils to potential erosion from water and wind. In contrast, perennial crops provide year-round soil cover and can have greater net primary productivity than annuals (Jarchow and Liebman, 2013) as they capture more photosynthetic radiation over the course of a year (Crews and DeHaan, 2015). This greater net primary productivity of perennial systems is used to support belowground structures, which can result in enhanced carbon sequestration (Lal, 2004; Crews and Rumsey, 2017), microbial community diversity (Culman *et al.*, 2010) and nutrient cycling (Crews, 2005). Even in small, targeted areas, native perennial cropping systems can provide multiple economic and ecosystem benefits when strategically integrated into croplands dominated by summer annual crops (Schulte *et al.*, 2017).

Soil carbon sequestration is an opportunity for agriculture to mitigate climate change and promote soil fertility, further supporting food production and environmental quality (Lal, 2004; Powlson *et al.*, 2011). Increased carbon storage of perennial cropping systems may be maximized by increasing species and functional group diversity (Fornara and Tilman, 2008).

© The Author(s), 2021. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

For example, adding C4 grasses and legumes increased carbon sequestration by 193 and 522% compared to monocultures of native perennial plants in a 12-year study in Minnesota, USA (Fornara and Tilman, 2008). Additionally, grasslands with a greater number of species from diverse functional groups may have greater net primary productivity (Loreau *et al.*, 2001), which can potentially result in carbon sequestration. While the relationship between diversity and productivity has been studied in native perennial systems, limited research has been conducted on carbon dynamics in agricultural land converted from annual crops to perennial polycultures. Managed perennial cropping systems can function differently than natural systems in response to plant diversity. For example, while greater species richness can lead to greater aboveground biomass in unmanaged systems (Loreau *et al.*, 2001), less diverse systems in an agronomically managed context, particularly switchgrass (*Panicum virgatum* L.) monocultures, have been shown to yield similar or more aboveground biomass than higher diversity plantings (Zilverberg *et al.*, 2014; Jungers *et al.*, 2015a). These findings are particularly relevant for the multifunctionality of managed lands because increased aboveground biomass is not just an indicator of greater provisioning services such as harvestable material—it can also imply greater regulating and supporting services such as carbon storage (Ferchaud *et al.*, 2016).

Soil microbial parameters are useful for integrating plant diversity/productivity with soil carbon storage potential for several reasons. Soil microbial biomass C (MBC) and microbial activity are often more responsive to agricultural management activities compared to slow-changing, recalcitrant C pools and can therefore be used as early indicators of soil organic matter change (Joergensen *et al.*, 1995; Mohammadi *et al.*, 2011). Various extracellular enzymes can be measured in soils to estimate specific soil microbial nutrient acquisition and decomposition activity. For example, soils with high levels of chitinase are expected to contain microbes active in N acquisition (Steinauer *et al.*, 2015). In addition to being sensitive to management on a short-term basis, microbial biomass and extracellular enzyme activities (EEA) are sensitive to changes in plant diversity and have been used to understand how plant species richness alters belowground activity. For example, in native prairies, plant diversity was positively correlated with cellobiohydrolase and 1,4-N-acetylglucosaminidase (Chung *et al.*, 2007). Similarly, increasing plant diversity in perennial systems increases soil microbial biomass (Wardle *et al.*, 1997; Thakur *et al.*, 2015; Chen *et al.*, 2019). Recent efforts to understand the mechanisms of soil carbon accumulation suggest that microbial biomass and extracellular enzyme pools are not only more responsive to short-term management, but may even be important drivers of increased soil organic carbon storage (Dwivedi *et al.*, 2019; Robertson *et al.*, 2019). For example, microbially-derived compounds assist in the association of organic matter to soil mineral particles to form a relatively persistent organic matter pool (Cotrufo *et al.*, 2019). Taken together, a broader analysis of species composition effects on microbial biomass and activity across different environments may provide greater insight regarding whether species composition leads to greater carbon storage in perennial agricultural systems.

Nitrogen (N) fertilization of perennial grassland bioenergy crops can increase productivity (Jungers *et al.*, 2015b) and affect soil microbial biomass and EEA (Li *et al.*, 2018; Yuan *et al.*, 2020). In a meta-analysis of various terrestrial ecosystems, Jian *et al.* (2016) showed that N fertilization increased hydrolase

activities which were positively associated with soil organic carbon. However, N fertilization at rates exceeding N removal with crop harvest has been linked to declines in soil C within annual crop rotations (Khan *et al.*, 2007). N fertilization can also reduce plant diversity in prairie plant communities (Isbell *et al.*, 2013), thus imparting an indirect effect on soil microbes and carbon dynamics (Revillini *et al.*, 2019). Quantifying the effects of N fertilization on soil microbial biomass and EEA across different plant species mixtures and locations varying in edaphic and climatic conditions can result in more accurate expectations of soil carbon dynamics across space.

Plant productivity, soil microbial biomass and activity, and soil carbon are important characteristics in perennial agricultural systems and may be influenced by plant species diversity and N fertilization. Our objective was to determine the effects of plant species diversity and composition (i.e., the number of species and functional groups of sown species in mixtures) and N fertilization on soil MBC, EEA, soil C change, and aboveground biomass productivity 11–12 years after the establishment of native prairie plant mixtures managed for bioenergy feedstock production. We hypothesized that increased plant species diversity and N fertilization would increase biomass yield and soil organic C by modulating soil physical and microbial parameters.

Materials and methods

Site description and experimental design

Data were collected from an existing perennial bioenergy cropping study established in 2006 and 2007 at Roseau, Becker, Lamberton and Waseca, Minnesota (Table 1). Details on field preparation, species selection and seeding rates for mixtures, and seedling establishment are described by Mangan *et al.* (2011). Treatments were arranged in a randomized complete block design with split plots and three replications per location. Main plots were fertilized with either 0 or 67 kg N ha⁻¹ yr⁻¹ by manually broadcasting ammonium nitrate each spring starting in 2008. Species mixture treatments were applied as sub plots and included switchgrass monoculture (SG), four-species grass mixture (GM), an eight-species legume/grass mixture (LG) and a 24-species legume/grass/forb mixture (HD). A list of individual species in each mixture is in Table S1. After sowing seeds for each species in the mixture treatments in June of 2006 for Becker, Lamberton and Waseca and 2007 for Roseau, the plant communities were not altered except for localized termination of perennial weeds [e.g., *Cirsium arvense* (L.) Scop]. All standing biomass was removed annually after senescence.

Plant diversity and aboveground biomass

Plant community composition was assessed annually from establishment to 2013 and in 2017 by recording the species identity and relative abundance as the percent ground cover occupied by each species in each plot. Species diversity was quantified using the Shannon diversity index (H'), where p_i is the proportion of species i based on percent cover. Aboveground biomass was measured by harvesting biomass to a height of 5 cm from two randomly placed 0.25-m² quadrats in each plot in October (following a killing frost of -2°C) in 2017. Biomass was dried at 60°C until constant moisture before weighing. Details on biomass harvest procedures can be found in Jungers *et al.* (2015a).

Table 1. Soil characteristics from 0 to 30.5 cm of the research locations prior to sowing perennial crop treatments in 2006 (Becker, Lamberton, Waseca) or 2007 (Roseau); summarized from Mangan *et al.* (2011).

Location	Soil type	Soil description	pH	Clay	OM	NO ₃ ⁻	S	P	K
				%	g kg ⁻¹	kg ha ⁻¹		mg kg ⁻¹	
Becker	Hubbard	Sandy, mixed, frigid Entic Haploboroll	6.6	7.2	14	8	11	30	108
Lamberton	Normania	Fine-loamy, mixed superactive mesic Aauic Hapludolls	5.7	33.9	36	14	18	71	601
Roseau	Bearden	Fine-silty, mixed, superactive, frigid Aeric Calciaquolls	7.2	9.8	16	7	24	18	121
Waseca	Webster	Fine-loamy, mixed, superactive, mesic Typic Endoaquolls	6.5	27.8	43	30	31	18	189

Soil collection and total soil carbon analysis

Data from soil sampling events during an initial baseline sampling in 2006–2007, a sampling from 2013 and a sampling in 2017 were used for this analysis. Baseline soil was collected at all sites in 2006 (southern sites including Becker, Lamberton and Waseca) and 2007 (Roseau) from 0 to 30 cm before initial site preparation or seeding (Mangan *et al.*, 2011).

Soil was collected in 2013 using a hydraulic soil probe (Giddings Machine Company, Inc., Colorado, USA) at all sites except Roseau (Jungers *et al.*, 2015a). Two 2.5-cm diameter cores were excavated from each plot, partitioned into 0–15, 15–30 and 30–60 cm depth intervals, and homogenized by depth interval in the field. Soil was dried at 60°C until constant weight prior to total C analysis.

Soil was collected in October 2017 using a hydraulic soil probe to a depth of 60 cm. Two cores from each plot, partitioned into 0–15, 15–30 and 30–60 cm depth intervals, were homogenized by depth interval in the field. A subsample of each depth was immediately frozen for extracellular enzyme analysis, and a subsample was kept at 4°C until microbial biomass could be determined. A subsample was also dried at 60°C until constant weight prior to total C analysis. Carbonates were measured with a 1 M HCl acid test. Total soil carbon C was determined using combustion analysis. Samples from 2017 were analyzed on an Elementar pyrocube (Elementar Americas, Inc., Ronkonkoma, NY, USA), with previous samples analyzed in a similar manner (Mangan *et al.*, 2011; Jungers *et al.*, 2017).

Microbial biomass carbon

MBC was determined by direct chloroform extraction using 0.5 M K₂SO₄ (Gregorich *et al.*, 1990; McDaniel and Grandy, 2016). For each soil sample, pairs of 10 ± 0.05 g sieved (4 mm) subsamples were weighed into 70 ml acid-washed test tubes. Gravimetric soil moisture was determined on a third 10 g sieved sample. In one sample from each pair, 1.0 ml chloroform was added, followed by the addition of 40 ml 0.5 M K₂SO₄ to both samples of each pair. All samples were then shaken at 250 rpm on an oscillating shaker for 4 h. After sediment settled, the supernatant was poured through rinsed #42 filters into acid-washed specimen cups, then transferred into labeled, unused scintillation vials. Extracts were immediately frozen until later analyzed on a TOC analyzer (TOC-V-CPN; Shimadzu Scientific Instruments Inc., Columbia, MD, USA) that determines the carbon concentration (ppm) of each extract. Microbial C was then calculated by

subtracting baseline values from values for fumigated samples and taking the soil mass into account. Microbial C is then reported as mg C kg soil⁻¹ (dry mass equivalent). Vials containing extracts from the Becker location were broken, and the entire site was extracted again from frozen soil. Although determining soil microbial biomass from frozen soil is not recommended, it has been shown that there is no difference in microbial biomass determined on frozen vs fresh soil in conditions similar to ours (Winter *et al.*, 1994).

Extracellular enzyme activity

Soil for analysis of EEA was kept frozen until the day of the assay. To determine EEA, the protocol outlined by German *et al.* (2011) was followed. In short, 0.5 ± 0.01 g soil was added to 50 ml of 50 mM pH 6.0 tris buffer. The soil slurry was sonicated and shaken vigorously immediately before 200 µl of slurry was added to 96-well plates in eight replicate wells. The activities of β-glucosidase (BG), N-acetyl-glucosaminidase (NAG), cellobiohydrolase (CBH) and acid phosphatase (PHOS) were determined by adding 50 µl of 4-methylumbelliferone linked enzyme substrates at saturating concentrations based on pre-optimization testing before the actual assay was performed (German *et al.*, 2011). Controls and standards included tris buffer alone (blank), substrate or soil slurry with tris buffer (negative controls), 4-methylumbelliferone with soil slurry (quench standards) and 4-methylumbelliferone with buffer (reference standards). After an hour of incubation at room temperature, 10 µl of NaOH solution was added to each well. Fluorescence was measured 1 min after NaOH addition on a microplate reading fluorometer with excitation filters at 365 and 450 nm. EEA was calculated as nmol substrate cleaved (as determined by converting fluorescence units using a reference standard curve) per gram of soil (oven dry equivalent) per hour.

Statistical analysis

Data were analyzed using linear mixed-effects models. Species mixture and N treatments were considered fixed categorical effects, and because of the split-plot design, N fertilization was also considered a random effect nested within block to account for pseudo-replication. Data from each location were analyzed separately because of differences in establishment year, soil types and weather. Statistical tests that included location as a fixed effect—along with all possible interactions between location and species mixture and N treatments—were conducted *ad hoc* to

quantitatively compare treatment effects across space. *F*-tests were used to determine the statistical significance of fixed effects and their interactions using $\alpha = 0.05$. Post-hoc mean comparisons were tested based on Tukey-adjusted *P* values to identify significant differences between levels of fixed effects. Pearson's correlation coefficients were calculated to determine relationships between plant community metrics (aboveground biomass and H') and final total soil C, and EEA. For correlation coefficients between any plant community variable and soil variable > 0.5 , a linear mixed-effects model was conducted to determine the direction and magnitude of the relationship. Statistical analysis was conducted in R using the nlme package (R Development Core Team, 2018; Pinheiro *et al.*, 2019).

Results

Effects of planted species richness and N fertilization on plant species diversity and aboveground biomass

The Shannon diversity index (H') varied by species mixture treatments only at the Waseca and Roseau locations. At Waseca, the SG treatment was more diverse (mean = 1.23, s.e. = 0.57) than the GM treatment (mean = 0.06, s.e. = 0.15), while all other treatments had similar H' (Table 2). At Roseau, H' of the HD treatment (mean = 0.65, s.e. = 0.12) was similar to the GM (mean = 0.76, s.e. = 0.10) but lower than the GL (mean = 1.30, s.e. = 0.14) and SG treatments (mean = 1.28, s.e. = 0.15). The three most abundant species based on percent cover for each treatment combination are presented in Supplemental Table S2.

At Waseca, biomass yield was affected by the interaction between species mixture and N fertilizer treatments (Table 2). A means comparison test of all combinations of species mixture and N fertilizer treatments showed that the unfertilized HD treatment yielded significantly more biomass (mean = 972.1 g m⁻², s.e. = 143.7) compared to the unfertilized GM treatment (mean = 237.1 g m⁻², s.e. = 83.9). Aboveground biomass yield was not affected by species mixture or N fertilization treatments at Becker, Lamberton or Roseau where yields averaged 561.0 (s.e. = 39.9), 479.2 (s.e. = 50.2) and 530.4 (s.e. = 72.8) g m⁻², respectively.

Effects of planted species richness and N fertilization on MBC and EEA

Species mixture and N fertilization treatments did not affect soil MBC at any location (Table 2) and averaged 993.7 (s.e. = 642.3), 318.5 (s.e. = 20.8), 1078.7 (s.e. = 285.2) and 113.1 (s.e. = 19.5) mg C kg soil⁻¹ at Becker, Lamberton, Roseau and Waseca, respectively. At Roseau, species mixture treatments had a significant effect on BG, CBH and PHOS enzyme activity levels (nmol g soil⁻¹ h⁻¹). The HD treatment had higher rates of BG than all other species mixture treatments, and higher CBH and PHOS activities compared to the GM treatment (Table 3). Nitrogen fertilization did not affect any enzyme activities (Table 2). Soil enzyme activities were not affected by either treatment at Lamberton or Waseca.

Species mixture treatments and the interaction between species mixture and N fertilization affected PHOS at Becker (Table 2), where only fertilized species mixture treatments affected PHOS levels. Fertilized SG had higher PHOS (mean = 230.0; s.e. = 28.0) than the fertilized HD (mean = 67.5; s.e. = 18.7) and fertilized LG treatments (mean = 103.0; s.e. = 12.9). Average PHOS activity

of the unfertilized treatments at Becker was 113 nmol g soil⁻¹ h⁻¹ (s.e. = 33.1). There were no treatment effects on NAG activity.

Effects of planted species richness and N fertilization on total soil C

At Becker there was a significant species mixture by N fertilization interaction for total C at 0–15 cm (Table 2); however, a means comparison test did not show significant differences. At Roseau and Waseca, species mixture effects were significant at the 30–60 cm depth. At Roseau, total soil C was greater in the GM and LG treatments compared to SG, and all three were similar to the HD treatment (Fig. 1). At Waseca, there was a significant interaction between species mixture and N fertilization treatments where the fertilized GM treatment had greater total soil C compared to the fertilized LG treatment but no other treatments were significantly different (Table 2; Fig. 2).

Treatment effects on the change in total soil C from the last soil sampling event (2013) to the end of the study (2017) only occurred at Becker (Table 2). At the 0–15 cm depth interval, there was an interaction between species mixture and N fertilization treatments; however, mean comparison tests did not indicate significant differences between any species mixture/N fertilizer treatment combinations (Fig. 2A). At the 30–60 cm depth, a species mixture by N fertilization interaction occurred due to a reduction in total soil C in the fertilized SG, which was different from the positive accrual of total soil C in the unfertilized GM (Fig. 2B).

Effects of planted species richness and N fertilization on soil chemical properties

There were significant effects of species mixture and the species mixture by N fertilizer interaction on soil pH at Becker and Roseau, but the treatment effects were inconsistent over the two locations (species mixture \times site *F*-value = 2.176, *P* = 0.037; Table 2; Fig. 3). At Becker, the LG treatment had a lower pH than the HD treatment, while the GM and LG treatments had a higher pH than the SG and HD treatments. At Roseau, the SG and HD treatments had a lower pH than the GM and LG treatments. The interaction between species mixture and N fertilizer at Roseau resulted in a significant difference between the fertilized SG (mean = 6.4, s.e. = 0.23) and unfertilized LG (mean = 7.4, s.e. = 0.15).

Species mixture treatment had a significant effect on soil P at Lamberton and Roseau and interacted with N fertilization at Waseca (Table 2). At Lamberton, soil P was lowest in the HD treatment (mean = 28.0 ppm, s.e. = 3.0) compared to the GM (mean = 35.2 ppm, s.e. = 5.3) and SG (mean = 36.8 ppm, s.e. = 5.0). At Roseau, soil P was lowest in the SG (mean = 2.0 ppm, s.e. = 0.2), compared to the LG and HD (same mean = 2.17 ppm, s.e. = 0.2), with the highest soil P in the GM (mean = 2.67 ppm, s.e. = 0.3). At Waseca, unfertilized SG had the highest soil P (mean = 13.3 ppm, s.e. = 1.3) compared to the unfertilized LG (mean = 5.7 ppm, s.e. = 0.9). There were no treatment effects on soil K.

Correlation analysis of response variables

Aboveground biomass was not correlated with any soil variable based on Pearson's correlation coefficients >0.5 or <-0.5 (Fig. 4). At Becker only, H' was negatively associated with BG

Table 2. ANOVA significance results (*P*-values) by site for aboveground biomass (AB), microbial biomass carbon (MBC), final soil pH, phosphorus (P) and potassium (K) at 0–15 cm, final (2017) total soil carbon at four depths (0–15, 15–30, 30–60, 60–90 cm), change in total soil carbon from 2013 to 2017 at four depths (0–15, 15–30, 30–60, 60–90 cm) and the activity of β -glucosidase (BG), cellobiohydrolase (CBH), N-acetylglucosaminidase (NAG), phosphatase (PHOS)

Location	Factor	MBC	Enzymes				Final total soil C			Change in total soil C			Soil properties			Plant community variables	
			BG	CBH	NAG	PHOS	0–15	15–30	30–60	0–15	15–30	30–60	pH	P	K	AB	H'
Becker																	
	Mixture (M)	0.490	0.624	0.486	0.916	0.018	0.351	0.169	0.572	0.714	0.075	0.006	0.022	0.695	0.270	0.916	0.058
	Nitrogen (N)	0.439	0.694	0.412	0.741	0.296	0.881	0.293	0.792	0.495	0.389	0.108	0.096	0.182	0.106	0.111	0.498
	M × N	0.483	0.333	0.774	0.083	0.012	0.041	0.179	0.135	0.007	0.464	0.033	0.097	0.808	0.350	0.857	0.025
Lamberton																	
	Mixture (M)	0.537	0.658	0.353	0.673	0.647	0.855	0.609	0.119	0.104	0.615	0.809	0.167	0.017	0.161	0.314	0.099
	Nitrogen (N)	0.349	0.894	0.815	0.804	0.726	0.726	0.298	0.752	0.703	0.995	0.702	0.197	0.251	0.332	0.370	0.369
	M × N	0.161	0.591	0.365	0.399	0.583	0.640	0.309	0.918	0.462	0.278	0.682	0.563	0.367	0.695	0.180	0.347
Roseau																	
	Mixture (M)	0.285	0.002	0.037	0.127	0.024	0.208	0.442	0.005	NA ^a	NA	NA	<0.001	0.046	0.405	0.499	0.007
	Nitrogen (N)	0.488	0.322	0.650	0.299	0.340	0.523	0.759	0.283	NA	NA	NA	0.072	0.397	0.137	0.520	0.274
	M × N	0.353	0.279	0.318	0.072	0.558	0.257	0.976	0.067	NA	NA	NA	0.001	0.517	0.342	0.882	0.848
Waseca																	
	Mixture (M)	0.505	0.688	0.858	0.981	0.769	0.480	0.635	0.008	0.156	0.864	0.291	0.738	0.056	0.884	0.061	0.028
	Nitrogen (N)	0.917	0.437	0.575	0.282	0.631	0.088	0.197	0.599	0.174	0.487	0.657	0.972	0.079	0.114	0.558	0.472
	M × N	0.540	0.487	0.354	0.193	0.211	0.855	0.822	0.026	0.978	0.167	0.514	0.954	0.036	0.461	0.046	0.335

^aNo soil collected at Roseau in 2013, therefore change in soil C not reported. *P*-values in bold text are significant 0.05.

Table 3. Mean and standard error of the mean (s.e.) of β -glucosidase (BG), cellobiohydrolase (CBH) and phosphatase (PHOS) soil enzyme activity ($\text{nmol g soil}^{-1} \text{h}^{-1}$) at the 0–15 cm soil depth interval in the switchgrass (SG), four-species grass mixture (GM), eight-species legume–grass mixture (LG) and high diversity (HD) species mixture treatments at Roseau, MN

Species mixture treatment	BG		CBH		PHOS	
	Mean	s.e.	Mean	s.e.	Mean	s.e.
SG	88.2b [†]	23.3	5.4ab	2.0	338.0ab	68.7
GM	57.6b	23.6	1.9b	0.3	187.9b	31.8
LG	68.8b	35.1	3.6ab	1.0	274.7ab	68.8
HD	174.0a	80.1	10.6a	3.2	457.2a	103.0

NAG activities were unresponsive to treatment and not included in this table (see Table 2).

[†]Means followed by the same letter within a column are not significantly different.




Fig. 1. Total soil carbon (C) in the 30–60 cm in the switchgrass (SG), four-species grass mixture (GM), eight-species legume–grass mixture (LG) and the 12-species high-diversity mixture (HD) at Roseau (A) and Waseca (B), MN. Mixture means are averaged over N fertilizer rates (0 or 67 kg N ha⁻¹) at Roseau and the mixture by N fertilizer rate interaction is shown for Waseca.

(Pearson's correlation = -0.50) and PHOS (Pearson's correlation = -0.58) enzyme activities.

Discussion

Aboveground biomass yield and species diversity

The effects of plant species diversity on productivity have been largely learned through experiments that carefully control and




Fig. 2. Change in total soil carbon (C) from 2013 to 2017 in the 0–15 cm (A) and 30–60 cm (B) soil profiles in the fertilized (67 kg N ha⁻¹) and unfertilized (0 kg N ha⁻¹) switchgrass (SG), four-species grass mixture (GM), eight-species legume–grass mixture (LG) and the 12-species high-diversity mixture (HD) at Becker, Minnesota.

maintain the number of species and relative abundance of species in mixtures (Tilman *et al.*, 2014). Our study is unique in that it represented a more typical scenario for a managed multispecies bioenergy crop; we did not maintain planted species richness or diversity but instead allowed the relative abundances of planted species to fluctuate and develop through time. Long-term patterns in species abundances and yields from this study can be found in Jungers *et al.* (2015a). In this study, plant surveys from the final year of the study (11 or 12 years after establishment) show (1) that some of the most abundant species in the mixture treatments were species that were not originally planted in the treatment, indicating that species recruitment was occurring, (2) plant diversity varied by species mixture and N fertilization across locations (a significant three-way interaction between species mixture, N fertilization and location; $F = 2.17$; $P = 0.037$) and (3) the most




Fig. 3. Soil pH in the top 0–15 cm in the switchgrass (SG), four-species grass mixture (GM), eight-species legume–grass mixture (LG) and the 12-species high-diversity mixture (HD) at Becker (A) and Roseau (B), Minnesota. Means are averaged over nitrogen fertilizer rates.

abundant species in the mixtures varied by location (Table S2). Recruitment of species into monocultures and loss of species in the polycultures contributed to the homogenization of H^+ across mixture treatments (Table 2). Despite not controlling for species richness or diversity, evidence of the positive diversity–productivity relationship commonly observed in natural system experiments was observed at the Waseca site, where the HD treatment yielded more biomass than the GM treatment.

Soil microbial C, extracellular enzymes and potential nutrient limitation

Although treatments were in place for more than a decade, we did not measure any differences in MBC across treatments. This was surprising given that a global meta-analysis recently demonstrated that as perennial stands age, a more pronounced positive relationship develops between plant species and MBC (Chen *et al.*, 2019) especially in stands >10 years old. Our results show that final plant community composition and diversity varied substantially from the planted species mixture treatments, and that these differences varied by location, which could explain why we did not observe a relationship between MBC and initial plant diversity treatments. We also saw no correlations between MBC, above-ground biomass or final plant diversity when measured at the end of the study.

Significant treatment effects on EEA were observed at the Roseau site, where we saw higher β -glucosidase (BG), cellobiohydrolase (CBH) and phosphatase (PHOS) activity in the HD treatment compared to all others. This was also one of the two sites where after 11–12 years, the HD treatment had lower plant diversity than the GL or SG treatments. Microbes are inherently carbon limited (Soong *et al.*, 2020) and BG and CBH are both carbon acquisition enzymes. At this northern Minnesota location, conditions would have been less ideal for microbial decomposition, with decreasing plant diversity over time with the dominance of grass species, high in recalcitrant carbon, microbial carbon limitation could have been higher. This in turn would have resulted in increased microbial investment in carbon acquisition enzymes, particularly BG which is known to be associated with more recalcitrant biomass (Caldwell, 2006). BG activity can be influenced by fertilization, but it can be relatively ephemeral in nature (Ajwa *et al.*, 1999; Bach and Hofmockel, 2015), thus the late fall timing of sampling may have precluded the observation of any differences between fertilized and unfertilized plots.

At the Roseau site, phosphatase (PHOS) activity was also highest in the HD treatment. The HD treatment at Roseau also had significantly lower pH and somewhat lower soil P, and was dominated by switchgrass more than other plant mixtures at that site or even across sites (Table S2). Grasses acquire phosphorus by relying heavily on acid phosphatases, organic acids and the fostering of microbial decomposition activity for P acquisition (Richardson and Simpson, 2011). Also supporting this idea, PHOS activity can be greater in soils under monocultures of grasses compared to monocultures of forbs (Fujita *et al.*, 2010). With a relatively lower pH from 7.0 to 6.8 on average in the HD, there also could have been a slight shift toward acid phosphatases, which are produced by plants and microbes, as opposed to purely microbially produced alkaline phosphatases (Spohn and Kuzyakov, 2013).

Nitrogen fertilization has been shown to consistently enhance PHOS activity (Marklein and Houlton, 2012), yet the effects of N on P uptake can be plant species-specific (Fujita *et al.*, 2010). Although we did not see a main effect of N fertilization on PHOS, there was an N fertilization \times mixture treatment interaction at Becker. Under N fertilized conditions, PHOS activity was greater in SG compared to LG and HD mixtures. Lower PHOS activity in the fertilized LG and HD mixtures compared to SG at Becker can be explained by the higher abundances of forb species in these treatments compared to SG (Table S2) and the notion that forbs are less dependent on extracellular enzymes for P acquisition than grasses. An interesting point at the Becker site is that plant species diversity was actually significantly negatively correlated with PHOS activity, and slightly positively correlated with soil pH (Fig. 4). The lack of mineralizable organic matter in the sandy soil at Becker could possibly tighten plant–soil–microbial relationships given that plant productivity is still just as high there. As mentioned, the most diverse plots at Becker were dominated by forbs, which likely do not produce or stimulate as much phosphatase production.

The plant–soil–microbial P acquisition relationships observed at Becker were also different than Roseau in terms of how plant species mixtures potentially altered soil pH. Soil acidification in the HD mixture at Roseau could have been driven by organic acid and phosphatase production by switchgrass, a dominant species in that mixture. At Becker where switchgrass never dominated mixtures as it did at the Roseau site, lower pH in the LG compared to the HD could have been associated with the higher




Fig. 4. Pearson's correlation coefficient matrix for microbial biomass C (MBC), β -glucosidase (BG), cellobiohydrolase (CBH), N-acetylglucosaminidase (NAG), phosphatase (PHOS), final total soil C at the 0–15, 15–30 and 30–60 depth intervals, soil pH and aboveground biomass (AB) and Shannon diversity index (H') at Becker (A), Lamberton (B), Roseau (C) and Waseca (D), MN.

legume abundance in LG (Fig. 4A). Effects of plant species on soil acidification may have been enhanced at Becker as a result of the lower buffering capacity inherent to the sandy soil there.

Total Soil C

We did not see changes in total soil C or differences in final total soil C across treatments at Lamberton, the site with the highest clay content and relatively high organic matter content

(Tables 1 and 2). The only location where changes in total soil C were observed was Becker, the location with the lowest clay and organic matter contents. The variability we observed in soil C across locations matches the patterns observed in previous studies. Studying an Alfisol and a Mollisol in the Upper Midwest, USA, Sprunger and Robertson (2018) found that the less fertile Alfisol was quicker to accumulate soil C across diversity levels, suggesting that differences in diversity effects on soil C were related to baseline soil C and clay content. Even long-term

biodiversity studies conducted on coarse-structured, low organic matter soils show minimal changes in soil C within the first 10 yrs of observation (Yang *et al.*, 2019). Changes in soil C after converting annual cropland to perennial grasslands can require decades to occur (McLauchlan *et al.*, 2006). Low soil C accumulation rates in early years following the transition from annual crops to perennials have been attributed to priming effects and subsequent losses in soil C (Jelinski and Kucharik, 2009; Cihacek and Ulmer, 2010). In our study, a comparison of final total soil C across treatments would capture a cumulative effect of the N fertilizer and species mixture treatments. Although our study allowed for more than 10 years of soil C treatment effects to accumulate, we found few changes in total soil C across treatments.

At Roseau, we observed greater total soil C at the 30–60 cm depth interval beneath GM and LG treatments compared to SG (Fig. 2A). Although we did not measure EEA below the 0–15 cm depth to match the total soil C differences, we do see that the GM and LG treatments had among the lowest BG and CBH activities. These differences relative to the other treatments do not exactly align with the relative differences in 30–60 cm total soil C, but this EEA trend could potentially help explain total soil C differences. Low BG and CBH activities indicate lower C mineralization of organic matter, a process that involves microbial respiration and subsequent C losses. Therefore, lower BG and CBH activities could indicate less decomposition, thus explaining the relatively higher total soil C observed in the 30–60 cm depth below the GH and LG treatments.

An interaction between species mixture and N fertilization treatments on final soil C at Waseca (Fig. 1B) could have been related to crop residue C:N content and mineralization potential. Final soil C was lower in the fertilized LG compared to the fertilized GM treatment. Fertilization may have increased the tissue N content of the grasses in LG, and in combination with N-rich tissues of legumes, may have resulted in lower C:N of tissues in LG compared to GM. The lower C:N of the fertilized LG residue may have decomposed more readily, thus resulting in lower soil C accumulation and final values compared to the GM treatment.

Our analysis of soil C change from 2013 to 2017 provides an indication of the more recent soil C accumulation trajectory. At Becker, we observed an increase in soil C from 2013 to 2017 at the 0–15 cm (Fig. 3A) depth. However, we observed soil C loss at the 30–60 cm depth in some treatment combinations including the fertilized SG and HD treatments. Differences in soil C accrual patterns by depth could be related to differences in the root depth distribution, whereas a concentration of high C:N root biomass at the 0–15 cm depth may have resulted in higher soil C accrual at that specific depth interval. Although belowground biomass was not sampled in this study, root tissue quality, biomass and distribution of biomass by depth can explain a large portion of variability in soil C dynamics compared to planted species diversity (Fornara and Tilman, 2008; Jungers *et al.*, 2017), thus explaining the lack of a species (Zemenchik and Albrecht, 2002) mixture treatment on total soil C in this study. Khan *et al.* (2007) also observed reductions in soil C at depths >30 cm under fertilized, long-term crop rotations including corn, oats, soybean and alfalfa.

Associations between plant community and soil variables

Plant community composition and diversity changed from the time that the species mixture treatments were established to the end of the study, which can explain the lack of differences in soil MBC, total C and EEA across species mixture treatments.

As a result, we tested for associations between soil variables and plant community metrics measured at the end of the study. We did not find any strong associations between plant community and soil variables. This analysis provided little evidence of a positive relationship between soil variables and plant diversity when measured at a single timepoint, with the few significant relationships already discussed above (see section ‘Soil microbial C, extracellular enzymes and potential nutrient limitation’). This can be explained by the potentially slow response of soil parameters to plant diversity. Soil variables measured at the end of this study may reflect the 11 or 12 years of accumulated plant diversity effects; effects that likely changed during succession throughout the life of the stands.

Conclusion

Soil microbial and C dynamics were inconsistent in response to the diversity of perennial plant mixtures and N fertilization in bioenergy cropping systems studied at four locations varying in soil and climate conditions after a decade or more since establishment. Inconsistencies across sites are likely related to site-specific changes in plant species composition through time, weather and soil conditions. For example, grass monocultures and mixtures dominated by grass species had higher PHOS activity compared to mixtures where forbs were more abundant. Similarly, we found evidence of soil C accrual at Becker, the location with the lowest clay and soil organic matter content. However, few soil microbial or C variables responded similarly across locations. By mimicking realistic management practices, such as not hand weeding or replanting absent species, we allowed plant succession to naturally introduce additional variability into this study to better reflect outcomes that would be observed in real-world production bioenergy fields. Increasing the number of sown species in perennial bioenergy cropping systems is likely to support other ecosystem services at various stages of the production lifespan; however, here we show that sown species diversity effects on soil C can vary widely across locations and through time.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S1742170521000302>

Acknowledgements. This work was supported by the Minnesota Turf Seed Council and the Minnesota Department of Agriculture. We thank Kevin Betts, Lindsay Wilson and Nicole Tautges for their technical assistance in the field and we also thank Fucui Li, Galen Bergquist, Carol Loopstra and Anna Crandall for assistance in the laboratory.

Conflict of interest. The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- Ajwa HA, Dell CJ and Rice CW (1999) Changes in enzyme activities and microbial biomass of tallgrass prairie soil as related to burning and nitrogen fertilization. *Soil Biology and Biochemistry* **31**, 769–777.
- Bach EM and Hofmockel KS (2015) Coupled carbon and nitrogen inputs increase microbial biomass and activity in prairie bioenergy systems. *Ecosystems* **18**, 417–427.
- Caldwell BA (2006) Effects of invasive scotch broom on soil properties in a Pacific coastal prairie soil. *Applied Soil Ecology* **32**, 149–152.
- Chen C, Chen HYH, Chen X and Huang Z (2019) Meta-analysis shows positive effect of plant diversity on microbial biomass and respiration. *Nature Communications* **10**, 1332.

- Chung H, Zak DR, Reich PB and Ellsworth DS** (2007) Plant species richness, elevated CO₂, and atmospheric nitrogen deposition alter soil microbial community composition and function. *Global Change Biology* **13**, 980–989.
- Cihacek L and Ulmer M** (2010) Effects of tillage on inorganic carbon storage in soils of the Northern Great Plains of the U.S. In Kimble JL, Lal R and Follett RE (eds), *Agriculture Practices and Policies for Carbon Sequestration in Soil*. Boca Raton, FL: CRC Press, pp. 63–70.
- Cotrufo MF, Ranalli MG, Haddix ML, Six J and Lugato E** (2019) Soil carbon storage informed by particulate and mineral-associated organic matter. *Nature Geoscience* **12**, 989–994.
- Cox TS, Glover JD, Van Tassel DL, Cox CM and DeHaan LR** (2006) Prospects for developing perennial grain crops. *BioScience* **56**, 649–459.
- Crews TE** (2005) Perennial crops and endogenous nutrient supplies. *Renewable Agriculture and Food Systems* **20**, 25–37.
- Crews TE and DeHaan LR** (2015) The strong perennial vision: a response. *Agroecology and Sustainable Food Systems* **39**, 500–515.
- Crews TE and Rumsey BE** (2017) What agriculture can learn from native ecosystems in building soil organic matter: a review. *Sustainability* **9**, 1–18.
- Culman SW, DuPont ST, Glover JD, Buckley DH, Fick GW, Ferris H and Crews TE** (2010) Long-term impacts of high-input annual cropping and unfertilized perennial grass production on soil properties and belowground food webs in Kansas, USA. *Agriculture, Ecosystems & Environment* **137**, 13–24.
- Dwivedi D, Tang J, Bouskill N, Georgiou K, Chacon SS and Riley WJ** (2019) Abiotic and biotic controls on soil organ-mineral interactions: developing model structures to analyze why soil organic matter persists. *Reviews in Mineralogy & Geochemistry* **85**, 329–348.
- Ferchaud F, Vitte G and Mary B** (2016) Changes in soil carbon stocks under perennial and annual bioenergy crops. *GCB Bioenergy* **8**, 290–306.
- Fornara DA and Tilman D** (2008) Plant functional composition influences rates of soil carbon and nitrogen accumulation. *Journal of Ecology* **96**, 314–322.
- Fujita Y, Robroek BJM, de Ruiter PC, Heil GW and Wassen MJ** (2010) Increased N affects P uptake of eight grassland species: the role of root surface phosphatase activity. *Oikos* **119**, 1665–1673.
- German DP, Weintraub MN, Grandy AS, Lauber CL, Rinkes ZL and Allison SD** (2011) Optimization of hydrolytic and oxidative enzyme methods for ecosystem studies. *Soil Biology and Biochemistry* **43**, 1387–1397.
- Gregorich EG, Wen G, Voroney RP and Kachanoski RG** (1990) Calibration of a rapid direct chloroform extraction method for measuring soil microbial biomass C. *Soil Biology and Biochemistry* **22**, 1009–1011.
- Isbell F, Reich PB, Tilman D, Hobbie SE, Polasky S and Binder S** (2013) Nutrient enrichment, biodiversity loss, and consequent declines in ecosystem productivity. *Proceedings of the National Academy of Sciences of the USA* **110**, 11911–11916.
- Jarchow ME and Liebman M** (2013) Nitrogen fertilization increases diversity and productivity of prairie communities used for bioenergy. *GCB Bioenergy* **5**, 281–289.
- Jelinski NA and Kucharik CJ** (2009) Land-use effects on soil carbon and nitrogen on a U.S. Midwestern floodplain. *Soil Science Society of America Journal* **73**, 217–225.
- Jian S, Li J, Chen J, Wang G, Mayes MA, Dzantor KE, Hui D and Luo Y** (2016) Soil extracellular enzyme activities, soil carbon and nitrogen storage under nitrogen fertilization: a meta-analysis. *Soil Biology and Biochemistry* **101**, 32–43.
- Joergensen RG, Anderson TH and Wolters V** (1995) Carbon and nitrogen relationships in the microbial biomass of soils in beech (*Fagus sylvatica* L.) forests. *Biology and Fertility of Soils* **19**, 141–147.
- Jungers JM, Clark AT, Betts K, Mangan ME, Sheaffer CC and Wyse DL** (2015a) Long-term biomass yield and species composition in native perennial bioenergy cropping systems. *Agronomy Journal* **107**, 1627–1640.
- Jungers JM, Sheaffer CC and Lamb JA** (2015b) The effect of nitrogen, phosphorus, and potassium fertilizers on prairie biomass yield, ethanol yield, and nutrient harvest. *BioEnergy Research* **8**, 279–291.
- Jungers JM, Eckberg JO, Betts K, Mangan ME, Wyse DL and Sheaffer CC** (2017) Plant roots and GHG mitigation in native perennial bioenergy cropping systems. *GCB Bioenergy* **9**, 326–338.
- Khan SA, Mulvaney RL, Ellsworth TR and Boast CW** (2007) The myth of nitrogen fertilization for soil carbon sequestration. *Journal of Environmental Quality* **36**, 1821–1832.
- Lal R** (2004) Soil carbon sequestration impacts on global change and food security. *Science (New York, N.Y.)* **304**, 1623–1627.
- Li J, Jian S, de Koff JP, Lane CS, Wang G, Mayes MA and Hui D** (2018) Differential effects of warming and nitrogen fertilization on soil respiration and microbial dynamics in switchgrass croplands. *GCB Bioenergy* **10**, 565–576.
- Loreau M, Naeem S, Inchausti P, Bengtsson J, Grime JP, Hector A, Hooper DU, Huston MA, Raffaelli D, Shmida B, Tilman D and Wardle DA** (2001) Biodiversity and ecosystem functioning: current knowledge and future challenges. *Science (New York, N.Y.)* **294**, 804–809.
- Mangan ME, Sheaffer C, Wyse DL, Ehlke NJ and Reich PB** (2011) Native perennial grassland species for bioenergy: establishment and biomass productivity. *Agronomy Journal* **103**, 509–519.
- Marklein AR and Houlton BZ** (2012) Nitrogen inputs accelerate phosphorus cycling rates across a wide variety of terrestrial ecosystems. *New Phytologist* **193**, 696–704.
- McDaniel MD and Grandy AS** (2016) Soil microbial biomass and function are altered by 12 years of crop rotation. *Soil* **2**, 583–599.
- McLauchlan KK, Hobbie SE and Post WM** (2006) Conversion from agriculture to grassland builds soil organic matter on decadal timescales. *Ecological Applications* **16**, 143–153.
- Mohammadi K, Heidari G, Khalesro S and Sohrabi Y** (2011) Soil management, microorganisms and organic matter interactions: a review. *African Journal of Biotechnology* **10**, 19840–19849.
- Pinheiro J, Bates D, DebRoy S and Sarkar D** (2019) nlme: linear and non-linear mixed effects models.
- Powelson DS, Gregory PJ, Whalley WR, Quinton JN, Hopkins DW, Whitmore AP, Hirsch PR and Goulding KWT** (2011) Soil management in relation to sustainable agriculture and ecosystem services. *Food Policy* **36**, 72–87.
- R Development Core Team** (2018) *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Revillini D, Wilson GWT, Miller RM, Lancione R and Johnson NC** (2019) Plant diversity and fertilizer management shape the belowground microbiome of native grass bioenergy feedstocks. *Frontiers in Plant Science* **10**, 1–18.
- Richardson AE and Simpson RJ** (2011) Soil microorganisms mediating phosphorus availability. *Plant Physiology* **156**, 989–996.
- Robertson AD, Paustian K, Ogle S, Wallenstein MD, Lugato E and Cotrufo FM** (2019) Unifying soil organic matter formation and persistence frameworks: the MEMS model. *Biogeosciences (Online)* **16**, 1225–1248.
- Schulte LA, Niemi J, Helmers MJ, Liebman M, Arbuckle JG, James DE, Kolka RK, O'Neal ME, Tomer MD, Tyndall JC, Asbjornsen H, Drobney P, Neal J, Van Ryswyk G and Witte C** (2017) Prairie strips improve biodiversity and the delivery of multiple ecosystem services from corn-soybean croplands. *Proceedings of the National Academy of Sciences* **114**, 11247–11252.
- Soong JL, Fuchsluger L, Marañon-Jimenez S, Torn MS, Janssens IA, Penuelas J and Richter A** (2020) Microbial carbon limitation: the need for integrating microorganisms into our understanding of ecosystem carbon cycling. *Global Change Biology* **26**, 1953–1961.
- Spohn M and Kuzyakov Y** (2013) Distribution of microbial- and root-derived phosphatase activities in the rhizosphere depending on P availability and C allocation—coupling soil zymography with 14C imaging. *Soil Biology and Biochemistry* **67**, 106–113.
- Sprunger CD and Robertson GP** (2018) Early accumulation of active fraction soil carbon in newly established cellulosic biofuel systems. *Geoderma* **318**, 42–51.
- Steinauer K, Tilman D, Wragg PD, Cesarz S, Cowles JM, Pritsch K, Reich PB, Weisser WW and Eisenhauer N** (2015) Plant diversity effects on soil microbial functions and enzymes are stronger than warming in a grassland experiment. *Ecology* **96**, 99–112.
- Thakur M, Milcu A, Manning P, Niklaus P, Roscher C, Power S, Reich P, Scheu S, Tilman D, Ai F, Guo H, Ji R, Pierce S, Ramirez N, Richter A,**

- Steinauer K, Strecker T, Vogel T and Eisenhauer N** (2015) Plant diversity drives soil microbial biomass carbon in grasslands irrespective of global environmental change factors. *Global Change Biology* **21**, 4076–4085.
- Tilman D, Isbell F and Cowles JM** (2014) Biodiversity and ecosystem functioning. *Annual Review of Ecology, Evolution, and Systematics* **45**, 471–493.
- Wardle DA, Bonner KI and Nicholson KS** (1997) Biodiversity and plant litter: experimental evidence which does not support the view that enhanced species richness improves ecosystem function. *Nordic Society Oikos* **79**, 247–258.
- Werling BP, Dickson TL, Isaacs R, Gaines H, Gratton C, Gross KL, Liere H, Malmstrom CM, Meehan TD, Ruan L, Robertson BA, Robertson GP, Schmidt TM, Schrottenboer AC, Teal TK, Wilson JK and Landis DA** (2014) Perennial grasslands enhance biodiversity and multiple ecosystem services in bioenergy landscapes. *Proceedings of the National Academy of Sciences of the USA* **111**, 1652–1657.
- Winter JP, Zhang Z, Tenuta M and Voroney RP** (1994) Measurement of microbial biomass by fumigation-extraction in soil stored frozen. *Soil Science Society of America Journal* **58**, 1645–1651.
- Yang Y, Tilman D, Furey G and Lehman C** (2019) Soil carbon sequestration accelerated by restoration of grassland biodiversity. *Nature Communications* **10**, 718.
- Yuan M, Duan J, Li J, Jian S, Gamage L, Dzantor KE, Hui D and Fay PA** (2020) Effects of nitrogen fertilization and bioenergy crop species on central tendency and spatial heterogeneity of soil glycosidase activities. *Scientific Reports* **10**, 19681.
- Zemenchik RA and Albrecht KA** (2002) Nitrogen use efficiency and apparent nitrogen recovery of Kentucky bluegrass, smooth brome grass, and orchardgrass. *Agronomy Journal* **94**, 421–428.
- Zilverberg CJ, Johnson WC, Owens V, Boe A, Schumacher T, Reitsma K, Hong CO, Novotny C, Volke M and Werner B** (2014) Biomass yield from planted mixtures and monocultures of native prairie vegetation across a heterogeneous farm landscape. *Agriculture, Ecosystems and Environment* **186**, 148–159.