



The association of overall diet quality with BMI and waist circumference by education level in Mexican men and women

Nancy López-Olmedo¹, Barry M Popkin¹, Michelle A Mendez² and Lindsey Smith Taillie^{1,*}

¹Department of Nutrition, Gillings School of Public Health, University of North Carolina, 123 West Franklin Street, CB # 8120 Carolina Square, Chapel Hill, NC 27516-3997, USA: ²Department of Environmental and Occupational Health, University of Pittsburgh, Pittsburgh, PA, USA

Submitted 3 May 2018: Final revision received 17 January 2019: Accepted 6 February 2019: First published online 13 June 2019

Abstract

Objective: The present study evaluated the association of two measures of diet quality with BMI and waist circumference (WC), overall and by education level, among Mexican men and women.

Design: We constructed two *a priori* indices of diet quality, the Mexican Diet Quality Index (MxDQI) and the Mexican Alternate Healthy Eating Index (MxAHEI), which we examined relative to BMI and WC. We computed sex-specific multivariable linear regression models for the total sample and by education level.

Setting: Mexico.

Participants: Mexican men (*n* 954) and women (*n* 1356) participating in the Mexican National Health and Nutrition Survey 2012.

Results: Total dietary scores were not associated with BMI in men and women, but total MxDQI was inversely associated with WC in men (−0.10, 95 % CI −0.20, −0.004 cm). We also found that some results differed by education level in men. For men with the lowest education level, a one-unit increase in total MxDQI and MxAHEI score was associated with a mean reduction in BMI of 0.11 (95 % CI −0.18, 0.04) and 0.18 (95 % CI −0.25, −0.10) kg/m², respectively. Likewise, a one-unit increase in total MxDQI and MxAHEI score was associated with a mean change in WC of −0.30 (95 % CI −0.49, −0.11) and −0.53 (95 % CI −0.75, −0.30) cm, respectively, in men with the lowest level of education. In women, the association of diet quality scores with BMI and WC was not different by education level.

Conclusions: Our findings suggest that a higher diet quality in men with low but not high education is associated with lower BMI and WC.

Keywords

BMI
Waist circumference
Diet quality
Mexican adults

Mexico is a middle-income country that has experienced a nutrition transition characterized by a decrease in the prevalence of different forms of undernutrition, whereas the prevalence of obesity has had one of the world's largest increases^(1,2). For instance, 76 % of women and 69 % of men had overweight or obesity in 2016, while 88 % of women and 65 % of men had abdominal obesity during the same year⁽³⁾.

Diets high in energy, saturated fat, sodium, refined carbohydrates or added sugars, but low in fruits, vegetables or whole-grain products, are thought to be the leading risk factors for morbidity and mortality from obesity and obesity-related diseases⁽⁴⁾. Previous studies in Mexico indicate that the consumption of energy-containing beverages

doubled from 1999 to 2006, whereas in 2012, the contribution of sugar-sweetened beverages and food products high in saturated fat or added sugars to total energy intake was about 10 and 16 %, respectively, in all age groups^(5,6). However, the approach of analysing single or a few nutrients or foods does not consider the complexity of dietary behaviours, as nutrients and foods are not eaten in isolation⁽⁷⁾.

Dietary indices, or scores, are the most common approaches to providing an overall rating of an individual's intake in reference to dietary recommendations used to examine the association between overall diet quality and health outcomes, including manifestation of type 2 diabetes and incidence or mortality due to CVD^(8–11). However, only

*Corresponding author. Email taillie@unc.edu



a few studies have analysed the relationship of overall diet quality with BMI or waist circumference (WC), finding inconsistent results^(12–15). Furthermore, there are pre-existing disparities in dietary intakes and obesity by socio-economic status in Mexico. In 2012, fruit and vegetable intake as well as the intakes of saturated fat and added sugars (including sugar-sweetened beverages) were higher among those with high compared with low socio-economic status^(6,16). Likewise, the prevalence of obesity was higher in adults with higher socio-economic status⁽¹⁷⁾. Nevertheless, it has not been explored whether the association of diet quality with BMI and WC differs by socio-economic status. Therefore, it is relevant to understand what role education level, a proxy of socio-economic status, plays in the relationship of diet quality with BMI and WC. Finally, the relationship of diet quality with BMI and WC stratified by sex has been barely studied, despite previous evidence showing that women may be more concerned with the quality of food⁽¹⁸⁾ and health consciousness might vary by education level⁽¹⁹⁾. Furthermore, in obesity research as well as in all biomedical research, it is recognized that weight gain as well as adipose tissue storage and metabolism may vary by sex^(20–22). In the present paper, we examine the association of diet quality with BMI and WC, overall and by education level, among Mexican men and women. The paper uses two different dietary quality indices: one based on the Mexican Dietary Guidelines (MxDG) and the other on foods and nutrients associated with cardiometabolic risk. The latter, the Alternate Healthy Eating Index (AHEI), is the only diet quality index validated against many different non-communicable diseases⁽²³⁾.

Methods

Study design and population

We used data from the National Health and Nutrition Survey (ENSANUT; from its Spanish acronym, Encuesta Nacional de Salud y Nutrición) 2012. We obtained information about sociodemographic characteristics, nutrition and health of 96 031 people from 50 528 randomly selected households. Dietary collection and assessment have been described elsewhere, but in brief, we collected dietary information in a random sub-sample (n 10 886) representing the national, regional (North/Central/South) and urban/rural population⁽²⁴⁾. We used the 24 h dietary recall developed by the US Department of Agriculture, adapting it to the Mexican context^(25,26). We calculated energy and nutrient intakes using the food composition database compiled by the National Institute of Public Health^(27–30).

We included non-pregnant and non-lactating adults aged 20–69 years with dietary information (n 2676). We excluded those without weight, height or WC measurements (n 107), without information on type 2 diabetes or smoking status (n 282), and without information on parity (n 3). Moreover, adults classified as underweight

(BMI < 18.5 kg/m²) were excluded (n 28). Although there is no established definition of low WC, we also decided to exclude men and women if their WC was ≤ 65 and ≤ 55 cm, respectively (n 4). Finally, we excluded those with a ratio of total energy intake to estimated energy requirement (in logarithmic scale) below -3 SD and above $+3$ SD (n 30), as previously described⁽¹⁶⁾. Some individuals were excluded for two or more variables; therefore, the total number of excluded participants does not equal the sum of all the excluded participants by variable. The analytic sample was composed of 2310 adults.

Variable definitions

BMI and waist circumference

Body weight was measured with light clothing using digital scales (model 872, Seca) with 0.1 kg precision and height using stadiometers (Dyna-top, model E-1, Mexico) with 0.1 cm precision. WC was measured using a fibreglass tape at the midpoint between the highest part of the iliac crest and the lowest part of the ribs' margin of the median axial line⁽³¹⁾. Trained personnel took the measurements using standard procedures^(32,33). We calculated BMI using the standard equation and categorized it based on WHO definitions⁽³⁴⁾. Men and women with a WC of ≥ 90 and ≥ 80 cm, respectively, were also categorized as having abdominal obesity, using as reference the classification by the International Diabetes Federation⁽³⁵⁾.

Mexican Diet Quality Index

We developed the Mexican Diet Quality Index (MxDQI) based on the MxDG, which were published in 2015⁽³⁶⁾. These guidelines recommend the number and size of servings for nine food groups, by age group and by total energy intake. We used the number of servings recommended for adults with a total energy intake of 8368 kJ/d (2000 kcal/d) as reference (for more details see the online supplementary material, Supplemental Table 1). We created thirteen MxDQI components based on all food groups except tap water, because the MxDG include only a suggested range of water consumption since water needs can vary by age, physical activity and weather (Table 1). Rather than number of servings, we used cut-off points recommended by the WHO, as well as recommendations for fat intake for the Mexican population, to define minimum and maximum scores for polyunsaturated fat, saturated fat and added sugars^(37–39). Finally, we considered sodium intake as an MxDQI component because the MxDG, consistent with international guidelines, recommend consuming no more than 2000 mg Na/d. The minimum score for sodium was based on the results of systematic reviews and meta-analyses⁽⁴⁰⁾. We defined scores between 0 (non-compliance) and 15 (intakes close to recommended) for each component. Specifically, we assigned a maximum score of 5 to those MxDQI components derived from the same food group (e.g. whole-grain and refined-grain

**Table 1** Mexican Diet Quality Index (MxDQI) and Mexican Alternate Healthy Eating Index (MxAHEI) components and criteria for scoring

Food component	MxDQI			MxAHEI		
	Maximum points	Criteria for minimum score (0)	Criteria for maximum score	Maximum points	Criteria for minimum score (0)	Criteria for maximum score
Adequacy						
Vegetables	10	0 servings	≥3 servings/2000 kcal	9	0 servings	≥5 servings
Whole fruit	10	0 servings	≥3 servings/2000 kcal	9	0 servings	≥4 servings
Whole-grain cereals						
Women	5	0 servings	≥3 servings/2000 kcal	9	0 g	≥75 g
Men						≥90 g
Legumes	10	0 servings	≥2 servings/2000 kcal	5	0 servings	≥1 serving
Seafood, poultry or eggs	5	<1 serving/2000 kcal†	≥2 servings/2000 kcal			
Nuts				5	0 servings	≥1 serving
Low-fat dairy	5	0 servings	≥3.5 servings/2000 kcal			
Polyunsaturated fat*	5	<6% of total energy intake	>10% of total energy intake	9	≤2% of total energy intake	≥10% of total energy intake
Long-chain (n-3) fats (EPA + DHA)				9	0 mg	≥250 mg
Moderation						
100% Fruit juices	5	>250 ml/2000 kcal	≤125 ml/2000 kcal			
Sugar-sweetened beverages				9	≥1 serving	0 servings
Refined grains	5	>3 servings/2000 kcal	≤1 serving/2000 kcal			
Red and processed meat	5	>1.5 servings/2000 kcal	≤0.5 serving/2000 kcal	9	≥1.5 serving	0 servings
Added sugars	15	>10% of total energy intake	<5% of total energy intake			
Sodium	15	>2 g/2000 kcal	≤1.5 g/2000 kcal	9	>2 g	≤1.5 g
Saturated fat	5	>10% of total energy intake	<7% of total energy intake			
Trans fat				9	≥4% of total energy intake	≤0.5% of total energy intake
Alcohol						
Women				9	≥2.5 drinks	0.5–1.5 drinks
Men					≥3.5 drinks	0.5–2.0 drinks
Total	100	0	100	100	0	110

*This dietary component in MxAHEI does not include long-chain (n-3) fats (EPA + DHA).

†2000 kcal = 8368 kJ.

cereals). We also assigned the added sugars and sodium components a maximum score of 15, given their high consumption in the Mexican population and therefore their potential impact on health (Table 1)^(41,42).

Mexican Alternate Healthy Eating Index

The AHEI includes twelve components scored from 0 (worst) to 5 or 10 (best), and the total AHEI-2010 score ranges from 0 (non-adherence) to 110 (perfect adherence)⁽²³⁾. For developing the Mexican Alternate Healthy Eating Index (MxAHEI), we used the original AHEI-2010 criteria for minimum and maximum score, except for sodium. We considered the same sodium criteria as for the MxDQI, since the cut-offs for sodium in AHEI-2010 were based on deciles of distribution in the population⁽²³⁾. Furthermore, we based serving sizes on the Mexican System of Food Equivalents⁽⁴³⁾. We scaled up the MxAHEI to 100 for comparisons with MxDQI (Table 1).

Covariates

The ENSANUT 2012 captured several variables through questionnaire; for our analysis, we considered education level, type 2 diabetes status, age, sex, parity, area and region of residence, assets index, type of housing, number of individuals in a household, marital status, smoking status and physical activity. We defined education level as no reading/writing skills, reading/writing skills or 3–9 years of school (elementary and middle school) and ≥10 years of school (high school or more), based on the reported grade completed and whether participants knew how to read and write. We selected these categories based on studies about the social and economic impact of illiteracy (including individuals with ≤2 years of elementary school) in Latin American and Caribbean regions^(44,45). The categorization of type 2 diabetes status (yes/no) was based on previous diagnosis by a doctor. We classified parity as none, 1–2, 3–4 and ≥5 pregnancies. Likewise, we categorized locations with <2500 inhabitants as rural and



locations with ≥ 2500 inhabitants as urban, and we defined regions as North, Central and South.* An assets index was constructed using factor analysis, where factor scores were estimated using a principal components approach, applied to household characteristics and assets⁽⁴⁶⁾. The index score was computed for each respondent and respondents were then classified into three categories (low, medium and high) using tertiles of the distribution of the assets index scores as cut-off points. Other variables were categorized as follows: type of housing as owned, rented or other (individuals occupied the property in another situation not specified); number of individuals in a household as 1–2, 3–4 or ≥ 5 ; and marital status as married/living together, divorced/separated/widowed or single. We categorized smoking status as current, former and never based on the questions: (i) 'Have you ever smoked at least one hundred cigarettes (5 packs) in your life?' and (ii) 'How many cigarettes you are smoking currently?' (possible answers: 'I am not currently smoking' or number and frequency). Finally, we assessed physical activity using the Spanish short version of the International Physical Activity Questionnaire and classified it based on WHO recommendations⁽⁴⁷⁾.

Statistical analysis

We conducted all analyses using the statistical software package Stata version 14.0 (2015). We used survey commands to account for survey design and weighting to generate nationally representative results. Statistical tests were two-tailed and considered significant at $P < 0.05$. We examined characteristics of anthropometric measurements, total diet scores, sociodemographic variables and lifestyle behaviours by sex and education level. We first performed sex-specific multivariable linear regression models for testing the statistically significant associations of MxDQI and MxAHEI scores, total and by component, with BMI and WC, adjusting for age (quadratic), total energy intake, smoking status, type 2 diabetes status, parity, area and region of residence, tertile of assets index, type of housing, number of individuals in a household, marital status and education level. Furthermore, we adjusted MxDQI models for alcohol intake. Models including dietary components as outcome variables were also adjusted for the other dietary components to account for the potential correlation among all components. To test whether the association of diet quality with BMI and WC was different for individuals with different education levels, we performed models that included interaction terms between diet score and the three defined levels of education. We performed global Wald

tests to determine whether any diet score coefficients differed across the three education levels in men and women.

Sensitivity analyses

We did not include physical activity in the main analyses due to the poor validity of the International Physical Activity Questionnaire short form for assessing moderate-to-vigorous physical activity among Mexican adults⁽⁴⁸⁾. However, physical activity could be an important confounder of the relationship between diet quality and anthropometry. Thus, we conducted sensitivity analyses to test whether the inclusion of crude physical activity data in models altered the associations of dietary indices with BMI and WC. Second, we conducted analyses in which corn tortilla was treated as a refined grain instead of a whole grain, since it is uncertain whether all corn tortillas are made with whole grains. Third, a set of sensitivity analyses focused on different approaches for examining the association of dietary components with BMI and WC. We tested two additional types of models, models that were not adjusted for other dietary components and models that adjusted for total score minus the dietary component of interest, to assess whether estimations were different from those obtained when adjusting for the other dietary components.

Results

Overall, the prevalence of overweight was slightly lower in women than in men, but the prevalence of obesity and abdominal obesity was higher in women than in men, as observed in the entire ENSANUT 2012 sample. Moreover, the mean of total MxDQI and MxAHEI score was about 40 in men and women; however, the total dietary scores were higher in men and women with the lowest education level. A higher proportion of men and women with low education level met the scoring criteria of several dietary components, including whole-grain cereals, legumes, red and processed meat, and saturated fat (Table 2).

Association of MxDQI and MxAHEI with BMI in men and women

The total MxDQI score was not associated with BMI among men and women. However, the vegetables as well as the seafood, poultry or eggs components were inversely associated with BMI in men. Likewise, the total MxAHEI score was not associated with BMI in both men and women. Nevertheless, the red and processed meat component was inversely associated with BMI among men, whereas the alcohol component was positively associated with BMI in women (Table 3).

*States by region are as follows. North: Baja California, Chihuahua, Coahuila, Durango, Nuevo León, Sinaloa, Sonora, Tamaulipas and Zacatecas; Central: Aguascalientes, Ciudad de México, Colima, Estado de México, Guanajuato, Hidalgo, Jalisco, Michoacán, Querétaro, San Luis Potosí and Tlaxcala; South: Campeche, Chiapas, Guerrero, Morelos, Oaxaca, Puebla, Tabasco, Veracruz and Yucatán.

**Table 2** Characteristics of Mexican men and women by education level. ENSANUT 2012 (*n* 2310)

	Men (<i>n</i> 954)						Women (<i>n</i> 1356)					
	No reading/writing skills (<i>n</i> 100)		Reading/writing skills or 3–9 years of school (<i>n</i> 621)		≥10 years of school (<i>n</i> 233)		No reading/writing skills (<i>n</i> 165)		Reading/writing skills or 3–9 years of school (<i>n</i> 910)		≥10 years of school (<i>n</i> 281)	
	Mean or %	SE	Mean or %	SE	Mean or %	SE	Mean or %	SE	Mean or %	SE	Mean or %	SE
Age (years)	48	2.0	43	0.9	36	1.4	51	1.4	43	0.6	36	0.9
BMI (kg/m ²)	27	0.5	27	0.3	27	0.5	29	0.8	30	0.2	29	0.5
Waist circumference (cm)	93	1.7	95	0.9	95	2.1	92	1.7	94	0.6	90	1.1
Total MxDQI score	45	2.2	42	0.9	35	1.3	50	2.0	40	0.7	37	1.0
Total MxAHEI score	42	1.5	40	0.6	37	0.8	45	1.3	40	0.4	38	0.7
BMI category												
Normal weight	34	6.5	37	3.0	34	4.6	33	6.0	20	1.9	25	3.6
Overweight	42	7.2	38	2.9	43	4.6	30	5.3	40	2.6	39	4.1
Obesity	24	6.6	25	2.3	24	3.7	36	5.6	41	2.5	36	4.0
Abdominal obesity	59	7.1	60	3.0	62	4.7	81	5.8	89	1.4	77	3.6
Proportion of individuals meeting the scoring criteria												
MxDQI components												
Vegetables	25	6.2	17	2.3	12	2.7	25	4.7	19	1.9	28	3.7
Whole fruit	6	2.4	7	1.9	11	2.7	9	2.6	6	1.2	7	1.9
Whole-grain cereals	82	5.0	74	2.6	55	4.8	86	3.7	71	2.3	45	3.9
Legumes	30	6.5	14	1.8	9	2.3	26	4.3	11	1.4	3	1.2
Seafood, poultry or eggs	30	6.1	37	2.8	33	4.3	33	5.3	43	2.5	38	4.0
Low-fat dairy	0	0.0	1	0.5	2	1.4	2	1.4	1	0.4	2	1.4
Polyunsaturated fat	66	6.6	71	2.8	67	4.7	69	5.6	62	2.5	69	3.9
100% Fruit juices	98	1.1	91	1.7	83	3.2	94	2.1	90	1.4	88	2.6
Refined grains	3	2.4	2	0.8	2	1.1	1	0.4	2	0.6	1	0.7
Red and processed meat	53	6.9	44	3.0	38	4.7	70	5.9	52	2.6	37	4.1
Added sugars	27	6.2	19	2.1	14	3.5	34	5.4	19	1.9	13	2.3
Sodium	23	5.2	22	2.4	13	3.3	30	5.4	19	1.9	13	2.8
Saturated fat	45	7.3	32	2.8	11	2.3	43	5.5	24	2.1	11	2.5
MxAHEI components												
Vegetables	4	2.8	7	1.5	4	2.0	4	2.1	4	1.0	7	2.2
Whole fruit	2	1.6	4	1.5	7	2.5	2	1.2	2	0.6	3	1.1
Whole-grain cereals	86	4.6	74	2.5	64	4.6	82	4.2	72	2.1	48	3.8
Legumes	38	6.8	33	2.5	19	3.3	35	5.1	21	2.0	12	2.2
Nuts	1	0.5	1	0.6	2	1.2	0	0.3	1	0.3	2	1.0
Polyunsaturated fat	41	6.9	34	3.0	29	4.2	32	6.0	26	2.2	34	4.0
Long-chain (<i>n</i> -3) fats (EPA + DHA)	7	2.7	6	1.3	7	2.7	4	1.5	4	1.0	11	3.0
Sugar-sweetened beverages	23	6.3	12	1.7	10	2.9	23	4.7	14	1.7	12	2.5
Red and processed meat	50	6.9	40	2.9	31	4.6	66	6.0	46	2.6	32	4.0
Sodium	27	5.5	21	2.3	15	3.4	52	6.1	35	2.3	22	3.3
Trans fat	98	1.3	95	1.1	85	4.0	96	1.8	91	1.8	89	3.1
Alcohol	3	1.7	3	0.8	4	2.4	0	0.0	0	0.0	0	0.0
Parity category												
None	–	–	–	–	–	–	65	5.5	38	2.6	39	4.0
1–2	–	–	–	–	–	–	6	3.4	23	2.1	36	3.8
3–4	–	–	–	–	–	–	6	2.3	17	2.1	20	3.5
≥5	–	–	–	–	–	–	24	4.6	22	1.9	5	1.6
Area												
Urban	47	7.1	68	2.4	88	2.3	56	5.6	73	1.7	84	2.8
Rural	53	7.1	32	2.4	12	2.3	44	5.6	27	1.7	16	2.8
Region												
North	12	4.0	20	1.8	23	3.2	11	3.1	20	1.4	18	2.6
Central	46	7.2	44	3.0	50	4.4	38	6.4	50	2.2	51	3.8
South	42	6.7	35	2.6	27	3.5	52	6.0	30	1.9	31	3.3
Tertile of assets index												
Low	62	7.4	39	2.7	13	2.8	59	6.1	30	2.1	11	2.2
Medium	27	7.1	33	2.9	25	3.9	26	5.1	35	2.3	20	2.9
High	11	4.4	28	3.0	63	4.4	15	5.8	35	2.7	69	3.4
Type of housing												
Owned	84	6.0	76	3.0	78	3.8	90	3.0	79	2.2	82	2.5
Rented	11	5.9	11	2.4	8	2.7	3	1.3	7	1.1	8	1.7
Other*	5	2.0	13	2.0	13	3.0	7	2.7	14	1.8	10	2.0

**Table 2** (Continued)

	Men (n 954)						Women (n 1356)					
	No reading/writing skills (n 100)		Reading/writing skills or 3–9 years of school (n 621)		≥10 years of school (n 233)		No reading/writing skills (n 165)		Reading/writing skills or 3–9 years of school (n 910)		≥10 years of school (n 281)	
	Mean or %	SE	Mean or %	SE	Mean or %	SE	Mean or %	SE	Mean or %	SE	Mean or %	SE
Number of individuals in a household												
1–2	29	5.6	13	1.5	10	2.6	17	3.2	13	1.4	8	1.7
3–4	29	6.6	39	2.7	56	4.8	49	5.8	42	2.4	54	4.2
≥ 5	42	7.2	49	2.9	34	4.7	34	5.5	45	2.6	38	4.2
Marital status												
Married/living together	74	7.0	81	2.6	59	4.8	67	6.0	71	2.3	55	4.2
Divorced/separated/widowed	7	2.7	4	0.9	8	2.7	19	5.3	17	2.0	11	3.1
Single	20	6.3	15	2.4	34	4.8	13	4.8	12	1.6	33	3.9
Smoking status												
Never	53	7.3	46	2.8	54	4.8	91	3.2	84	2.0	72	3.8
Former	22	5.3	29	2.6	25	4.1	3	1.8	11	1.8	13	2.7
Current	25	6.6	25	2.4	21	4.2	7	2.8	5	1.0	14	3.2
Physical activity												
	n 97		n 593		n 212		n 158		n 859		n 264	
Inactive	7	3.3	12	2.0	17	3.7	16	4.1	16	1.9	15	2.4
Moderately active	12	4.9	11	2.2	9	2.4	12	4.3	14	1.6	9	2.4
Active	81	5.6	77	3.0	74	4.2	72	5.4	70	2.3	75	3.2

ENSANUT, National Health and Nutrition Survey, MxDQI, Mexican Diet Quality Index, MxAHEI, Mexican Alternate Healthy Eating Index.

Data for age, BMI, waist circumference, total MxDQI score and total MxAHEI score are presented as mean and SE; all other data are presented as % and SE.

*They occupied the property in another situation not specified.

Table 3 Association of Mexican Diet Quality Index (MxDQI) and Mexican Alternate Healthy Eating Index (MxAHEI) with BMI in men and women. ENSANUT 2012 (n 2310)

	Men (n 954)			Women (n 1356)		
	β	95 % CI	P value	β	95 % CI	P value
MxDQI						
Total	-0.02	-0.05, 0.003	0.09	-0.02	-0.04, 0.01	0.23
Dietary components*						
Vegetables	-0.14	-0.25, -0.02	0.02	0.04	-0.08, 0.16	0.50
Whole fruit	0.12	-0.04, 0.27	0.13	0.05	-0.07, 0.17	0.44
Whole-grain cereals	-0.07	-0.28, 0.14	0.50	-0.04	-0.27, 0.18	0.70
Legumes	-0.05	-0.15, 0.05	0.32	-0.09	-0.20, 0.02	0.12
Seafood, poultry or eggs	-0.24	-0.42, -0.06	0.01	-0.17	-0.36, 0.02	0.08
Low-fat dairy	-0.08	-0.46, 0.31	0.69	0.38	-0.10, 0.85	0.12
Polyunsaturated fat	-0.01	-0.10, 0.09	0.89	0.08	-0.02, 0.17	0.11
100% Fruit juices	0.01	-0.25, 0.28	0.92	0.07	-0.21, 0.34	0.65
Refined grains	0.09	-0.12, 0.31	0.40	0.04	-0.17, 0.24	0.74
Red and processed meat	-0.14	-0.31, 0.03	0.10	-0.16	-0.36, 0.04	0.11
Added sugars	-0.01	-0.07, 0.05	0.81	-0.004	-0.07, 0.06	0.88
Sodium	-0.04	-0.12, 0.04	0.32	-0.02	-0.09, 0.05	0.60
Saturated fat	0.08	-0.10, 0.27	0.37	-0.12	-0.34, 0.09	0.25
MxAHEI						
Total	-0.04	-0.08, 0.004	0.08	-0.03	-0.07, 0.01	0.15
Dietary components*						
Vegetables	-0.12	-0.28, 0.05	0.17	0.09	-0.07, 0.25	0.29
Whole fruit	0.17	-0.04, 0.38	0.11	0.05	-0.10, 0.20	0.51
Whole-grain cereals	-0.04	-0.14, 0.07	0.51	-0.02	-0.13, 0.09	0.69
Legumes	0.04	-0.14, 0.22	0.66	-0.19	-0.38, 0.01	0.06
Nuts	0.47	0.07, 0.87	0.02	0.20	-0.68, 1.07	0.66
Polyunsaturated fat	-0.12	-0.27, 0.03	0.11	0.11	-0.02, 0.23	0.09
Long-chain (n-3) fats	0.01	-0.12, 0.15	0.87	-0.01	-0.17, 0.15	0.92
Sugar-sweetened beverages	-0.04	-0.15, 0.06	0.40	-0.03	-0.14, 0.07	0.55
Red and processed meat	-0.10	-0.19, -0.003	0.04	-0.08	-0.18, 0.01	0.08
Sodium	0.03	-0.09, 0.14	0.67	-0.06	-0.16, 0.04	0.23
Trans fat	0.26	-0.81, 1.33	0.63	-0.41	-1.82, 1.00	0.57
Alcohol	0.01	-0.12, 0.15	0.84	1.45	0.55, 2.35	< 0.01

ENSANUT, National Health and Nutrition Survey.

*Adjusted for the other dietary components.

Table 4 Association of total Mexican Diet Quality Index (MxDQI) and Mexican Alternate Healthy Eating Index (MxAHEI) with waist circumference in men and women. ENSANUT 2012 (*n* 2310)

	Men (<i>n</i> 954)			Women (<i>n</i> 1356)		
	β	95 % CI	<i>P</i> value	β	95 % CI	<i>P</i> value
MxDQI						
Total	-0.10	-0.20, -0.004	0.04	-0.02	-0.08, 0.04	0.50
Dietary components*						
Vegetables	-0.40	-0.82, 0.02	0.07	0.12	-0.16, 0.40	0.39
Whole fruit	0.40	-0.39, 1.19	0.32	0.17	-0.13, 0.46	0.27
Whole-grain cereals	-0.43	-1.20, 0.35	0.28	0.13	-0.38, 0.65	0.62
Legumes	-0.14	-0.47, 0.18	0.39	-0.13	-0.39, 0.12	0.31
Seafood, poultry or eggs	-0.88	-1.60, -0.17	0.02	-0.30	-0.71, 0.10	0.14
Low-fat dairy	-0.45	-1.64, 0.75	0.46	0.78	-0.17, 1.74	0.11
Polyunsaturated fat	-0.07	-0.46, 0.31	0.71	0.12	-0.12, 0.35	0.33
100% Fruit juices	0.25	-0.58, 1.07	0.55	0.12	-0.47, 0.71	0.69
Refined grains	0.23	-0.58, 1.03	0.58	0.14	-0.33, 0.60	0.56
Red and processed meat	-0.54	-1.06, -0.03	0.04	-0.32	-0.81, 0.17	0.19
Added sugars	-0.05	-0.28, 0.18	0.66	-0.02	-0.16, 0.13	0.82
Sodium	-0.18	-0.49, 0.14	0.27	-0.04	-0.21, 0.12	0.61
Saturated fat	0.49	-0.06, 1.03	0.08	-0.40	-0.89, 0.10	0.12
MxAHEI						
Total	-0.15	-0.28, -0.01	0.03	-0.07	-0.16, 0.02	0.13
Dietary components*						
Vegetables	-0.25	-1.07, 0.58	0.56	0.12	-0.24, 0.49	0.51
Whole fruit	0.64	-0.49, 1.76	0.27	0.19	-0.21, 0.59	0.34
Whole-grain cereals	-0.25	-0.61, 0.11	0.17	0.05	-0.20, 0.30	0.69
Legumes	0.09	-0.61, 0.78	0.81	-0.35	-0.80, 0.09	0.12
Nuts	0.64	-0.53, 1.81	0.28	0.59	-1.54, 2.71	0.59
Polyunsaturated fat	-0.47	-1.17, 0.23	0.19	0.20	-0.13, 0.53	0.24
Long-chain (<i>n</i> -3) fats	-0.06	-0.57, 0.46	0.83	-0.18	-0.59, 0.23	0.39
Sugar-sweetened beverages	-0.13	-0.58, 0.33	0.58	-0.15	-0.37, 0.07	0.19
Red and processed meat	-0.36	-0.67, -0.04	0.03	-0.15	-0.40, 0.10	0.24
Sodium	0.20	-0.20, 0.60	0.33	-0.17	-0.41, 0.07	0.17
Trans fat	0.57	-2.64, 3.78	0.73	-0.76	-4.37, 2.84	0.68
Alcohol	-0.06	-0.48, 0.35	0.77	0.95	0.24, 1.67	0.01

ENSANUT, National Health and Nutrition Survey.

*Adjusted for the other dietary components.

Association of MxDQI and MxAHEI with waist circumference in men and women

Among men, a one-unit increase in total MxDQI and total MxAHEI scores was associated with a reduction in WC of 0.10 (95 % CI 0.004, 0.20) cm and 0.15 (95 % CI 0.01, 0.28) cm, respectively. With regard to specific components, we observed that seafood, poultry or eggs as well as red and processed meat components were inversely associated with WC. In women, dietary scores were not associated with WC (Table 4).

Association of MxDQI and MxAHEI with BMI by education level in men and women

We found that the association of total MxDQI and total MxAHEI scores with BMI was different by education level in men (*P* for interaction < 0.05). A one-unit increase in total MxDQI and total MxAHEI scores was associated with a mean reduction in BMI of 0.11 (95 % CI 0.04, 0.18) kg/m² and 0.18 (95 % CI 0.10, 0.25) kg/m², respectively, for men with the lowest level of education. There was no association between the dietary scores and BMI for higher-educated men. With regard to components, we observed that the legumes and refined grains components were

inversely associated with BMI in men with lowest education level (*P* for interaction < 0.05; Table 5).

In women, the association of total MxDQI score with BMI was not different by education level. However, vegetables, whole fruit, as well as seafood, poultry or eggs components were inversely associated with BMI in women with no reading/writing skills (*P* for interaction < 0.05). Likewise, the association between total MxAHEI score and BMI was not different by education level. Nevertheless, the whole fruit component was inversely associated with BMI among women with the lowest education level, and the long-chain (*n*-3) fats component was inversely associated with BMI in women with ≥ 10 years of school (Table 5).

Association of MxDQI and MxAHEI with waist circumference by education level in men and women

As observed for BMI, the association between total MxDQI score and WC was different by education level in men (*P* for interaction < 0.05). The total MxDQI score was inversely associated with BMI in men with the lowest and highest education level (*P* < 0.05). Among specific components,

Table 5 Association of Mexican Diet Quality Index (MxDQI) and Mexican Alternate Healthy Eating Index (MxAHEI) with BMI in men and women by education level. ENSANUT 2012 (*n* 2310)

	Education level									<i>P</i> value for interaction*
	No reading/writing skills			Reading/writing skills or 3–9 years of school			≥10 years of school			
	β	95 % CI	<i>P</i> value	β	95 % CI	<i>P</i> value	β	95 % CI	<i>P</i> value	
Men										
<i>n</i>	100			621			233			
MxDQI score										
Total	-0.11	-0.18, -0.04	< 0.01	-0.01	-0.04, 0.03	0.70	-0.04	-0.08, 0.01	0.10	0.02
Dietary components†										
Vegetables	-0.17	-0.47, 0.13	0.27	-0.17	-0.31, -0.03	0.02	-0.07	-0.29, 0.16	0.56	0.74
Whole fruit	-0.05	-0.43, 0.32	0.78	0.13	-0.03, 0.29	0.11	0.13	-0.17, 0.43	0.41	0.66
Whole-grain cereals	-0.29	-0.76, 0.18	0.22	-0.005	-0.26, 0.25	0.97	-0.16	-0.48, 0.15	0.31	0.48
Legumes	-0.34	-0.55, -0.12	< 0.01	0.01	-0.12, 0.13	0.91	-0.07	-0.26, 0.13	0.51	0.02
Seafood, poultry or eggs	-0.23	-0.67, 0.20	0.29	-0.12	-0.34, 0.09	0.26	-0.47	-0.82, -0.12	0.01	0.24
Low-fat dairy	-0.16	-2.05, 1.73	0.87	-0.08	-0.55, 0.39	0.74	-0.07	-0.67, 0.52	0.81	1.00
Polyunsaturated fat	-0.15	-0.37, 0.08	0.20	-0.003	-0.10, 0.10	0.95	0.03	-0.18, 0.24	0.78	0.44
100 % Fruit juices	-0.66	-1.41, 0.08	0.08	-0.10	-0.45, 0.26	0.60	0.20	-0.12, 0.51	0.22	0.08
Refined grains	-0.48	-0.91, -0.04	0.03	0.18	-0.07, 0.44	0.16	0.08	-0.28, 0.43	0.67	0.02
Red and processed meat	-0.63	-1.04, -0.21	< 0.01	-0.12	-0.34, 0.11	0.30	-0.09	-0.44, 0.26	0.62	0.07
Added sugars	-0.10	-0.23, 0.04	0.16	0.002	-0.08, 0.08	0.97	0.003	-0.1, 0.11	0.94	0.42
Sodium	-0.05	-0.20, 0.10	0.50	-0.003	-0.09, 0.09	0.95	-0.13	-0.28, 0.01	0.08	0.27
Saturated fat	-0.15	-0.61, 0.32	0.54	0.10	-0.12, 0.33	0.37	0.12	-0.24, 0.48	0.53	0.60
MxAHEI score										
Total	-0.18	-0.25, -0.10	< 0.01	-0.02	-0.07, 0.03	0.38	-0.04	-0.11, 0.04	0.33	< 0.01
Dietary components†										
Vegetables	-0.24	-0.63, 0.16	0.24	-0.18	-0.36, -0.002	0.05	0.03	-0.45, 0.51	0.90	0.65
Whole fruit	-0.25	-0.62, 0.12	0.19	0.15	-0.07, 0.37	0.18	0.29	-0.21, 0.78	0.25	0.12
Whole-grain cereals	-0.19	-0.44, 0.06	0.14	-0.01	-0.16, 0.14	0.89	-0.12	-0.31, 0.06	0.18	0.37
Legumes	-0.39	-0.79, 0.01	0.06	0.14	-0.09, 0.37	0.24	-0.07	-0.41, 0.27	0.68	0.06
Nuts	0.27	-0.36, 0.90	0.40	0.53	-0.15, 1.21	0.12	0.44	0.07, 0.81	0.02	0.84
Polyunsaturated fat	-0.29	-0.64, 0.05	0.10	-0.05	-0.22, 0.12	0.59	-0.25	-0.65, 0.14	0.21	0.31
Long-chain (<i>n</i> -3) fats	-0.13	-0.47, 0.21	0.46	0.07	-0.11, 0.25	0.44	-0.05	-0.36, 0.26	0.75	0.52
Sugar-sweetened beverages	0.18	-0.40, 0.75	0.55	0.19	-0.08, 0.47	0.17	0.04	-0.25, 0.33	0.78	0.71
Red and processed meat	-0.33	-0.56, -0.10	0.01	-0.11	-0.24, 0.02	0.11	-0.07	-0.27, 0.14	0.51	0.17
Sodium	0.03	-0.21, 0.28	0.78	0.06	-0.10, 0.22	0.43	-0.07	-0.27, 0.13	0.49	0.56
<i>Trans</i> fat	13.64	7.64, 19.63	< 0.01	0.40	-0.82, 1.61	0.52	0.18	-1.61, 1.96	0.85	< 0.01
Alcohol	-0.22	-0.52, 0.08	0.15	0.01	-0.24, 0.25	0.94	0.06	-0.12, 0.24	0.50	0.29



Table 5 (Continued)

	Education level									P value for interaction*
	No reading/writing skills			Reading/writing skills or 3–9 years of school			≥10 years of school			
	β	95 % CI	P value	β	95 % CI	P value	β	95 % CI	P value	
Women										
<i>n</i>	165			910			281			
MxDQI score										
Total	−0.06	−0.14, 0.03	0.18	−0.002	−0.03, 0.03	0.89	−0.04	−0.10, 0.02	0.24	0.35
Dietary components†										
Vegetables	−0.36	−0.69, −0.03	0.04	0.11	−0.03, 0.25	0.13	0.03	−0.20, 0.26	0.79	0.03
Whole fruit	−0.35	−0.68, −0.02	0.04	0.06	−0.10, 0.21	0.46	0.16	−0.04, 0.35	0.11	0.03
Whole-grain cereals	−0.67	−1.35, 0.01	0.05	−0.01	−0.25, 0.23	0.93	−0.01	−0.42, 0.40	0.96	0.17
Legumes	−0.17	−0.52, 0.18	0.34	−0.03	−0.16, 0.10	0.61	−0.23	−0.45, −0.01	0.04	0.29
Seafood, poultry or eggs	−0.59	−1.26, 0.09	0.09	0.02	−0.20, 0.25	0.84	−0.52	−0.84, −0.19	< 0.01	0.01
Low-fat dairy	0.79	−0.04, 1.62	0.06	0.06	−0.35, 0.48	0.76	0.67	−0.49, 1.84	0.26	0.24
Polyunsaturated fat	0.08	−0.28, 0.44	0.65	0.13	0.01, 0.24	0.03	−0.03	−0.22, 0.15	0.72	0.34
100 % Fruit juices	−0.31	−1.40, 0.78	0.57	0.21	−0.03, 0.44	0.08	−0.19	−0.86, 0.48	0.58	0.35
Refined grains	−0.10	−0.80, 0.60	0.78	0.07	−0.18, 0.33	0.57	−0.02	−0.39, 0.35	0.92	0.85
Red and processed meat	−0.05	−1.03, 0.94	0.93	−0.10	−0.32, 0.12	0.38	−0.34	−0.64, −0.05	0.02	0.33
Added sugars	0.07	−0.17, 0.31	0.58	−0.04	−0.11, 0.04	0.31	0.05	−0.06, 0.17	0.37	0.32
Sodium	0.02	−0.20, 0.25	0.84	−0.02	−0.10, 0.06	0.60	−0.03	−0.16, 0.10	0.64	0.92
Saturated fat	−0.28	−1.09, 0.54	0.50	−0.13	−0.36, 0.10	0.28	−0.04	−0.38, 0.30	0.82	0.83
MxAHEI score										
Total	−0.03	−0.19, 0.13	0.69	0.001	−0.05, 0.05	0.97	−0.11	−0.20, −0.02	0.02	0.10
Dietary components†										
Vegetables	−0.54	−1.08, 0.01	0.06	0.13	−0.08, 0.34	0.22	0.19	−0.13, 0.51	0.24	0.06
Whole fruit	−0.60	−1.12, −0.07	0.03	0.04	−0.18, 0.26	0.73	0.25	−0.02, 0.51	0.07	0.01
Whole-grain cereals	−0.37	−0.76, 0.02	0.07	−0.04	−0.17, 0.09	0.55	−0.03	−0.27, 0.20	0.79	0.27
Legumes	−0.38	−1.08, 0.31	0.28	−0.04	−0.27, 0.19	0.72	−0.46	−0.82, −0.10	0.01	0.11
Nuts	−0.84	−2.38, 0.71	0.29	0.64	−0.66, 1.94	0.33	−0.23	−1.16, 0.70	0.63	0.33
Polyunsaturated fat	0.21	−0.36, 0.77	0.47	0.18	0.01, 0.35	0.04	−0.004	−0.26, 0.25	0.97	0.49
Long-chain (<i>n</i> -3) fats	0.21	−0.55, 0.97	0.59	0.20	−0.02, 0.42	0.08	−0.35	−0.56, −0.14	< 0.01	< 0.01
Sugar-sweetened beverages	0.62	−0.85, 2.08	0.41	0.16	−0.11, 0.44	0.25	0.17	−0.17, 0.51	0.33	0.84
Red and processed meat	−0.05	−0.60, 0.51	0.87	−0.05	−0.16, 0.07	0.42	−0.22	−0.38, −0.05	0.01	0.19
Sodium	0.14	−0.25, 0.54	0.48	−0.08	−0.20, 0.05	0.24	−0.15	−0.33, 0.03	0.10	0.40
Trans fat	−4.37	−13.99, 5.25	0.37	−0.82	−2.66, 1.03	0.39	0.75	−1.54, 3.04	0.52	0.40
Alcohol‡	–	–	–	0.84	0.23, 1.45	0.01	2.47	1.91, 3.03	< 0.01	< 0.01

ENSANUT, National Health and Nutrition Survey.

*The *P* value for interaction represents the statistical significance of the interaction term to test whether the association between diet quality and BMI is different for low- v. high-educated individuals.

†Adjusted for the other dietary components.

‡Blank cells indicate variable was omitted from model.



legumes as well as red and processed meat MxDQI components were inversely associated with WC among men with no reading/writing skills (P for interaction < 0.05). Likewise, the association between total MxAHEI and WC was different by education level (P for interaction < 0.05). The total MxAHEI score was inversely associated with WC in men with no reading/writing skills. The red and processed meat MxAHEI component was inversely associated with WC in men with lowest education level (P for interaction < 0.05 ; Table 6).

The association between total MxDQI score and WC in women was not different by education level. However, the whole fruit component was inversely associated with WC among women with no reading/writing skills, whereas the seafood, poultry or eggs component was inversely associated with WC among women with ≥ 10 years of school (P for interaction < 0.05). Likewise, we observed that whole fruit and nuts MxAHEI components were inversely associated with WC in women with the lowest education level, while the long-chain ($n-3$) fats component was inversely associated with WC among women with ≥ 10 years of school (Table 6).

Results of sensitivity analyses

Estimated associations of total MxDQI and MxAHEI scores with BMI and WC were similar to those observed when total MxDQI and MxAHEI models were further adjusted for physical activity. However, when physical activity was included as covariate, the associations between dietary scores and WC were no longer statistically significant in men, whereas in women these associations were statistically significant in the MxAHEI models (see online supplementary material, Supplemental Table 2). Likewise, estimated associations of total MxDQI and MxAHEI scores with BMI and WC were similar to those observed when corn tortilla was considered as refined instead of whole grain (Supplemental Table 2). The whole-grain cereals MxDQI component was positively associated with BMI in men, whereas the refined grain MxDQI component was positively associated with BMI in women when corn tortilla was considered as refined grain (Supplemental Table 3). Supplemental Tables 4–7 present the results of testing the association of diet quality scores with BMI and WC without adjusting for the other dietary components and adjusting for total scores minus the dietary component of interest, which were similar to those obtained using models adjusted for the other dietary components.

Discussion

To our best knowledge, the present study is the first that examines the association of diet quality with BMI and WC using two different dietary quality scores, one based on the most recent MxDG and the other on foods and

nutrients associated with cardiometabolic risk. We found that both total dietary quality scores were not associated with BMI in men and women. However, some dietary components were inversely associated with BMI, including the vegetables and seafood, poultry or eggs MxDQI components, as well as the red and processed meat MxAHEI component in men. We also observed that the total dietary scores were inversely associated with WC in men, but not in women. We specifically found that the seafood, poultry or eggs as well as red and processed meat components were inversely associated with WC in men.

The present study is also one of the first to examine the association of diet quality with BMI and WC by education level, which we considered relevant as a first exploration given the disparities in dietary intakes and obesity prevalence by socio-economic status already documented in Mexican adults. We found that for men with the least education, total high diet quality was associated with lower BMI and WC. In contrast, there was no association between total diet quality and BMI or WC in higher-educated men.

It is not clear from the current study why better total diet quality predicts lower BMI and WC for lower-educated but not higher-educated men. Several studies have found that individuals with lower socio-economic status more frequently undertake behaviours that could be detrimental to health than those of higher socio-economic status, such as low physical activity, smoking or poor sleep duration^(49,50). Thus, one possibility is that improvements in diet quality offset the association of these other behaviours with health in low-educated men. A second possibility might be that for low-educated men, diet quality is a proxy for better health behaviour overall; men who seek out healthy diets despite the norm may also be more likely to engage in other health-promoting behaviours, and this combined behaviour leads to better weight outcomes. A third possibility relates more specifically to the role of physical activity: men with less education may have jobs that are more physically demanding in comparison to higher-educated men; higher diet quality scores together with higher work-related physical activity could be associated with a reduction of BMI and WC. Previous studies in low- and middle-income countries have found that individuals with less education were more active than groups with more education⁽⁵¹⁾ and occupational activity has been found to be one of the major contributors to the intensity of overall physical activity⁽⁵²⁾. Unfortunately, we were unable to include physical activity in the main analyses due to the poor validity of the International Physical Activity Questionnaire short form in Mexican adults⁽⁴⁸⁾, so we were able to examine only crude physical activity in our models. More research is needed to understand how diet interacts with other key lifestyle components, such as physical activity, sedentary activity, occupation and sleep duration, to understand the pathway between diet and obesity across the population.



Table 6 Association of Mexican Diet Quality Index (MxDQI) and Mexican Alternate Healthy Eating Index (MxAHEI) with waist circumference in men and women by education level. ENSANUT 2012 (n 2310)

	Education level									P value for interaction*
	No reading/writing skills			Reading/writing skills or 3–9 years of school			≥10 years of school			
	β	95 % CI	P value	β	95 % CI	P value	β	95 % CI	P value	
Men										
n	100			621			233			
MxDQI score										
Total	-0.30	-0.49, -0.11	< 0.01	-0.03	-0.16, 0.1	0.66	-0.20	-0.35, -0.05	0.01	0.03
Dietary components†										
Vegetables	-0.46	-1.37, 0.44	0.32	-0.73	-1.18, -0.28	< 0.01	0.32	-0.73, 1.37	0.55	0.21
Whole fruit	-0.18	-1.13, 0.77	0.71	0.28	-0.25, 0.81	0.31	0.74	-0.98, 2.45	0.40	0.52
Whole-grain cereals	-0.79	-2.50, 0.92	0.37	-0.25	-1.12, 0.62	0.57	-0.69	-2.08, 0.70	0.33	0.80
Legumes	-0.91	-1.56, -0.26	0.01	0.06	-0.35, 0.47	0.77	-0.32	-1.17, 0.53	0.46	0.03
Seafood, poultry or eggs	-0.86	-2.20, 0.49	0.21	-0.45	-1.22, 0.32	0.26	-1.76	-3.45, -0.06	0.04	0.35
Low-fat dairy	0.55	-7.57, 8.67	0.89	-0.01	-1.35, 1.33	0.99	-0.96	-2.76, 0.83	0.29	0.67
Polyunsaturated fat	-0.58	-1.22, 0.05	0.07	0.05	-0.30, 0.39	0.79	-0.18	-1.21, 0.85	0.73	0.18
100 % Fruit juices	-3.99	-8.07, 0.09	0.06	-0.39	-1.05, 1.12	0.95	0.69	-0.33, 1.71	0.18	0.08
Refined grains	-0.85	-2.28, 0.58	0.24	0.35	-0.49, 1.20	0.41	0.34	-1.26, 1.93	0.68	0.28
Red and processed meat	-2.01	-3.30, -0.71	< 0.01	-0.09	-0.76, 0.57	0.78	-1.04	-2.66, 0.57	0.21	0.01
Added sugars	-0.33	-0.75, 0.10	0.13	-0.001	-0.31, 0.30	0.99	-0.07	-0.45, 0.32	0.72	0.46
Sodium	-0.05	-0.46, 0.37	0.83	-0.003	-0.32, 0.31	0.98	-0.70	-1.36, -0.04	0.04	0.09
Saturated fat	0.07	-1.46, 1.59	0.93	0.75	0.05, 1.44	0.04	-0.27	-1.69, 1.15	0.71	0.43
MxAHEI score										
Total	-0.53	-0.75, -0.30	< 0.01	-0.08	-0.25, 0.09	0.37	-0.22	-0.46, 0.02	0.07	0.01
Dietary components†										
Vegetables	-0.53	-1.66, 0.61	0.37	-0.81	-1.40, -0.22	0.01	1.03	-1.82, 3.87	0.48	0.47
Whole fruit	-0.86	-1.98, 0.26	0.13	0.26	-0.43, 0.94	0.46	1.64	-1.38, 4.66	0.29	0.12
Whole-grain cereals	-0.54	-1.47, 0.39	0.26	-0.04	-0.47, 0.38	0.84	-0.68	-1.47, 0.12	0.10	0.26
Legumes	-1.12	-2.34, 0.11	0.07	0.48	-0.38, 1.34	0.27	-0.50	-1.98, 0.97	0.50	0.04
Nuts	1.09	-0.97, 3.15	0.30	0.83	-0.90, 2.56	0.35	0.44	-0.98, 1.86	0.54	0.85
Polyunsaturated fat	-1.01	-2.02, -0.01	0.05	-0.09	-0.81, 0.63	0.81	-1.26	-3.48, 0.95	0.26	0.17
Long-chain (n-3) fats	-0.90	-1.83, 0.03	0.06	0.34	-0.35, 1.03	0.34	-0.64	-1.70, 0.43	0.24	0.06
Sugar-sweetened beverages	1.07	-0.40, 2.54	0.16	0.22	-0.52, 0.97	0.55	-0.48	-1.56, 0.59	0.38	0.15
Red and processed meat	-1.14	-1.83, -0.45	< 0.01	-0.23	-0.66, 0.20	0.30	-0.55	-1.44, 0.34	0.22	0.06
Sodium	0.40	-0.29, 1.08	0.25	0.26	-0.30, 0.83	0.36	0.02	-0.94, 0.97	0.97	0.81
Trans fat	42.06	23.61, 60.51	< 0.01	1.94	-2.11, 5.99	0.35	-0.28	-5.20, 4.65	0.91	< 0.01
Alcohol	-0.76	-1.93, 0.41	0.21	-0.12	-0.84, 0.61	0.75	0.14	-0.43, 0.71	0.63	0.39

Table 6 (Continued)

	Education level									P value for interaction*
	No reading/writing skills			Reading/writing skills or 3–9 years of school			≥10 years of school			
	β	95 % CI	P value	β	95 % CI	P value	β	95 % CI	P value	
Women										
<i>n</i>		165			910			281		
MxDQI score										
Total	−0.08	−0.25, 0.09	0.36	−0.003	−0.07, 0.07	0.92	−0.04	−0.18, 0.10	0.58	0.67
Dietary components†										
Vegetables	−0.76	−1.51, −0.001	0.05	0.23	−0.11, 0.56	0.18	0.21	−0.33, 0.74	0.45	0.05
Whole fruit	−0.87	−1.53, −0.21	0.01	0.22	−0.15, 0.58	0.24	0.41	−0.11, 0.92	0.12	0.01
Whole-grain cereals	−1.55	−3.09, −0.02	0.05	0.04	−0.54, 0.62	0.89	0.54	−0.38, 1.46	0.25	0.06
Legumes	−0.10	−0.85, 0.64	0.79	−0.03	−0.33, 0.26	0.83	−0.50	−1.08, 0.07	0.08	0.34
Seafood, poultry or eggs	−1.08	−2.50, 0.33	0.13	0.10	−0.40, 0.61	0.69	−1.08	−1.85, −0.30	0.01	0.01
Low-fat dairy	2.84	1.06, 4.61	< 0.01	0.18	−0.88, 1.25	0.73	0.92	−1.05, 2.89	0.36	0.03
Polyunsaturated fat	0.14	−0.57, 0.85	0.70	0.15	−0.11, 0.42	0.27	0.03	−0.43, 0.48	0.90	0.89
100 % Fruit juices	−1.50	−3.87, 0.87	0.22	0.48	−0.21, 1.16	0.17	−0.40	−1.45, 0.66	0.46	0.15
Refined grains	0.50	−1.02, 2.02	0.52	0.05	−0.51, 0.61	0.87	0.25	−0.60, 1.09	0.57	0.82
Red and processed meat	0.09	−2.10, 2.29	0.93	−0.07	−0.59, 0.45	0.79	−1.03	−1.83, −0.23	0.01	0.08
Added sugars	0.05	−0.42, 0.52	0.83	−0.09	−0.26, 0.09	0.32	0.14	−0.15, 0.43	0.35	0.36
Sodium	0.11	−0.44, 0.66	0.69	−0.05	−0.25, 0.15	0.66	−0.11	−0.38, 0.16	0.43	0.79
Saturated fat	−0.35	−2.03, 1.33	0.68	−0.47	−1.00, 0.06	0.08	−0.17	−1.13, 0.79	0.73	0.83
MxAHEI score										
Total	−0.11	−0.39, 0.17	0.46	0.003	−0.11, 0.11	0.95	−0.24	−0.41, −0.06	0.01	0.06
Dietary components†										
Vegetables	−1.24	−2.50, 0.03	0.06	0.16	−0.31, 0.64	0.50	0.47	−0.27, 1.20	0.21	0.07
Whole fruit	−1.48	−2.59, −0.37	0.01	0.26	−0.31, 0.83	0.38	0.55	−0.18, 1.28	0.14	0.01
Whole-grain cereals	−0.91	−1.78, −0.04	0.04	0.02	−0.30, 0.35	0.89	0.24	−0.27, 0.75	0.35	0.06
Legumes	−0.51	−2.01, 0.98	0.50	−0.12	−0.65, 0.42	0.67	−0.96	−1.84, −0.09	0.03	0.23
Nuts	−4.15	−7.57, −0.73	0.02	2.05	−1.34, 5.44	0.24	−0.69	−2.09, 0.72	0.34	0.04
Polyunsaturated fat	0.48	−0.66, 1.62	0.41	0.26	−0.15, 0.68	0.21	0.06	−0.64, 0.76	0.87	0.78
Long-chain (<i>n</i> -3) fats	0.26	−1.52, 2.04	0.77	0.32	−0.17, 0.81	0.20	−1.10	−1.8, −0.40	< 0.01	< 0.01
Sugar-sweetened beverages	0.06	−2.15, 2.28	0.96	0.002	−0.59, 0.60	1.00	0.18	−0.63, 1.00	0.66	0.94
Red and processed meat	0.03	−1.22, 1.28	0.96	−0.01	−0.30, 0.28	0.94	−0.59	−1.05, −0.13	0.01	0.06
Sodium	0.31	−0.55, 1.16	0.48	−0.19	−0.49, 0.10	0.19	−0.36	−0.80, 0.08	0.11	0.39
Trans fat	−12.4	−33.02, 8.22	0.24	−2.60	−6.68, 1.48	0.21	5.11	−4.23, 14.45	0.28	0.17
Alcohol‡	–	–	–	0.70	−0.04, 1.44	0.06	1.46	0.44, 2.48	0.01	0.21

ENSANUT, National Health and Nutrition Survey.

*The P value for interaction represents the statistical significance of the interaction term to test whether the association between diet quality and waist circumference is different for low- v. high-educated individuals.

†Adjusted for the other dietary components.

‡Blank cells indicate variable was omitted from model.



In our primary models, the association of total MxDQI and MxAHEI scores with BMI and WC was not different by education level in women, but the pattern of the associations tended to be the opposite than in men. We found a statistically significant association of total MxAHEI score with BMI and WC in women with ≥ 10 years of education. Although associations were not statistically significant in women with lower education levels, BMI and WC estimations were in the same direction as those observed in higher-educated women. Sex differences in associations between risk factors and obesity have been reported elsewhere, and several explanations have been proposed given the complexity of the subject matter, including differences in the duration and intensity of occupational physical activity. Studies suggest that higher-educated women could have higher physical activity levels than lower-educated women, which in turn can be associated with lower BMI or WC if women improve the quality of their diet^(53,54). One of the potential reasons why we did not find different results by education may be because lower-educated women could also have levels of housework-related physical activity that interact with diet to reduce BMI and WC, but not enough to find statistically significant associations. However, the association between total MxAHEI score and WC differed by education in women after adjusting for physical activity. Future research using valid measurements of physical activity will be needed to understand how diet might interact with physical activity in women.

We specifically observed that the association of different dietary components with BMI or WC was different by education level in men and women. Higher consumption of legumes and lower intakes of refined grains and red and processed meat were inversely associated with BMI or WC among men with the lowest education level, whereas among women in the same level of education as men, a higher consumption of fruit and vegetables was associated with low BMI or WC. A potential explanation of these results is the heterogeneity in the consumption of dietary components among men and women. Given the truncated nature of the diet quality indices, it is possible that, for instance, the consumption of fruit and vegetables was even higher in women than in men, and therefore associated with lower BMI or WC, even though the proportion of individuals who met the scoring criteria of each one of the dietary components was similar between men and women by education level. Another possible reason for the differences by sex is that women and men completed the 24 h dietary recall differently, potentially resulting in different measurement error. Furthermore, we found that different dietary components were associated with BMI and WC. For instance, red and processed meat, in both MxDQI and MxAHEI, was consistently associated with WC but not with BMI. Similar results were observed in a previous study in Iranian adults⁽⁵⁵⁾. Therefore, it is possible that the WC measure is more sensitive to the changes in

some dietary factors, compared with other measures of general obesity such as BMI.

Limitations

It is important to acknowledge the limitations of our study. First, the potential reverse causal link between diet quality and obesity cannot be dismissed, given the nature of the cross-sectional study design. Second, the current study was based on a single 24 h recall, which does not allow between- and within-person variability to be distinguished, and therefore usual intake cannot be estimated. Specifically, episodically consumed dietary components, such as alcohol, could be misrepresented and this might be why we were unable to find consistent associations of dietary quality with BMI and WC. Despite the limitation of using this method, some cross-sectional studies have found associations between diet quality and health outcomes using a single 24 h recall^(12,56,57), which may indicate that some food components and weights were not the most adequate to estimate the diet quality in Mexican adults. For instance, it could be useful to categorize tortillas as cooked and fried, since both types of tortilla can be highly consumed but their associations with BMI and WC may be opposite. Therefore, future direction should include the modifications of these indices or the creation of other indices that better reflect changes in BMI and WC in Mexican adults. The FFQ could provide a better estimation of usual intakes and therefore of usual diet quality. However, the FFQ used in ENSANUT was too aggregated and would result in misclassification because many details of dietary intake were not measured, and the quantification of intake was not as accurate as with 24 h dietary recalls. Recent studies have shown that the combined use of multiple 24 h dietary recalls and FFQ provides data superior to use of either method alone^(58–60), but this was not possible for the present study since these two dietary methods were collected in different sub-samples. Therefore, we recommend the collection in the future of both dietary instruments in the same sub-sample for a better estimation of the usual diet quality index. Finally, we selected education level as the measure of socio-economic status, rather than occupation or income. Education level can determine the occupation, and education level and occupation are jointly associated with income level⁽⁶¹⁾. However, these three proxies of socio-economic status could also have an independent role in predicting health-related behaviours. We considered it inappropriate to analyse occupation because of the complexity of understanding the potential sex differences, since 78 % of men reported they had paid work the week prior to the interview, whereas 62 % of women reported unpaid work in the same period. Moreover, income was measured using an open-ended question. Income is more difficult to estimate in low- to middle-income countries because of greater reliance on the informal economy and self-employment⁽⁶²⁾. However, we used

an asset-based household index as a control variable in our models. This index has been recommended by economists as a key proxy of household income, especially in low- to middle-income countries, including Mexico^(62,63). Despite the limitations, the results presented provide an overview of how diet quality is associated with BMI and WC and how these associations can be potentially different by education level in a representative sample of Mexican men and women, which offers insights for understanding diet–health outcome disparities in Mexico.

Conclusion

Results show total dietary quality scores were not associated with BMI in men and women, whereas the MxAHEI score was inversely with WC in men but not in women. Furthermore, the association of total MxDQI and MxAHEI scores with BMI and WC was different by education level in Mexican men. In Mexican women, the association of diet quality scores with BMI and WC did not differ by education. These results provide insights when implementing public actions that prevent or ameliorate the high prevalence of obesity in Mexican adults, considering the potential differences by sex and education.

Acknowledgements

Acknowledgements: The authors thank Dr Phil Bardsley for exceptional assistance with the data management, Ms Emily Ford Yoon and Ms Frances Dancy Burton for administrative assistance, and Ms Denise Ammons for graphics support. *Financial support:* This work was funded primarily by Bloomberg Philanthropies (grants to the University of North Carolina and Centro de Investigación en Nutrición y Salud, Instituto Nacional de Salud Pública), with support from the National Institutes of Health (NIH; grant number R01DK108148), the Robert Wood Johnson Foundation (grant number 71698), and the Carolina Population Center and its NIH Center grant (grant number P2C HD050924). N.L.-O. is supported by a Mexican National Council for Science and Technology (CONACyT) scholarship. Funders had no role in the design, analysis or writing of this article. *Conflict of interest:* None. *Authorship:* N.L.-O., B.M.P. and L.S.T. were responsible for study design, results interpretation and writing the manuscript. N.L.-O. was responsible for performing statistical analysis and drafting and editing of the manuscript. All authors contributed to study design and critical revision of the manuscript. All authors approved the final manuscript. *Ethics of human subject participation:* This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human subjects were approved by the Ethics Committee

of the Mexican National Institute of Public Health. Written informed consent was obtained from all subjects under study.

Supplementary material

To view supplementary material for this article, please visit <https://doi.org/10.1017/S136898001900065X>

References

1. Kroker-Lobos MF, Pedroza-Tobias A, Pedraza LS *et al.* (2014) The double burden of undernutrition and excess body weight in Mexico. *Am J Clin Nutr* **100**, issue 6, 1652S–1658S.
2. Gómez-Dantés H, Fullman N, Lamadrid-Figueroa H *et al.* (2016) Dissonant health transition in the states of Mexico, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* **388**, 2386–2402.
3. Hernández-Ávila M, Rivera-Domarco J, Shamah T *et al.* (2016) *Encuesta Nacional de Salud y Nutrición de Medio Camino 2016. Informe Final de Resultados (National Health and Nutrition Survey 2016. Final Results)*. Cuernavaca, Mexico: National Institute of Public Health.
4. Lim SS, Vos T, Flaxman AD *et al.* (2012) A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* **380**, 2224–2260.
5. Stern D, Smith LP, Zhang B *et al.* (2014) Changes in waist circumference relative to body mass index in Chinese adults, 1993–2009. *Int J Obes (Lond)* **38**, 1503–1510.
6. Aburto TC, Pedraza LS, Sanchez-Pimienta TG *et al.* (2016) Discretionary foods have a high contribution and fruit, vegetables, and legumes have a low contribution to the total energy intake of the Mexican population. *J Nutr* **146**, issue 9, 1881s–1887s.
7. Arvaniti F & Panagiotakos DB (2008) Healthy indexes in public health practice and research: a review. *Crit Rev Food Sci Nutr* **48**, 317–327.
8. Schwingshackl L & Hoffmann G (2015) Diet quality as assessed by the Healthy Eating Index, the Alternate Healthy Eating Index, the Dietary Approaches to Stop Hypertension score, and health outcomes: a systematic review and meta-analysis of cohort studies. *J Acad Nutr Diet* **115**, 780–800.e5.
9. Kourlaba G & Panagiotakos DB (2009) Dietary quality indices and human health: a review. *Maturitas* **62**, 1–8.
10. George SM, Ballard-Barbash R, Manson JE *et al.* (2014) Comparing indices of diet quality with chronic disease mortality risk in postmenopausal women in the Women's Health Initiative Observational Study: evidence to inform national dietary guidance. *Am J Epidemiol* **180**, 616–625.
11. Harmon BE, Boushey CJ, Shvetsov YB *et al.* (2015) Associations of key diet-quality indexes with mortality in the Multiethnic Cohort: the Dietary Patterns Methods Project. *Am J Clin Nutr* **101**, 587–597.
12. Sundararajan K, Campbell MK, Choi YH *et al.* (2014) The relationship between diet quality and adult obesity: evidence from Canada. *J Am Coll Nutr* **33**, 1–17.
13. Pate RR, Ross SET, Liese AD *et al.* (2015) Associations among physical activity, diet quality, and weight status in US adults. *Med Sci Sports Exerc* **47**, 743–750.
14. Asghari G, Mirmiran P, Yuzbashian E *et al.* (2017) A systematic review of diet quality indices in relation to obesity. *Br J Nutr* **117**, 1055–1065.



15. Gregory CO, McCullough ML, Ramirez-Zea M *et al.* (2009) Diet scores and cardio-metabolic risk factors among Guatemalan young adults. *Br J Nutr* **101**, 1805–1811.
16. López-Olmedo N, Carriquiry AL, Rodríguez-Ramírez S *et al.* (2016) Usual intake of added sugars and saturated fats is high while dietary fiber is low in the Mexican population. *J Nutr* **146**, issue 9, 1856S–1865S.
17. Pedroza-Tobias A & Rivera-Dommarco JA (2013) Prevalencia de obesidad en adultos Mexicanos, ENSANUT 2012 (Prevalence of obesity in Mexican adults, ENSANUT 2012). *Salud Publica Mex* **55**, Suppl. 2, S151–S160.
18. Arganini C, Saba A, Comitato R *et al.* (2012) Gender differences in food choice and dietary intake in modern western societies. In *Public Health – Social and Behavioral Health*, pp. 83–102 [J Maddock, editor]. Rijeka, Croatia: InTech.
19. Vogel C, Lewis D, Ntani G *et al.* (2017) The relationship between dietary quality and the local food environment differs according to level of educational attainment: a cross-sectional study. *PLoS One* **12**, e0183700.
20. Lovejoy JC & Sainsbury A (2009) Sex differences in obesity and the regulation of energy homeostasis. *Obes Rev* **10**, 154–167.
21. Palmer BF & Clegg DJ (2015) The sexual dimorphism of obesity. *Mol Cell Endocrinol* **402**, 113–119.
22. Whitaker KM, Choh AC, Lee M *et al.* (2016) Sex differences in the rate of abdominal adipose accrual during adulthood: the Fels Longitudinal Study. *Int J Obes (Lond)* **40**, 1278–1285.
23. Chiuvè SE, Fung TT, Rimm EB *et al.* (2012) Alternative dietary indices both strongly predict risk of chronic disease. *J Nutr* **142**, 1009–1018.
24. Gutiérrez JP, Rivera-Dommarco J, Shamah-Levy T *et al.* (2012) *Encuesta Nacional de Salud y Nutrición 2012. Resultados Nacionales*. Cuernavaca, México: Instituto Nacional de Salud Pública.
25. Conway JM, Ingwersen LA, Vinyard BT *et al.* (2003) Effectiveness of the US Department of Agriculture 5-step multiple-pass method in assessing food intake in obese and nonobese women. *Am J Clin Nutr* **77**, 1171–1178.
26. Blanton CA, Moshfegh AJ, Baer DJ *et al.* (2006) The USDA Automated Multiple-Pass Method accurately estimates group total energy and nutrient intake. *J Nutr* **136**, 2594–2599.
27. Agricultural Research Service, Food Surveys Research Group (2010) USDA Food and Nutrient Database for Dietary Studies, 4.1. <https://www.ars.usda.gov/northeast-area/beltsville-md/beltsville-human-nutrition-research-center/food-surveys-research-group/docs/fndds-download-databases/> (accessed January 2017).
28. Ledesma Solano J, Chávez Villasana A, Pérez-Gil Romo F *et al.* (2010) *Composición de Alimentos 'Miriam Muñoz de Chávez'. Valor Nutritivo de los Alimentos de Mayor Consumo (Food Composition 'Miriam Muñoz de Chávez'. Nutrition Value of the Most Consumed Foods)*, 2a ed. México, DF: McGraw-Hill.
29. US Department of Agriculture, Agricultural Research Service, Nutrient Data Laboratory (2013) National Nutrient Database for Standard Reference, Release 24. <https://www.ars.usda.gov/northeast-area/beltsville-md/beltsville-human-nutrition-research-center/nutrient-data-laboratory/docs/sr24-homepage/> (accessed January 2017).
30. Villalpando S, Ramírez-Silva I, Bernal Medina D *et al.* (2007) *Tablas de Composición de Ácidos Grasos de Alimentos Frecuentes en la Dieta Mexicana (Tables of Fatty Acids Composition of Common Foods in the Mexican Diet)*. Cuernavaca, Mexico: National Institute of Public Health.
31. World Health Organization (2008) *Waist Circumference and Waist-Hip Ratio*. Geneva: WHO.
32. Lohman T, Roche A & Martorell R (1988) *Anthropometric Standardization Reference Manual*. Champaign, IL: Human Kinetics.
33. Habicht JP (1974) Standardization of quantitative epidemiological methods in the field. *Bol Oficina Sanit Panam* **76**, 375–384.
34. World Health Organization (2000) *Obesity: Preventing and Managing the Global Epidemic. Report of a WHO Consultation. WHO Technical Report Series* no. 894. Geneva: WHO.
35. Alberti K, Eckel RH, Grundy SM *et al.* (2009) Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention, National Heart, Lung, and Blood Institute, American Heart Association, World Heart Federation, International Atherosclerosis Society, and International Association for the Study of Obesity. *Circulation* **120**, 1640–1645.
36. Bonvecchio-Arenas A, Fernández-Gaxiola AC, Plazas-Belusteguigoitia M *et al.* (2015) *Guías Alimentarias y de Actividad Física en Contexto de Sobrepeso y Obesidad en la Población Mexicana (Dietary and Physical Activity Guidelines in the Context of Overweight and Obesity in the Mexican Population)*. México, DF: Academia Nacional de Medicina.
37. Bourges H, Casanueva E & Rosado J (2008) *Recomendaciones de Ingestión de Nutrientes para la Población Mexicana: Bases Fisiológicas. Tomo 2 (Recommendations of Nutrient Intake for the Mexican Population: Physiological Basis. Vol. 2)*. México, DF: Editorial Medica Panamericana.
38. World Health Organization (2015) *Guideline: Sugars Intake for Adults and Children*. Geneva: WHO.
39. World Health Organization & Food and Agriculture Organization of the United Nations (2003) *Diet, Nutrition and Prevention of Chronic Diseases. Report of a Joint WHO/FAO Expert Consultation. WHO Technical Report Series* no. 916. Geneva: WHO.
40. Micha R, Shulkin ML, Peñalvo JL *et al.* (2017) Etiologic effects and optimal intakes of foods and nutrients for risk of cardiovascular diseases and diabetes: systematic reviews and meta-analyses from the Nutrition and Chronic Diseases Expert Group (NutriCODE). *PLoS One* **12**, e0175149.
41. Sánchez-Pimienta TG, Batis C, Lutter CK *et al.* (2016) Sugar-sweetened beverages are the main sources of added sugar intake in the Mexican population. *J Nutrition* **146**, issue 9, 1888S–1896S.
42. Vallejo M, Colin-Ramirez E, Rivera Mancia S *et al.* (2017) Assessment of sodium and potassium intake by 24 h urinary excretion in a healthy Mexican cohort. *Arch Med Res* **48**, 195–202.
43. Pérez-Lizaur AB (2014) Sistema Mexicano de alimentos equivalentes. In *Dietas Normales y Terapéuticas: Los Alimentos en la Salud y la Enfermedad*, 6a ed. México, DF: McGraw-Hill.
44. Martínez R & Fernández A (2010) *Impacto Social y Económico del Analfabetismo: Modelo de Análisis y Estudio Piloto*. Santiago: Comisión Económica para América Latina y el Caribe.
45. Robles JN & Navarro DM (2013) Analfabetismo en México: una deuda social. *Rev Int Estad Geogr* **3**, 5–17.
46. Gutiérrez JP (2008) *Clasificación por Niveles Socioeconómicos de los Hogares Entrevistados para la Encuesta Nacional de Salud y Nutrición 2006: Nota Metodológica (Classification of the Interviewed Households by Socioeconomic Levels for the National Health and Nutrition Survey 2006: Methodological Note)*. Cuernavaca, Mexico: National Institute of Public Health.
47. Medina C, Janssen I, Campos I *et al.* (2013) Physical inactivity prevalence and trends among Mexican adults: results from the National Health and Nutrition Survey (ENSANUT) 2006 and 2012. *BMC Public Health* **13**, 1063.



48. Medina C, Barquera S & Janssen I (2013) Validity and reliability of the International Physical Activity Questionnaire among adults in Mexico. *Rev Panam Salud Publica* **34**, 21–28.
49. Pampel FC, Krueger PM & Denney JT (2010) Socioeconomic disparities in health behaviors. *Annu Rev Sociol* **36**, 349–370.
50. Pepper GV & Nettle D (2014) Socioeconomic disparities in health behaviour: an evolutionary perspective. In *Applied Evolutionary Anthropology: Darwinian Approaches to Contemporary World Issues*, pp. 225–243 [MA Gibson and DW Lawson, editors]. New York: Springer.
51. Allen L, Williams J, Townsend N *et al.* (2017) Socioeconomic status and non-communicable disease behavioural risk factors in low-income and lower-middle-income countries: a systematic review. *Lancet Glob Health* **5**, e277–e289.
52. Chen M, Wu Y, Narimatsu H *et al.* (2015) Socioeconomic status and physical activity in Chinese adults: a report from a community-based survey in Jiaying, China. *PLoS One* **10**, e0132918.
53. Smith LP, Ng SW & Popkin BM (2014) No time for the gym? Housework and other non-labor market time use patterns are associated with meeting physical activity recommendations among adults in full-time, sedentary jobs. *Soc Sci Med* **120**, 126–134.
54. Vargas AS, Merino ALH & Hernández IP (2015) La participación laboral femenina y el uso del tiempo en el cuidado del hogar en México. *Contaduría y Administración* **60**, 651–662.
55. Dabbagh-Moghadam A, Mozaffari-Khosravi H, Nasiri M *et al.* (2017) Association of white and red meat consumption with general and abdominal obesity: a cross-sectional study among a population of Iranian military families in 2016. *Eat Weight Disord* **22**, 717–724.
56. Tande DL, Magel R & Strand BN (2010) Healthy Eating Index and abdominal obesity. *Public Health Nutr* **13**, 208–214.
57. Guo X, Warden BA, Paeratakul S *et al.* (2004) Healthy Eating Index and obesity. *Eur J Clin Nutr* **58**, 1580–1586.
58. Carroll RJ, Midthune D, Subar AF *et al.* (2012) Taking advantage of the strengths of 2 different dietary assessment instruments to improve intake estimates for nutritional epidemiology. *Am J Epidemiol* **175**, 340–347.
59. Conrad J & Nothlings U (2017) Innovative approaches to estimate individual usual dietary intake in large-scale epidemiological studies. *Proc Nutr Soc* **76**, 213–219.
60. Freedman LS, Midthune D, Arab L *et al.* (2018) Combining a food frequency questionnaire with 24-hour recalls to increase the precision of estimation of usual dietary intakes—evidence from the validation studies pooling project. *Am J Epidemiol* **187**, 2227–2232.
61. Galobardes B, Morabia A & Bernstein MS (2001) Diet and socioeconomic position: does the use of different indicators matter? *Int J Epidemiol* **30**, 334–340.
62. Howe LD, Galobardes B, Matijasevich A *et al.* (2012) Measuring socio-economic position for epidemiological studies in low-and middle-income countries: a methods of measurement in epidemiology paper. *Int J Epidemiol* **41**, 871–886.
63. Rutstein SO & Johnson K (2004) *The DHS Wealth Index*. *DHS Comparative Reports* no. 6. Calverton, MD: ORC Macro.