

Employment status and temporal patterns of energy intake: Nutrition and Health Survey in Taiwan, 2005–2008

Chun-An Chau¹, Wen-Harn Pan² and Hsin-Jen Chen^{1,*}

¹Institute and Department of Public Health, School of Medicine, National Yang-Ming University, No. 155, Sec. 2, Linong Street, Medical Building II, R213, Beitou District, Taipei 112, Taiwan, Republic of China; ²Institute of Biomedical Sciences, Academia Sinica, Taipei, Taiwan, Republic of China

Submitted 8 February 2017; Final revision received 25 July 2017; Accepted 3 August 2017; First published online 21 September 2017

Abstract

Objective: To (i) identify the major temporal patterns of energy intake among adults; (ii) examine the association between employment status and the patterns; and (iii) examine the association between dietary quality and the patterns.

Design: Secondary analysis based on the cross-sectional population-based nutrition survey in Taiwan, 2005–2008. Based on energy intake levels at six time intervals of a day derived from 24 h recall data, we applied cluster analysis to identify major temporal patterns of energy intake. Self-reported employment status was categorized into six groups: full-time, part-time, no job, student, homemaker and retired. Multinomial logistic regression models were fitted to test the association between temporal patterns of energy intake and employment groups.

Setting: Non-institutionalized community dwellers.

Subjects: Non-pregnant adults (≥ 19 years old) with total energy intake between 2092 and 20 920 kJ/d (500 and 5000 kcal/d; n 4508).

Results: Five major patterns were identified, which can be seen as the traditional meal pattern and its variants. About 20% of adults had the traditional pattern. The most prevalent pattern was the delayed morning meal pattern (33%), which had lower Ca and P intakes than the traditional pattern. About 14% of adults had the delayed lunchtime pattern, which had lower protein, PUFA, fibre, Ca, P, vitamin D and vitamin E intakes than the traditional. Adjusted prevalence of the delayed lunchtime pattern was highest among full-time students (34%), followed by part-time workers (24%), and was lower in retired (8%), homemakers (11%) and full-time employed adults (12%).

Conclusion: Adults' temporal patterns of energy intake, which varied with their employment status, affected their dietary quality.

Keyword
Dietary pattern
Feeding behaviour
Nutrition surveys
Cluster analysis
Meal timing

Temporal patterns of eating have been identified as a novel risk factor for CVD⁽¹⁾. Meal frequency and timing and the temporal distribution of energy intake have been associated with adiposity, obesity, and metabolic regulations such as changes in blood pressure, insulin resistance and lipid profiles^(1,2). For instance, a randomized controlled trial in Japan found that greater energy intake in the evening or night-time snacking may lead to elevated LDL cholesterol⁽³⁾. A cohort study in the UK suggested a relationship between fat intake in the evening and a higher incidence of high blood pressure⁽⁴⁾. In addition, hard outcomes such as type 2 diabetes and cardiovascular mortality were associated with a lower meal frequency in prospective studies^(5,6). Misalignment between the timing of nutrient intake and the circadian rhythm may be the core mechanism of the health impact of temporal patterns of eating⁽⁷⁾.

For adults, employment status is a fundamental factor that influences people's dietary quality and nutritional status^(8–11). Work hours constrain a worker's daily routine, subsequently shaping a person's meal habits such as timing/skipping of meals and snacks^(12,13). Meals such as breakfast, night eating and snacks have been associated with dietary quality^(14–19). However, work-related time constraints should affect the timing of multiple eating occasions of the day. Thus, focusing on a single meal or a certain eating occasion would lose sight of the temporal patterns of eating, as was done in most of the previous studies that attempted to examine the health and/or dietary effects of a single meal^(20–22). To identify the temporal patterns of eating in a free-living population would be the first step towards knowing the relationship between meal time and health impact.

*Corresponding author: Email hsinjenchen@ym.edu.tw

However, temporal patterns of eating are complex and have multiple dimensions: meal frequency, meal timing, regularity, and the dietary composition of meals and snacks⁽²³⁾. Dimension-reduction methods are common tools for researchers to mine the hidden pattern factors that are highly interrelated with certain unknown correlation structures, and therefore are usually used for dietary pattern research⁽²⁴⁾. Recent studies have applied latent class analysis and dynamic clustering analysis to identify the patterns of times when eating episodes occur in a day^(23,25). The study in Australia discovered the usual patterns of meals and snack times. Three patterns called 'conventional', 'late lunch' and 'grazing' were identified, but these patterns were based on the timing of eating occasions without considering the energy intake amounts⁽²³⁾. The other study in the USA accounted for energy, but the whole picture of the temporal distribution of energy intake was unavailable⁽²⁵⁾. Moreover, little is known about the temporal patterns of eating in Asian populations, especially as working hours are notoriously long in East Asian countries^(26,27). The present study aimed to identify adults' major temporal patterns of energy intake based on the 'timed' 24 h dietary recall data in the Nutrition and Health Survey in Taiwan (NAHSIT), 2005–2008. Furthermore, we examined the association between employment status and temporal patterns of energy intake and investigated whether dietary quality in terms of nutrient densities varied by temporal patterns of energy intake.

Methods

Study population

NAHSIT is a representative survey of the Taiwanese non-institutionalized and non-pregnant population that aimed to monitor the nutritional status and health of the population. To obtain representative estimates through a feasible survey method, the survey adopted a multistage, clustered and stratified complex sampling scheme with a probability proportional to size procedure. The survey of years 2005–2008 interviewed 6189 participants of all ages and the response rate was 65% (6189 respondents out of 9570 sampled subjects)^(28,29).

Dietary assessment

A one-day 24 h dietary recall was used to assess the participants' food intake of the previous day⁽³⁰⁾. The recall consisted of multiple passes: (i) the respondent recalled all dishes eaten and the time of eating in the past 24 h from the interview; (ii) the respondent was asked to think about easily forgotten foods and beverages such as dietary supplements and alcohol; (iii) the respondent recalled the ingredients and quantities of food consumption; and (iv) the interviewer reviewed the records with the interviewee. Trained interviewers conducted the computer-assisted personal interview recall. Food models, tableware

models and pictures of common packages of beverages were used to assist the respondent in remembering the volume of food intake. Food items recorded in the dietary recall were then converted to nutrients based on the food composition tables⁽³¹⁾. Participants whose dietary recall had extreme values of energy intake were excluded; thus, the final analysis included adults (aged 19 years or older) whose total energy intake ranged from 2092 to 20 920 kJ/d (500 to 5000 kcal/d; n 4508), a range commonly applied in observational studies to exclude extreme values of energy intake^(32–34).

The recall recorded the participants' time of eating the reported foods, and we used this information to determine the participants' meal times. The term 'meal time' refers to the time of any eating occasion during the day. The recall did not record the meal type associated with the time of food consumption, and thus the time of eating traditional meals such as breakfast, lunch or dinner was unknown. We divided a day into six intervals of 4 h: 05.00–08.59 hours, 09.00–12.59 hours, 13.00–16.59 hours, 17.00–20.59 hours, 22.00–00.59 hours and 01.00–04.59 hours. The time intervals applied were based on a previously published study by de Castro⁽³⁵⁾ and these time intervals overlap with the general population's usual meal times. Participants' recalled food intakes were grouped into these six time intervals. The energy intake levels of the six time intervals were then calculated.

Employment status assessment

The independent variable of interest was employment status. This was based on the participants' self-report on a question of employment status. The question contained the following responses: full-time job, part-time job, self-employed, no stable job, unemployed, no job due to disease, never worked, students, housekeeping and retired. We then categorized these into six groups: full-time (full-time and self-employed), part-time (part-time and no stable job), no job (unemployed, no job due to disease, never worked), student, homemaker and retired.

Covariates

The participants' sex, age, education level and health status were collected in the interview. Their residence was categorized into eight strata based on the counties' urbanization level, geography and ethnicity distribution (e.g. mountainous area, islands, Hakka–dense townships, East coast). Disease history was an indicator variable of being diagnosed with chronic disease(s) that could affect one's diet: hypertension, dyslipidaemia, diabetes, renal disease, heart disease, stroke, fatty liver, gout, peptic ulcer disease or cancer. In addition, since diet could vary between the weekdays and weekend, we further consider the day of dietary recall (weekday *v.* weekend) as a covariate. Income was not considered because: (i) income is conceptually a mediator between employment status

and temporal patterns of eating; and (ii) 32% of self-reported income data was missing. Interview dates were categorized into the following seasons: March–May, June–August, September–November and December–February.

Temporal pattern identification

People who had a similar distribution of energy intake over six time periods of a day were considered to have a similar temporal pattern^(24,36). The pattern categories of the study population were identified based on the similarity of the energy intake levels of the six 4 h intervals. Every participant's energy intakes of the six time periods can be expressed as a six-element vector. To prevent a right-skewed distribution of energy intake, we used log-scale energy intake of every time interval for cluster analysis, while the energy intake of 0 was replaced with 1 so that the log scale can be 0. Hierarchical clustering analysis was used and the Euclidean distance method was used to characterize the closeness among the observations in the six-dimensional data space. The clustering among the observations was then determined using Ward's method. Five temporal patterns of energy intake in the study population were decided upon because: (i) the five clusters explained 55% of data variation; (ii) including one more (the sixth) cluster additionally explained only 0.5% of data variation; and (iii) the six clusters increased the complexity of pattern interpretation and reduced the sample size of every cluster. To confirm the result of cluster categorization, the whole data were split into two data sets and the same five-cluster patterns were reproduced. Then, we examined the average energy intake level at every time period of the day to describe and interpret the five temporal patterns of energy intake.

Statistical analysis

Descriptive statistics were used to describe the differences and ANOVA and χ^2 tests were used to test the differences in basic characteristics among people of the five temporal patterns of energy intake. To examine whether people of different employment statuses had different probabilities of the five patterns, we applied multinomial logit models to estimate the conditional predicted probabilities of the five temporal patterns for every employment status group with adjustments for age, sex, education level, geographic residence and weekday/weekend. Finally, we applied general linear regression models to estimate the adjusted mean nutrient densities, which were expressed as nutrient amount per 4184 kJ (1000 kcal) of total energy intake. The models for estimating nutrient densities were adjusted for age, sex, employment status, education level, geographic residence and weekday/weekend. All statistical analyses were conducted using the statistical software package SUDAAN version 11 (2012) to account for the sampling weight and design effect.

Results

We identified five major temporal patterns of energy intake: 30.7% of adults had the traditional meal pattern (T), 17.6% had the traditional meal pattern plus afternoon eating (TA), 15.6% had the traditional meal pattern plus afternoon and night eating (TAN), 25.6% had a delayed morning meal pattern (DM) and 10.5% had a delayed lunchtime pattern (DL; Fig. 1). The mean (95% CI) total energy intake of patterns T, TA, TAN, DM and DL were, respectively: 7393 (7037, 7749) kJ (1767 (1682, 1852) kcal); 8598 (7950, 9242) kJ (2055 (1900, 2209) kcal); 8816 (8222, 9410) kJ (2107 (1965, 2249) kcal); 8297 (8037, 8556) kJ (1983 (1921, 2045) kcal); and 7096 (6544, 7653) kJ (1696 (1564, 1829) kcal). Pattern T was characterized by large energy intakes in the periods 05.00–08.59, 09.00–12.59 and 17.00–20.59 hours, which are the traditional times for serving breakfast, lunch and dinner. Meanwhile, energy intakes in the off-time meals were close to zero for pattern T. The energy intake distribution of pattern TA was similar to that of pattern T, while the difference between the two patterns was the extra energy intake in the afternoon (13.00–16.59 hours) for the former. Pattern TAN had energy intakes in the three traditional meal times plus afternoon and night (22.00–00.59 hours); thus this pattern had the highest whole-day energy intake among the five patterns. Pattern DM had a low energy intake in the morning (05.00–08.59 hours) and high energy intakes at noon and afternoon periods. Pattern DL was characterized by a very low energy intake at 09.00–12.59 hours, followed by great amounts of energy intake during the afternoon and the evening.

As Table 1 shows, men were more likely to have patterns TAN and DL than women, and women were more likely to have pattern TA than men. Younger adults were more likely to have patterns DM and DL and less likely to have patterns T and TA than the older groups. The distribution of education level by the five temporal patterns of energy intake may reflect the age distributions since older birth cohorts had, in general, fewer years of education than the younger cohort in Taiwan. Adults who reported a history of chronic disease(s) were more likely to have patterns T and TA and less likely to have patterns DM and DL than adults who did not have a disease. Nevertheless, we did not find significant weekday/weekend differences or seasonal variations in the five temporal patterns of energy intake.

Regarding employment status, after adjusting for sex, age, education level, history of chronic diseases, geography and weekday/weekend, we estimated the adjusted proportions of the temporal patterns for the employment status groups (Table 2). This showed that 22.4% of the employed group, 19.9% of homemakers, 19.2% of the retired group and only 4.0% of students had pattern T. The proportion having pattern T in students was significantly lower than that in other employment statuses. At the same time, 33.3% of




Fig. 1 Distribution of energy intake at six time periods of the day by temporal pattern of energy intake among non-institutionalized, non-pregnant, community-dwelling adults (n 4508) aged ≥ 19 years, Nutrition and Health Survey in Taiwan, 2005–2008: (a) traditional meal pattern (T; n 1384, 20.3%*); (b) traditional meal pattern plus afternoon eating (TA; n 795, 14.3%*); (c) traditional meal pattern plus afternoon and night eating (TAN; n 704, 17.7%*); (d) delayed morning meal pattern (DM; n 1154, 33.0%*); (e) delayed lunchtime pattern (DL; n 471, 14.5%*). Value labels near the line present the mean energy intake of the period (with its 95% CI represented by a vertical bar) in kilocalories; to convert to kilojoules, multiply by 4.184. *Weighted percentage

students, 23.6% of part-time employees, 12.0% of full-time employees, 10.5% of homemakers and 8.3% of the retired population had the DL pattern. Sensitivity analyses were done on the following subgroups: women, adults ≥ 45 years old and young adults (19–30 years old). In general, the distribution of the temporal patterns of energy intake by employment status were similar to those found in the whole group analysis (see online supplementary material).

After adjusting for the energy intake level, nutrient density levels also differed by temporal patterns of energy intake (Table 3). Pattern T had the highest intakes of protein, PUFA, Ca, P, vitamin D and vitamin E, and the lowest carbohydrate intake compared with the other temporal patterns. Pattern TA had the highest carbohydrate and the lowest Mg intakes among the five temporal patterns of energy intake. Pattern DL had the lowest nutrient density, including protein, PUFA, dietary fibre, Ca, P, vitamin D and vitamin E. These comparisons were based on models that were adjusted for covariates

including age, sex, education level, employment status, history of chronic diseases, geography and weekday of the dietary recall.

Discussion

Based on 24 h dietary recall, we identified five major temporal patterns of energy intake in Taiwan adults and found that the patterns varied with employment status in this economically developed society. These patterns can be categorized as different variants of pattern T with varying levels of nutrient quality. In general, the nutrient density, including that of protein, PUFA, fibre, Ca, P, vitamins D and E, was higher for pattern T than for pattern DL. Approximately 20% of adults had pattern T and 14% had pattern DL. The most prevalent temporal pattern of energy intake was pattern DM (33%), which had lower nutrient densities of Ca and P than pattern T.

Table 1 Basic characteristics of the participants by temporal pattern of energy intake; non-institutionalized, non-pregnant, community-dwelling adults (*n* 4508) aged ≥ 19 years, Nutrition and Health Survey in Taiwan, 2005–2008

	Temporal pattern of energy intake*											
	All (<i>n</i> 4508)		T (<i>n</i> 1384)		TA (<i>n</i> 795)		TAN (<i>n</i> 704)		DM (<i>n</i> 1154)		DL (<i>n</i> 471)	
	<i>n</i>	%†	<i>n</i>	%†	<i>n</i>	%†	<i>n</i>	%†	<i>n</i>	%†	<i>n</i>	%†
Sex (<i>P</i> = 0.0322)‡												
Men	2244	54.3	704	52.5	352	47.4	390	61.6	555	52.0	243	60.2
Women	2264	45.7	680	47.5	443	52.6	314	38.4	599	48.0	228	39.8
Age (<i>P</i> < 0.0001)‡												
19–30 years	750	33.0	99	17.3	72	18.2	127	34.6	307	39.0	145	53.9
31–44 years	763	27.2	141	21.7	119	24.7	146	30.0	273	33.3	84	19.7
45–64 years	1492	28.3	441	36.1	270	37.7	277	29.5	356	22.3	148	20.6
≥ 65 years	1503	11.5	703	25.0	334	19.3	154	5.9	218	5.4	94	5.7
Education (<i>P</i> = < 0.0001)‡												
None	471	4.1	237	9.2	89	5.6	46	2.1	66	2.0	33	2.3
Middle school or lower	1905	26.9	698	36.9	380	34.9	265	22.4	402	21.6	160	22.9
High school or lower	1546	43.9	328	33.5	255	42.9	285	49.7	489	48.6	189	41.9
College or higher	583	25.1	120	20.4	71	16.6	107	25.8	197	27.8	88	32.9
Missing, <i>n</i> 3												
Employment status (<i>P</i> < 0.0001)‡												
Full-time job	1880	53.8	472	52.8	301	51.2	368	62.0	528	55.5	211	44.1
Part-time job	211	5.5	48	2.9	32	3.9	34	7.0	55	5.3	42	9.0
No job	382	8.5	113	7.0	49	5.8	58	8.7	115	9.6	47	10.5
Student	132	8.3	11	1.3	6	3.4	15	4.9	66	10.1	34	23.0
Retired	1040	10.4	423	18.3	232	17.6	137	8.1	179	6.2	69	4.5
Homemaker	859	13.5	316	17.6	173	18.1	91	9.4	211	13.3	68	9.0
Missing, <i>n</i> 4												
History of chronic disease (<i>P</i> < 0.0001)‡												
None	2243	67.5	567	57.1	350	58.1	366	69.2	679	72.6	281	77.8
Any	2036	32.5	742	43.0	406	41.9	306	30.8	422	27.4	160	22.2
Missing, <i>n</i> 229												
Day of dietary recall (<i>P</i> = 0.1065)‡												
Weekdays	3563	79.1	1121	84.6	644	82.2	547	81.9	889	76.6	362	70.7
Weekend	945	20.9	263	15.4	151	17.8	157	18.1	265	23.4	109	29.3
Season (<i>P</i> = 0.2005)‡												
Mar–May	1241	31.1	390	35.8	235	30.2	206	33.4	292	29.4	118	26.6
Jun–Aug	1123	29.4	357	32.4	208	28.9	168	24.8	274	30.0	116	29.9
Sep–Nov	1124	22.4	365	20.3	191	19.3	175	28.7	275	20.4	118	24.8
Dec–Feb	1020	17.1	272	11.5	161	21.5	155	13.1	313	20.1	119	18.7

*T, traditional meal pattern; TA, traditional meal pattern plus afternoon eating; TAN, traditional meal pattern plus afternoon and night eating; DM, delayed morning meal pattern; DL, delayed lunchtime pattern.

†Weighted statistics.

‡From χ^2 test.

Table 2 Adjusted distribution* of the five temporal patterns of energy intake† by employment status among non-institutionalized, non-pregnant, community-dwelling adults (*n* 4508) aged ≥ 19 years, Nutrition and Health Survey in Taiwan, 2005–2008

Employment status	Distribution (%)				
	T	TA	TAN	DM	DL
Full-time job	22.37 ^D	14.40	19.11	32.08	12.03 ^D
Part-time job	11.33	10.21	23.16	31.72	23.58 ^{E,F}
No job	15.22	10.37	18.72	38.99	16.70
Student	4.01 ^{A,E,F}	11.62	10.37	39.67	34.34 ^{A,E,F}
Retired	19.18 ^D	17.22	19.87	35.40	8.33 ^{B,D}
Homemaker	19.88 ^D	15.18	16.78	37.64	10.51 ^{B,D}

A,B,C,D,E,F Superscript letters (A = full-time job; B = part-time job; C = unemployed; D = student; E = retired; F = homemaker) indicate the groups that had a significantly different proportion of the temporal energy intake pattern compared with this employment status group. For example, 22.37% of full-time employees had pattern T, and the D superscript indicates 22.37% was significantly different from the proportion of students having pattern T. The α level for pairwise comparisons was set at 0.0033 to address the multiple comparisons.

*Adjusted for sex, age, education level, chronic disease, geography and weekday of dietary recall. The model took the sampling weight and study design into account.

†T, traditional meal pattern; TA, traditional meal pattern plus afternoon eating; TAN, traditional meal pattern plus afternoon and night eating; DM, delayed morning meal pattern; DL, delayed lunchtime pattern.

‘Time to eat’ is embedded in the daily schedule, which is highly determined by employment status. Greater tendency to have traditional meal times in full-time

employees was probably due to the time frame of daily work-related activities. For other employment statuses, off-time meals were more likely to happen. A study on

Table 3 Adjusted* mean nutrient densities of the five temporal patterns of energy intake† among non-institutionalized, non-pregnant, community-dwelling adults (*n* 4508) aged ≥19 years, Nutrition and Health Survey in Taiwan, 2005–2008

	T		TA		TAN		DM		DL		P‡
	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	
Protein (g/4184 kJ§)	44.0 ^E	0.7	41.3	0.9	41.9	1.0	42.2	0.6	40.0 ^A	1.6	<0.0001
Fat (g/4184 kJ)	35.3	0.8	33.5	0.6	33.8	0.9	34.5	0.7	33.2	0.8	0.0153
Carbohydrate (g/4184 kJ)	126.9 ^B	1.9	135.1 ^{A,D}	1.7	131.0	2.6	127.7 ^B	1.8	131.7	2.7	0.0008
SFA fats (g/4184 kJ)	11.0	0.3	11.0	0.2	11.2	0.4	11.0	0.3	11.0	0.3	0.9963
MUFA (g/4184 kJ)	12.7	0.3	12.2	0.2	12.5	0.4	12.5	0.4	12.4	0.3	0.2647
PUFA (g/4184 kJ)	11.4 ^{C,E}	0.3	10.2	0.3	10.0 ^A	0.2	10.9 ^E	0.3	9.8 ^{A,D}	0.3	0.0001
Dietary fibre (g/4184 kJ)	9.4 ^E	0.2	10.1	0.8	8.8	0.3	8.2	0.2	7.5 ^A	0.5	0.0088
Cholesterol (mg/4184 kJ)	178.8	6.9	160.8	5.7	164.8	6.2	174.9	6.1	162.3	12.0	0.0397
Ca (mg/4184 kJ)	341.7 ^{C,D,E}	12.4	319.9 ^E	13.9	305.9 ^A	9.2	295.5 ^A	6.4	272.3 ^{A,B}	12.7	0.0001
P (mg/4184 kJ)	621.4 ^{D,E}	9.4	600.5	17.2	584.8	12.3	584.2 ^A	8.8	557.3 ^A	16.9	<0.0001
Mg (mg/4184 kJ)	150.2	2.5	153.1 ^D	4.1	141.0	2.5	139.7 ^B	2.6	135.7	6.9	0.0009
Na (mg/4184 kJ)	2030.8	71.3	1839.4	67.5	1872.0	93.5	2014.9	66.7	1945.4	77.7	0.0583
K (mg/4184 kJ)	1518.5	34.0	1540.7	55.1	1420.9	35.9	1389.0	26.3	1357.0	57.1	0.0074
Zn (mg/4184 kJ)	5.4	0.1	5.3	0.2	5.1	0.1	5.4	0.2	5.1	0.3	0.1153
Fe (mg/4184 kJ)	8.6	0.2	8.0	0.3	8.1	0.2	8.4	0.2	8.1	0.5	0.3344
Retinol equivalents (µg/4184 kJ)	633.2	39.1	578.9	48.4	504.2	28.4	486.6	28.6	511.4	55.8	0.0126
Thiamin (mg/4184 kJ)	0.7	0.0	0.6	0.0	0.7	0.0	0.6	0.0	0.6	0.0	0.7055
Riboflavin (mg/4184 kJ)	0.8	0.0	0.7	0.0	0.7	0.0	0.7	0.0	0.7	0.1	0.1032
Niacin (mg/4184 kJ)	11.0	0.3	10.6	0.4	11.3	0.4	10.6	0.3	10.3	0.4	0.2108
Vitamin B ₆ (mg/4184 kJ)	1.0	0.0	1.0	0.0	1.0	0.0	1.0	0.0	0.9	0.0	0.6892
Vitamin B ₁₂ (µg/4184 kJ)	3.7	0.3	2.7	0.2	3.9	0.6	3.5	0.3	3.1	0.5	0.0218
Vitamin C (mg/4184 kJ)	100.0	6.4	139.9	26.7	94.7	3.7	83.4	3.2	90.1	7.4	0.0838
Vitamin D (µg/4184 kJ)	4.5 ^{C,E}	0.3	3.9	0.3	3.5 ^A	0.2	3.6	0.3	2.9 ^A	0.3	<0.0001
Tocopherol equivalents (mg/4184 kJ)	4.6 ^E	0.2	4.3	0.1	4.2	0.1	4.3	0.1	3.9 ^A	0.1	0.0097

^{A,B,C,D,E}Superscript letters (A = pattern T; B = pattern TA; C = pattern TAN; D = pattern DM; E = pattern DL) indicate the temporal energy intake pattern groups that had a significantly different density of this nutrient compared with this temporal energy intake pattern. For example, the protein nutrient density of pattern T was 44.0 g/4184 kJ, and the E superscript indicates 44.0 g/4184 kJ was significantly different from the protein nutrient density of pattern DL. The α level for pairwise comparisons was set at 0.0033 to address the Bonferroni multiple comparisons.

*Adjusted for sex, age, education level, employment status, chronic disease, geography, and weekday of dietary recall.

†T, traditional meal pattern; TA, traditional meal pattern plus afternoon eating; TAN, traditional meal pattern plus afternoon and night eating; DM, delayed morning meal pattern; DL, delayed lunchtime pattern.

‡P value from ANOVA testing for between-group differences.

§4184 kJ = 1000 kcal.

Northern European populations found that students were more likely to have a daily rhythm of eating that was 'unsynchronized' with the conventional meal times of the general population⁽³⁷⁾. A survey of college students in the USA showed 54% of students snacking in the afternoon and 73% snacking at night⁽³⁸⁾. In Taiwan, we found that full-time students were also the least likely to follow traditional meal times and, on the contrary, were most likely to have a delayed lunchtime compared with other employment statuses. The differences between employment statuses were more obvious for the 19- to 30-year-old group, suggesting that employment status could be an important determinant for young adults to maintain a structured and regular meal pattern and pinpointing college youths' unconventional meal times.

However, the situation may be opposite for the middle-aged and older adults. A survey of German adults aged 45–75 years found that the employed group was more likely to have irregular meal times than those who did not work, and longer working hours were associated with greater odds of having irregular meal times⁽³⁹⁾. This is similar to our findings in subgroup analysis for older adults (≥45 years old). Moreover, the prevalence of pattern DL appeared to be higher in those who had a job than in those who did not work. These findings imply that work-

life time conflict could be critical to the middle-aged population's meal times. On the other hand, among the older adults who did not work, the no job group was, in fact, more likely to delay their morning meals, suggesting a tendency to skip breakfast compared with the other groups. The higher tendency for the no job group to skip a morning meal might reflect a lower food security⁽⁴⁰⁾ or a lower appetite in the morning due to night eating that was associated with unemployment-related stress in other studies^(41,42).

The five temporal patterns of energy intake had different nutrient quality patterns. Especially, the two off-time patterns (DM and DL) we found in the present study had significantly lower nutrient densities than the T pattern. The DM pattern may imply breakfast skipping, which would be followed by more food intake in the subsequent snack or meal times. Eating a morning meal was associated with a lower subsequent energy intake in the afternoon and evening, but it is still controversial whether breakfast skipping might thus lower the total energy intake of the whole day^(19,43). Additionally, foods chosen for breakfast may differ from foods chosen for later meals⁽⁴⁴⁾. People tend to eat wholegrain foods, fruit and vegetables for breakfast⁽¹⁵⁾, and therefore breakfast eaters have higher fibre intake than breakfast skippers^(17,18). In the

present study, the DM pattern had lower Ca, P and Mg intakes than the T pattern. This is similar to the findings of other studies where more frequent breakfast was associated with higher dairy and Ca intakes^(45,46). The other off-time meal pattern was the DL pattern, which was found to have the worst nutrient quality compared with other temporal patterns of energy intake in Taiwan. This may be associated with the larger amounts of snacks in the afternoon and the evening. A study on Korean workers found that meal skippers were more likely to eat many snacks⁽¹⁶⁾. The time of day when foods are consumed may also influence overall nutrient intake; people tend to consume greater amounts of energy during eating occasions at later times of the day⁽³⁵⁾. In addition, snacking may also play a role: people tend to snack in the afternoon and the evening, and they are less likely to choose fruits at snack times⁽⁴⁷⁾. Since snacks are usually carbohydrate-dense and result in a greater total energy intake^(48,49), snacking more in the afternoon and the evening is associated with higher empty calories and lowers nutrient density. As the literature suggests, eating desserts and sweetened beverages for snacks would lower the dietary quality⁽¹⁴⁾.

Strengths and limitations

The present study was based on a representative sample of Taiwanese adults and the results may reflect what occurred in Taiwan. Moreover, the cluster analysis identified common temporal patterns of energy intake based on six 4 h intervals of a day, rather than based on the self-reported meal types. This approach can prevent cultural or habitual influences on the self-identified meal types and focus on the temporal distribution of energy intake throughout the day⁽²⁵⁾. Nevertheless, there are several limitations to be considered before making conclusions. First, the dietary intake was based on a one-day recall. The one-day snapshot may not represent a person's usual dietary intake, but a group's one-day dietary information may reveal the group's common dietary conditions. Thus, the results of the study should be interpreted at the population level. Nevertheless, if multiple days of recall could be collected, it would be possible to examine whether the temporal patterns of energy intake vary between days and to explore the determinants. Second, the employment status was based on individuals' self-report. Since the self-report error in employment status should be unrelated to the dependent variables, the temporal patterns of energy intake and nutrient quality, the impact of the self-report error would attenuate the association between employment status and temporal patterns of energy intake. Since there was no information for the work time schedule, we could not examine the influence of work shifts on temporal patterns of energy intake. The results based on the sample that pooled shift workers and others together may underestimate the probability of the off-time patterns in the

employed population. Third, it is unknown what meal the eating occasion was for and this makes it difficult to describe the patterns. In the present study, the patterns were named based on the meal time, rather than meal type, to prevent misinterpretation of the temporal patterns of energy intake into meal patterns. Fourth, employment status is highly associated with household income, but 32% of respondents had missing values of income level. Based on the complete data, *post hoc* analysis that additionally adjusted for income did not affect the results substantially (data not shown). However, future study is needed to understand the following pathways: whether time use, income or other factors of employment status influence the temporal patterns of eating. Fifth, the study was based on Taiwanese adults and the findings may not generalize to other populations.

Conclusion

The present study identified five major temporal patterns of energy intake in Taiwan. Adults who ate the traditional meal pattern had a better dietary (nutrient) quality compared with those who ate the other patterns. The traditional meal pattern was more likely to be practised in full-time workers, homemakers and retired people, compared with the other employment statuses. The worst quality of nutrient intake was found with the delayed lunchtime pattern, which was more common in adult students and part-time workers. The study suggests that the health disparity associated with nutrition might be introduced by the temporal patterns of energy intake. Further research is needed to understand how meal timing and temporal eating patterns would be associated with health status.

Acknowledgements

Financial support: This original research was partly supported by the Taiwan Ministry of Science and Technology (MOST 104-2320-B-010-042, 105-2314-B-010-012, 106-2314-B-010-010). Data analysed in this article were collected by the Nutrition and Health Survey in Taiwan (NAHSIT 2005–2008) sponsored by the Department of Health in Taiwan (DOH94-fs-6-4), a survey carried out by the Center for Survey Research of Academia Sinica, directed by Professor Wen-Harn Pan. The Office of Nutrition Survey, the Institute of Biomedical Sciences, Academia Sinica is responsible for data distribution. The assistance provided by the institutes and efforts made by all contributed to the survey are greatly appreciated. The views expressed herein are solely those of the authors. *Conflict of interest:* The authors have no competing interest to disclose. *Authorship:* C.-A.C. analysed the data and drafted the manuscript; W.-H.P. conceived the idea and reviewed the manuscript critically; H.-J.C. conceived the idea, analysed the data, and

wrote and reviewed the manuscript. *Ethics of human subject participation*: The study was based on anonymous open domain data set, and was approved by the National Yang-Ming University Internal Review Board.

Supplementary material

To view supplementary material for this article, please visit <https://doi.org/10.1017/S1368980017002476>

References

1. St-Onge MP, Ard J, Baskin ML *et al.* (2017) Meal timing and frequency: implications for cardiovascular disease prevention: a scientific statement from the American Heart Association. *Circulation* **135**, e96–e121.
2. Ekmekcioglu C & Touitou Y (2011) Chronobiological aspects of food intake and metabolism and their relevance on energy balance and weight regulation. *Obes Rev* **12**, 14–25.
3. Hibi M, Masumoto A, Naito Y *et al.* (2013) Nighttime snacking reduces whole body fat oxidation and increases LDL cholesterol in healthy young women. *Am J Physiol Regul Integr Comp Physiol* **304**, R94–R101.
4. Almoosawi S, Prynne CJ, Hardy R *et al.* (2013) Time-of-day of energy intake: association with hypertension and blood pressure 10 years later in the 1946 British Birth Cohort. *J Hypertens* **31**, 882–892.
5. Chen H-J, Wang Y & Cheskin LJ (2016) Relationship between frequency of eating and cardiovascular disease mortality in US adults: the NHANES III follow-up study. *Ann Epidemiol* **26**, 527–533.
6. Mekary RA, Giovannucci E, Willett WC *et al.* (2012) Eating patterns and type 2 diabetes risk in men: breakfast omission, eating frequency, and snacking. *Am J Clin Nutr* **95**, 1182–1189.
7. Johnston JD (2014) Physiological responses to food intake throughout the day. *Nutr Res Rev* **27**, 107–118.
8. Ali SM & Lindstrom M (2006) Socioeconomic, psychosocial, behavioural, and psychological determinants of BMI among young women: differing patterns for underweight and overweight/obesity. *Eur J Public Health* **16**, 325–331.
9. Alkerwi A, Vernier C, Sauvageot N *et al.* (2015) Demographic and socioeconomic disparity in nutrition: application of a novel correlated component regression approach. *BMJ Open* **5**, e006814.
10. Lee KW, Song WO & Cho MS (2016) Dietary quality differs by consumption of meals prepared at home vs. outside in Korean adults. *Nutr Res Pract* **10**, 294–304.
11. Vernay M, Malon A, Oleko A *et al.* (2009) Association of socioeconomic status with overall overweight and central obesity in men and women: the French Nutrition and Health Survey 2006. *BMC Public Health* **9**, 215.
12. Smith KJ, McNaughton SA, Cleland VJ *et al.* (2013) Health, behavioral, cognitive, and social correlates of breakfast skipping among women living in socioeconomically disadvantaged neighborhoods. *J Nutr* **143**, 1774–1784.
13. Suzuki A, Sakurazawa H, Fujita T *et al.* (2016) Overeating, late dinner, and perceived stress in Japanese workers. *Obes Res Clin Pract* **10**, 390–398.
14. Barnes TL, French SA, Harnack IJ *et al.* (2015) Snacking behaviors, diet quality, and body mass index in a community sample of working adults. *J Acad Nutr Diet* **115**, 1117–1123.
15. Kant AK & Graubard BI (2015) Within-person comparison of eating behaviors, time of eating, and dietary intake on days with and without breakfast: NHANES 2005–2010. *Am J Clin Nutr* **102**, 661–670.
16. Kim SY & Kim SM (2010) Energy intake and snack choice by the meal patterns of employed people. *Nutr Res Pract* **4**, 43–50.
17. Myhre JB, Loken EB, Wandel M *et al.* (2015) Meal types as sources for intakes of fruits, vegetables, fish and whole grains among Norwegian adults. *Public Health Nutr* **18**, 2011–2021.
18. Pereira MA, Erickson E, McKee P *et al.* (2011) Breakfast frequency and quality may affect glycemia and appetite in adults and children. *J Nutr* **141**, 163–168.
19. Purslow LR, Sandhu MS, Forouhi N *et al.* (2008) Energy intake at breakfast and weight change: prospective study of 6,764 middle-aged men and women. *Am J Epidemiol* **167**, 188–192.
20. Bi H, Gan Y, Yang C *et al.* (2015) Breakfast skipping and the risk of type 2 diabetes: a meta-analysis of observational studies. *Public Health Nutr* **18**, 3013–3019.
21. Mathias KC, Jacquier E & Eldridge AL (2016) Missing lunch is associated with lower intakes of micronutrients from foods and beverages among children and adolescents in the United States. *J Acad Nutr Diet* **116**, 667–676.e6.
22. O'Neil CE, Byrd-Bredbenner C, Hayes D *et al.* (2014) The role of breakfast in health: definition and criteria for a quality breakfast. *J Acad Nutr Diet* **114**, 2 Suppl., S8–S26.
23. Leech RM, Worsley A, Timperio A *et al.* (2017) Temporal eating patterns: a latent class analysis approach. *Int J Behav Nutr Phys Act* **14**, 3.
24. Hu FB (2002) Dietary pattern analysis: a new direction in nutritional epidemiology. *Curr Opin Lipidol* **13**, 3–9.
25. Eicher-Miller HA, Khanna N, Boushey CJ *et al.* (2016) Temporal dietary patterns derived among the adult participants of the National Health and Nutrition Examination Survey 1999–2004 are associated with diet quality. *J Acad Nutr Diet* **116**, 283–291.
26. Lee S, McCann D & Messenger JC (2007) *Working Time Around the World*. Geneva: International Labour Organization.
27. Tsai M-C, Nitta M, Kim S-W *et al.* (2016) Working overtime in East Asia: convergence or divergence? *J Contemp Asia* **46**, 700–722.
28. Tu SH, Chen C, Hsieh YT *et al.* (2011) Design and sample characteristics of the 2005–2008 Nutrition and Health Survey in Taiwan. *Asia Pac J Clin Nutr* **20**, 225–237.
29. Pan W-H & Tu S-H (2008) *2004–2008 Nutrition and Health Survey in Taiwan (NAHSIT) (D00090) [Data file]*. Taipei: Survey Research Data Archive, Center for Survey Research, Research Center for Humanities and Social Sciences, Academia Sinica.
30. Pan WH, Wu HJ, Yeh CJ *et al.* (2011) Diet and health trends in Taiwan: comparison of two nutrition and health surveys from 1993–1996 and 2005–2008. *Asia Pac J Clin Nutr* **20**, 238–250.
31. Wu SJ, Pan WH, Yeh NH *et al.* (2011) Trends in nutrient and dietary intake among adults and the elderly: from NAHSIT 1993–1996 to 2005–2008. *Asia Pac J Clin Nutr* **20**, 251–265.
32. Bruening M, van Woerden I, Todd M *et al.* (2016) A mobile ecological momentary assessment tool (devilSPARC) for nutrition and physical activity behaviors in college students: a validation study. *J Med Internet Res* **18**, e209.
33. Howarth NC, Murphy SP, Wilkens LR *et al.* (2006) Dietary energy density is associated with overweight status among 5 ethnic groups in the multiethnic cohort study. *J Nutr* **136**, 2243–2248.
34. Zhang Z, Gillespie C, Welsh JA *et al.* (2015) Usual intake of added sugars and lipid profiles among the US adolescents: National Health and Nutrition Examination Survey, 2005–2010. *J Adolesc Health* **56**, 352–359.
35. de Castro JM (2004) The time of day of food intake influences overall intake in humans. *J Nutr* **134**, 104–111.

36. Mooi E & Sarstedt M (2011) Cluster analysis. In *A Concise Guide to Market Research: The Process, Data, and Methods using IBM SPSS Statistics*, pp. 237–284. Berlin/Heidelberg: Springer.
37. Lund TB & Gronow J (2014) Deconstruction or continuity? The daily rhythm of eating in Denmark, Finland, Norway and Sweden in 1997 and 2012. *Appetite* **82**, 143–153.
38. Driskell JA, Kim YN & Goebel KJ (2005) Few differences found in the typical eating and physical activity habits of lower-level and upper-level university students. *J Am Diet Assoc* **105**, 798–801.
39. Muff C, Reinhardt JD, Erbel R *et al.* (2011) Who is at risk of irregular meal intake? Results from a population-based study. *J Public Health* **19**, 453–462.
40. Chun IA, Ryu SY, Park J *et al.* (2015) Associations between food insecurity and healthy behaviors among Korean adults. *Nutr Res Pract* **9**, 425–432.
41. Allison KC, Lundgren JD, O'Reardon JP *et al.* (2008) The Night Eating Questionnaire (NEQ): psychometric properties of a measure of severity of the Night Eating Syndrome. *Eat Behav* **9**, 62–72.
42. Andersen GS, Stunkard AJ, Sorensen TI *et al.* (2004) Night eating and weight change in middle-aged men and women. *Int J Obes Relat Metab Disord* **28**, 1338–1343.
43. Levitsky DA & Pacanowski CR (2013) Effect of skipping breakfast on subsequent energy intake. *Physiol Behav* **119**, 9–16.
44. Birch LL, Billman J & Richards SS (1984) Time of day influences food acceptability. *Appetite* **5**, 109–116.
45. Lai I-J (2006) Application of an ecological model to dairy product consumption behavior among eighth graders in Taipei, Taiwan. PhD Thesis, Oregon State University.
46. Lee S & Reicks M (2003) Environmental and behavioral factors are associated with the calcium intake of low-income adolescent girls. *J Am Diet Assoc* **103**, 1526–1529.
47. Cross AT, Babicz D & Cushman LF (1994) Snacking patterns among 1,800 adults and children. *J Am Diet Assoc* **94**, 1398–1403.
48. Bellisle F, Dalix AM, Mennen L *et al.* (2003) Contribution of snacks and meals in the diet of French adults: a dietary study. *Physiol Behav* **79**, 183–189.
49. Chapelot D (2011) The role of snacking in energy balance: a biobehavioral approach. *J Nutr* **141**, 158–162.