

Using Local Group galaxies to investigate the influence of blending on Cepheid distances and the cosmological distance scale

Barbara J. Mochejska

*Harvard-Smithsonian Center for Astrophysics, 60 Garden St,
Cambridge, MA 02138, USA*

Lucas M. Macri

*National Optical Astronomy Observatories, North Cherry Ave., Tucson,
AZ 85719, USA*

Dimitar D. Sasselov, Krzysztof Z. Stanek

*Harvard-Smithsonian Center for Astrophysics, 60 Garden St,
Cambridge, MA 02138, USA*

Abstract. We investigate the influence of blending on the Cepheid distance scale using two Local Group galaxies, M31 and M33. Blending leads to systematically low distances to galaxies observed with the HST, and therefore to systematically high estimates of H_0 . High-resolution HST images are compared to our ground-based data, obtained as part of the DIRECT project, for a sample of 22 Cepheids in M31 and 102 Cepheids in M33. For a sample of 22 Cepheids in M31, the average (median) flux contribution from luminous companions not resolved on the ground-based images in the V -band, S_V , is about 19% (12%) of the flux of the Cepheid. For 102 Cepheids in M33 the average (median) values of S_V , S_I , S_B are 23% (13%), 28% (20%), 28% (15%). For 64 Cepheids in M33 with periods in excess of 10 days the average (median) S_V , S_I , S_B are 16% (7%), 23% (12%), 20% (10%).

1. Introduction

As the number of extragalactic Cepheids discovered with HST continues to increase and the value of H_0 is sought from distances based on these variables, it becomes even more important to understand various possible systematic errors which could affect the extragalactic distance scale. Currently, the most important systematic is a bias in the distance to the Large Magellanic Cloud, which provides the zero-point calibration for the Cepheid distance scale (Udalski 2000; Fitzpatrick et al. 2003). Another possible systematic, the metallicity dependence of the Cepheid Period-Luminosity (PL) relation, is also very much an open issue, with empirical determinations ranging from 0 to $-0.4 \text{ mag dex}^{-1}$ (Sasselov et al. 1997; Udalski et al. 2001).




Figure 1. The cumulative probability distributions of the blending parameter S_V for the artificial Cepheids (dashed line) and the Cepheid catalog (solid).

We define *blending* as the close projected association of a Cepheid with one or more intrinsically luminous stars, which cannot be detected within the observed point-spread function (PSF) by photometric analysis. Blending is thus a phenomenon different from *crowding* or *confusion noise*; the latter occurs in stellar fields with a crowded and complex background due to the random superposition of stars with different luminosities.

We investigate the effects of stellar blending on the Cepheid distance scale by studying two Local Group spiral galaxies, M31 and M33. We identify some of the Cepheids, discovered by the DIRECT project (Stanek et al. 1999, Mochejska et al. 1999) on archival HST-WFPC2 images and compare them to our ground-based data to estimate the impact of blending on our photometry, taking advantage of their superior resolution.

2. The blending catalogs

We have adopted three criteria that a companion to a Cepheid has to fulfill to be included into our catalog as a blend. The star has to: (1) be located at a distance less than $0''.75$ from the Cepheid (half the typical full width at half maximum on ground-based images); (2) be undetected by DAOPHOT in our ground-based images; and (3) contribute at least 4% (for M31) or 6% (for M33) of the flux of the Cepheid in the same filter.




Figure 2. Blending/distance bias as a function of the summing radius/distance. The average and median S_V are indicated with open and filled symbols, respectively. The Ferrarese et al. (2000) results are indicated with asterisks.

To quantify blending we have used the parameter S_F , defined in Mochejska et al. (2000) as the sum of all flux contributions from blends in filter F normalized to the flux of the Cepheid: $S_F = \sum_{i=1}^{N_F} \frac{f_i}{f_C}$ where f_i is the flux of the i -th blend, f_C the flux of the Cepheid on the HST image and N_F the total number of blends. Table 1 shows the blending statistics for M31 and M33. The detailed blending catalogs can be found in Mochejska et al. (2000, 2001). The cumulative probability distribution of blending in M33 in the V filter is plotted in Fig. 1.

Table 1. Blending statistics for M31 and M33

Period range	S_V			S_I			S_B		
	avg	med	N	avg	med	N	avg	med	N
M31 all periods	0.19	0.12	22						
M33 all periods	0.24	0.14	95	0.30	0.21	62	0.29	0.15	57
M33 $P < 10$ d	0.37	0.25	35	0.43	0.29	20	0.47	0.26	18
M33 $P > 10$ d	0.16	0.07	60	0.23	0.14	42	0.20	0.10	39

3. Crowding vs blending – artificial star tests

If we assume that a Cepheid is associated with other luminous stars located in its proximity, then moving it to a randomly chosen position on the image will break that association. In the former case, the Cepheid will be subject to blending; in the latter, to crowding.

To estimate the influence of crowding we have generated an artificial catalog: for each Cepheid observed on a WFPC2 image we generated a list of 100 random positions and determined the contribution from companions at that location.

Fig. 1 shows the cumulative probability distributions for S_V drawn from the artificial crowding catalog (dashed line) and the Cepheid blending catalog (solid line). We have divided the sample of M33 Cepheids into two bins in period at $P = 10$ d and three bins in surface brightness, corresponding to regions near the nucleus, inside the spiral arms and in between them.

For Cepheids located in regions of lowest surface brightness blending is stronger than crowding. The two effects appear comparable in magnitude for Cepheids located in intermediate surface brightness regions. For Cepheids located in the highest surface brightness regions blending is weaker than crowding, most likely due to selection effects.

This comparison indicates that the importance of blending, relative to crowding, very likely increases with decreasing surface brightness. This is not unexpected, as young stars are known to cluster (Harris & Zaritsky 1999). Increasing the level of crowding will tend to obscure this effect.

4. Indications for remote galaxies

Using the HST M33 data as the template we have obtained an estimate of the effect that blending would have on this galaxy if it were observed at further distances. By increasing the radius around the Cepheid for summing the contributions from the blends we can simulate the deterioration of resolution due to the increasing distance to the galaxy. We have restricted ourselves to the long period Cepheids ($P > 10$ d), as they are preferred for determining distances. We have also rejected all Cepheids with $S_V > 45\%$, assuming that they will be recognized as blended based on the shape of the light curve and rejected from the sample. The bias in distance due to blending is illustrated in Fig. 2 as a function of distance. The distance underestimate increases from 0%–3% to 8% at 10–15 Mpc and levels off at 9%–10% at 25–30 Mpc. This indicates that blending could potentially be a substantial source of error in the Cepheid distance scale.


References

- Fitzpatrick, E.L., et al. 2003, *ApJ*, 587, 685
 Harris, J., Zaritsky, D. 1999, *AJ*, 117, 2831
 Mochejska, B.J., et al. 1999, *AJ*, 118, 2211
 Mochejska, B.J., Macri, L.M., Sasselov, D.D., Stanek, K.Z. 2000, *AJ*, 120, 810
 Mochejska, B.J., Macri, L.M., Sasselov, D.D., Stanek, K.Z. 2001, *astro-ph/0103440*
 Sasselov, D.D., et al. 1997, *A&A*, 324, 471
 Stanek, K.Z., et al. 1999, *AJ*, 117, 2810
 Udalski, A., et al. 2000, *Acta Astronomica*, 50, 307
 Udalski, A., et al. 2001, *Acta Astronomica*, 51, 221

Discussion

Alves: When you add flux from neighbouring stars, do you add all of it? Or do you account for these stars contributing to the “sky” background which is subtracted by doing photometry?

Mochejska: We add all of the flux - it all counts as part of the blend.



Barbara Mochejska