

Mini-spiral as source of material for Sgr A* in bright state

Bożena Czerny,¹ Vladimír Karas,² Devaky Kunneriath,²
and Tapas K. Das³

¹Copernicus Astronomical Center, Bartycka 18, 00-716 Warsaw, Poland
email: bcz@camk.edu.pl

²Astronomical Institute, Academy of Sciences,
Boční II 1401, CZ-14131 Prague, Czech Republic
email: vladimir.karas@cuni.cz, devaky@ig.cas.cz

³Harish Chandra Research Institute, Allahabad 211019, India
email: tapas@mri.ernet.in

Abstract. The question of the origin of the gas supplying the accretion process is pertinent especially in the context of enhanced activity of Galactic Center during the past few hundred years, seen now as echo from the surrounding molecular clouds, and the currently observed new cloud approaching Sgr A*. We discuss the so-called Galactic Center mini-spiral as a possible source of material feeding the supermassive black hole on a 0.1 parsec scale. The collisions between individual clumps reduce their angular momentum, and set some of the clumps on a plunging trajectory.

We conclude that the amount of material contained in the mini-spiral is sufficient to sustain the luminosity of Sgr A* at the required level. The accretion episodes of relatively dense gas from the mini-spiral passing through a transient ring mode at $\sim 10^4$ Rg provide a viable scenario for the bright phase of Galactic Center.

Keywords. Galaxy: center; Accretion, accretion disks; Black hole physics

1. Introduction

The supermassive black hole at the Milky Way center (Sgr A*) has the mass of about $4.4 \times 10^6 M_\odot$ (Genzel *et al.* 2010). It presently remains in a very quiet state. However, observations of the X-ray reflection from molecular clouds seem to imply that Sgr A* was orders of magnitude brighter only a few hundred years ago (see Ponti *et al.* 2010, and further references therein). It is interesting to note the recent discovery of a cloud identified at the distance of only a few thousands gravitational radii; see Gillessen *et al.* (2012). The cloud moves along a highly eccentric orbit, and so the latter authors suggest that will reach the minimum distance from the Sgr A* in 2013. Here we consider the mini-spiral as a possible source of the intermittently inflowing material.

2. Model

The Galactic Center black hole is surrounded by the minispiral, a complex system of gas and dust (Zhao *et al.* 2010, Kunneriath *et al.* 2012). The currently adopted explanation assumes a superposition of three elongated clumpy streams of material in a roughly Keplerian motion. The Northern Arm and the Eastern Arm appear to collide about 0.1–0.2 pc behind Sgr A* (Zhao *et al.* 2010). The hot gas can accrete directly onto the black hole. The Bondi radius for a cold gas is much smaller, so some kind of dissipative losses of angular momentum in collisions are needed.




Figure 1. Two examples of model fits that are consistent with data points from Terrier *et al.* (2010), Yu *et al.* (2011). Left: events without reprocessing (single event: $t_{\text{event}} = 1990.7$, $t_{\text{decay}} = 12$ years, dashed blue; a series of events $t_{\text{event}} = 1960\text{--}1995.5$, $t_{\text{decay}} = 9$ years). Right: single flare event with a decay time-scale $t_{\text{decay}} = 9$ years (blue continuous line – intrinsic profile ($t_{\text{event}} = 1969$), green dashed line – the scattered signal assuming the cloud size 5 pc.

The next stage of the evolution proceeds by action of viscosity. The time evolution of a gas located initially in a ring can be described analytically, and the accretion rate at the inner edge of the newly formed disk is described by the formula (Zdziarski *et al.* 2009)

$$\dot{M}(\tau) = \frac{M_{\text{ring}}}{4\pi R_{\text{circ}}^2 t_{\text{visc}}} \frac{(2\mu^2)^\mu}{\Gamma(\mu)} \tau^{-1-\mu} \exp\left(-\frac{2\mu^2}{\tau}\right), \quad (2.1)$$

where μ describes the viscosity, and τ_{visc} depends on the circularization radius. We adopt $\mu = 2/3$ in further considerations. We model the decay timescale of the $K\alpha$ reflection from the molecular cloud B2 assuming a range of accretion events distributed in time and/or in angular momentum. We also consider the smearing of the signal by the reprocessing due to a finite size of the cloud.

3. Results

We considered several examples of the model lightcurves, with and without the effect of reprocessing by the extended region of the B2 molecular cloud. Fig. 1 left panel shows the evolution for the case of a narrow accretion ring, while the right panel shows the result for a whole range of angular momentum. For a slowly cooling material, the fitted decay time-scale 12 years (without reprocessing) implies an accretion event settling as a ring $\sim 1.4 \times 10^4 R_g$, i.e. only by factor 20 less than the inner edge of the mini-spiral, and the infalling matter can retain as much as 20 per cent of the initial angular momentum. Reprocessing slightly decreases this value to 9 years and introduces a considerable delay of the event. For further details see our paper (A&A submitted).

References

- Genzel, R., Eisenhauer, F., & Gillesen, S. 2010, *Rev. Mod. Phys.*, 82, 3121
 Gillesen, S. *et al.* 2009, *ApJ*, 692, 1075
 Kunneriath, D. *et al.* 2012, *A&A*, 538, A127
 Ponti, G. *et al.* 2010, *ApJ*, 714, 732
 Terrier, R. *et al.* 2010, *ApJ*, 719, 143
 Yu, Y. W. *et al.* 2011, *MNRAS*, 411, 2002
 Zdziarski, A. A., Kawabata, R., & Mineshige, S. 2009, *MNRAS*, 399, 1633
 Zhao, J. H. *et al.* 2010, *ApJ*, 723, 1097