

## Recent Results from the CANGAROO Project

S.A. Dazeley, P.G. Edwards, J.R. Patterson, G.P. Rowell, M. Sinnott,  
G.J. Thornton, C. Wilkinson

*CANGAROO Collaboration, Australia,*  
*Department of Physics and Mathematical Physics,*  
*University of Adelaide, 5005.*

T. Hara, N. Hayashida, H. Katsumata, T. Kifune, T. Konishi,  
Y. Matsubara, T. Matsuoka, Y. Mizumoto, M. Mori, M. Muraishi,  
Y. Muraki, T. Oda, S. Ogio, T. Ohsaki, T. Sako, K. Sakurazawa,  
R. Susukita, A. Suzuki, T. Tamura, T. Tanimori, S. Yanagita,  
T. Yoshida and T. Yoshikoshi

*CANGAROO Collaboration, Japan,*  
*Institute for Cosmic Ray Research,*  
*University of Tokyo, Tokyo 188.*

### 1. The CANGAROO Project

The Collaboration of Australia and Nippon for a Gamma Ray Observatory in the Otback operates two large telescopes at Woomera (South Australia), which detect the Čerenkov light images produced in the atmosphere by electron-positron cascades initiated by very high energy ( $\sim 1$  TeV or  $10^{12}$  eV) gamma rays. These gamma rays arise from a different mechanism than at EGRET energies: inverse Compton (IC) emission from relativistic electrons.

The spoke-like images are recorded by a multi-pixel camera which facilitates the rejection of the large numbers of oblique and ragged cosmic ray images. A field of view  $\sim 3.5^\circ$  is required. The Australian team operates a triple 4 m diameter mirror telescope, BIGRAT, with a 37 photomultiplier tube camera and energy threshold 600 GeV. The Japanese operate a single, highly accurate 3.8 m diameter f/1 telescope and high resolution 256 photomultiplier tube camera. In 1998 a new 7 m telescope is planned for Woomera with a design threshold  $\sim 200$  GeV.

### 2. Egret pulsars

#### 2.1. The Crab

The Crab, a synchrotron nebula and pulsar at a distance of 1.5 kpc, is a standard candle for TeV work. We have observed it from Woomera, where it reaches a maximum elevation of  $38^\circ$ . At an energy threshold of 7 TeV, a  $4\sigma$  excess of gamma-ray events was found (Tanimori et al. 1994). More recent observations have confirmed this result.

## 2.2. PSR B1706-44

This isolated southern 102 ms pulsar is only the second pulsar known to emit  $\sim 1$  TeV photons so its confirmation and further study are important. Our observations were made in 1992-1993 and show steady non-periodic emission in both years with a combined statistical significance of  $12\sigma$  (Kifune et al. 1995). The measured flux of  $1 \times 10^{-11}$  photons  $\text{cm}^{-2} \text{s}^{-1}$  above 1 TeV is approximately 2 orders of magnitude below the extrapolated EGRET spectrum. Using BIGRAT data alone, the upper limit is  $4.6 \times 10^{-11}$  photons  $\text{cm}^{-2} \text{s}^{-1}$  above 600 GeV.

## 2.3. The Vela pulsar

The Vela pulsar is young (11,000 y) and relatively close ( $\sim 500$  pc). It is by far the brightest GeV gamma ray source observed by EGRET. The GeV emission is almost 100% pulsed. Recently Rosat observations below 2 keV have revealed a relativistic jet extending SSW from the pulsar along the magnetic axis out to  $0.4^\circ$  (Markwardt and Ogelman, 1995). emission. The TeV data show unpulsed gamma-ray emission from a position offset by  $\sim 0.05^\circ$  from the pulsar, near the jet. An excess was detected each year over the three year period 1993-1995. The jet may cause shock acceleration of energetic particles which could result in the emission of TeV gamma rays. Further work is needed to confirm this important result.

## 2.4. PSR B1055-52

PSR 1055-52 is a 197 ms EGRET pulsar at a distance of 1.5 kpc. It is notable as a very efficient converter of rotational energy into GeV gamma rays. However, no TeV emission has so far been observed in 2 years of CANGAROO observations.

## 3. Other pulsars

PSR 1259-63 is a binary pulsar. At periastron in January 1994, it passed very close to its companion. During these 4 days observation a  $3\sigma$  excess was observed by Cangaroo. Further observations are needed at the next periastron in 1997.

PSR J0437-4715 is a binary 5.757 ms pulsar. The system is only 140 parsecs away. The spin down energy flux density,  $\dot{E}/4\pi D^2$  is the same order of magnitude as PSR B1706-44. It has not been detected by EGRET. Our search for emission from PSR 0437-47 at the radio period yielded no evidence for TeV gamma ray emission. An upper limit for a pulsed flux above  $\sim 1$  TeV of  $5 \times 10^{-11}$  photons  $\text{cm}^{-2} \text{s}^{-1}$  was obtained by BIGRAT.

## References

- Kifune, T. et al., 1995, ApJ, 438, L91.  
Markwardt, C.B. and Ogelman, H., 1995, Nature 375, 40.  
Tanimori, T. et al., 1994, ApJ, 429, L61.