

On the Physical Mechanism Underlying PC1 and Baldwin Effect in Quasars

Xiao-Bo Dong¹, Ting-Gui Wang¹, Jian-Guo Wang^{2,1}, Xiaohui Fan³,
Huiyuan Wang¹, Hongyan Zhou¹, and Weimin Yuan²

¹Center for Astrophysics, University of Science and Technology of China
Email: xbdong@ustc.edu.cn

²Yunnan Observatory, Chinese Academy of Sciences, Kunming, Yunnan 650011, China

³Steward Observatory, The University of Arizona, Tucson, AZ 85721, USA

Keywords. galaxies: active, quasars: emission lines, quasars: general

Recent work (Baskin & Laor 2004; Dong *et al.* 2009a,b) suggests that the Eddington ratio ($\ell \equiv L/L_{\text{Edd}}$) is the origin of all the significant first-order object-to-object variations of quasar spectral properties from the zeroth-order similarity of AGN spectra; specifically, this includes the PC1 of Boroson & Green (1992), the classic or inverse Baldwin effect (Baldwin 1977), and even blueshifting (i.e., blue asymmetry) of high-ionization emission lines (Dong *et al.* 2009c).

To understand such unification, we shift our focus from the detailed physics of the accretion process (microphysics) to “statistical physics” (macrophysics) of the broad-line clouds (Dong *et al.*, in preparation; cf. Korista 1999). With more realistic constraints (e.g., the distribution of the number of line-emitting clouds with ℓ and other physical parameters), we believe that the first-order regularities and even the second-order effects of quasar emission lines may be reproduced exactly by future “locally optimally-emitting cloud” models (LOC; Baldwin *et al.* 1995).

On one hand, for clouds gravitationally bound around the central engine, there is a lower limit on the column density N_{H} that is set by and roughly scales with L/L_{Edd} (Dong *et al.* 2009a). On the other hand, in accretion-powered radiation systems such as quasars, L/L_{Edd} is closely related to \dot{m} ; thus the increase in L/L_{Edd} requires an increase in the gas supply. It is reasonable to conclude that at least a significant part of the gas required originates in the line-emitting clouds. Thus, as L/L_{Edd} increases, the total mass of line-emitting clumpy gas increases. Combining the above two factors, *the distribution of the line-emitting clouds is ultimately regulated by L/L_{Edd} .*

Note that the strength of high-ionization lines (e.g., C IV) and optically thick lines (e.g., Ly α and Mg II) is roughly proportional to the summed area of the illuminated surface of line-emitting clouds while that of low-ionization, optically thin lines (e.g., narrow-line and broad-line optical Fe II) is roughly proportional to the summed volume of the clouds. Thus, generally, the EW of high-ionization lines and optically thick lines correlates negatively with L/L_{Edd} whereas that of low-ionization, optically thin lines correlates positively. The exact $\log\text{EW}-\log L/L_{\text{Edd}}$ slope for a particular emission line depends on both the cloud distribution and the atomic physics — the latter is the very physics underlying the LOC model.

References

- Baldwin, J. A. 1977, *ApJ*, 214, 679
Baldwin, J. A., *et al.* 1995, *ApJ*, 455, 119
Baskin, A. & Laor, A. 2004, *MNRAS*, 350, L31
Boroson, T. A & Green, R. F. 1992, *ApJS*, 80, 109
Dong, X.-B., Wang, T., Wang, J., Fan, X., Wang, H., Zhou, H., & Yuan, W. 2009a, *ApJ*, 703, L1
Dong, X., *et al.* 2009b, submitted to *ApJ* [arXiv: 0903.5020]
Dong, X., *et al.* 2009c, in *The Origin and Physical Mechanism of the Ensemble Baldwin Effect*, ed. W. Wang, Z. Yang, & Z. Chen (San Francisco: ASP), p. 83
Korista, K. T. 1999 in *Quasars and Cosmology*, (San Francisco: ASP), p. 165