



Association between dietary patterns and glycaemic control in a middle-aged Chinese population

Xiao-Ming Shen¹, Long Shu², Yi-Qian Huang³, Xiao-Yan Zhang², Pei-Fen Zheng^{2,3} and Qin Zhu^{2,3,*}

¹Department of Endocrinology, The No.1 People's Hospital of Pinghu, Pinghu, Zhejiang, The People's Republic of China: ²Department of Nutrition, Zhejiang Hospital, Lingyin Road Number 12, Xihu District, Hangzhou 310013, Zhejiang, The People's Republic of China: ³Department of Digestion, Zhejiang Hospital, Lingyin Road Number 12, Xihu District, Hangzhou 310013, Zhejiang, The People's Republic of China

Submitted 25 February 2021: Final revision received 16 August 2021: Accepted 6 September 2021

Abstract

Objective: The present work was performed to analyse the association of dietary patterns with glycaemic control (Hb A1c < 7 %) in a large group of Chinese adults aged between 45 and 59 years.

Design: Habitual dietary intakes in the preceding 12 months were assessed by well-trained interviewers using a validated semi-quantitative FFQ. Factor analysis with principal component method was used to obtain the dietary patterns, and the associations between dietary patterns and glycaemic control were determined using multivariable logistic regression models. Poor glycaemic control was defined as HbA1c \geq 7.0.

Setting: Despite decades of research, data on the relationship between dietary patterns and glycaemic control (HbA1c < 7 %) in China are sparse.

Participants: A total of 1739 participants aged 45–59 years from Hangzhou were included in the final analysis.

Results: Three dietary patterns were ascertained and labelled as traditional southern Chinese, Western and grains-vegetables patterns. After controlling of the possible confounders, participants in the highest quartile of Western pattern scores had greater OR for HbA1c \geq 7.0 (OR = 1.05; (95 % CI 1.000, 1.095); $P = 0.048$) than did those in the lowest quartile. Compared with those in the lowest quartile of grains-vegetables pattern, participants in the highest quartile had lower OR for HbA1c \geq 7.0 (OR = 0.82; (95 % CI 0.720, 0.949); $P = 0.038$). Besides, no significant relationship between the traditional southern Chinese pattern and HbA1c \geq 7.0 was observed ($P > 0.05$).

Conclusions: This study indicated that the Western pattern was associated with a higher risk, and the grains-vegetables pattern was associated with a lower risk for HbA1c \geq 7.0. Future prospective studies are needed to confirm our findings.

Keywords
Dietary patterns
Glycaemic control
Factor analysis
Cross-sectional study
China

Diabetes mellitus (DM) is a global public health issue, with more than 400 million occurring annually worldwide⁽¹⁾. According to the Global Burden of Disease report, DM was recognised as one of the leading causes of years of life lost and has an important effect on Asian countries⁽²⁾. In China, the overall prevalence of diabetes has increased markedly in the last 10 years, with approximately 114.4 million adults having diabetes and 490 million having pre-diabetes⁽³⁾. Obviously, the high prevalence of diabetes carried considerable economic and health burden to the whole Chinese society⁽⁴⁾. It is well known that DM is

considered as a complex and multifactorial chronic disease that may be influenced by multiple factors, including the family history of DM, genetic factors, sedentary lifestyle, obesity, cigarette smoking and dietary factors^(5,6). Therefore, preventing hyperglycaemia and lowering diabetes risk may be effective for the prevention of diabetes in Chinese adults.

Previous studies have suggested that diet and lifestyle modification are modifiable risk factors for the prevention of DM⁽⁷⁾. Traditionally, nutrition research has been focusing on single nutrients or specific foods in isolation.

*Corresponding author. Email 13105714487@163.com

© The Author(s), 2021. Published by Cambridge University Press on behalf of The Nutrition Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike licence (<https://creativecommons.org/licenses/by-nc-sa/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the same Creative Commons licence is included and the original work is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use.

Numerous studies have investigated the association between the consumption of individual nutrients or foods and food groups and the risk of DM^(8–15). However, given the complexity of diets and the potential interactions between food components, studying individual nutrients or foods cannot accurately and comprehensively reveal the effect of daily diets on DM⁽¹⁶⁾. Meanwhile, in daily life, people do not eat isolated nutrients but consume meals containing many combinations of different foods and nutrients⁽¹⁷⁾. In this context, dietary pattern analysis has emerged as an alternative, holistic approach to study the relationship between overall diet and related health consequences, taking into account the complexity of overall diet and potentially facilitate nutritional recommendations⁽¹⁸⁾.

Over the last couple of decades, a rapid nutritional transition characterised by higher intake of sugar and animal-source foods and lower intake of cereals and vegetables is the major contributor to the epidemic of chronic diseases in China⁽¹⁹⁾. Although there is compelling evidence on the associations between dietary patterns and non-communicable diseases, e.g. obesity, hypertension and metabolic syndrome in the Chinese population^(20–23), little is known about the role of dietary patterns in glycaemic control^(24–26). Heretofore, only one study has investigated the associations between dietary patterns and glycated Hb A1c (HbA1c) in the Chinese population⁽²⁶⁾. Also, to our knowledge, no previous studies have examined the impact of empirical dietary patterns on glycaemic control among middle-aged people in China. Against this background, the purpose of this article was to ascertain dietary patterns using factor analysis and subsequently to evaluate the associations between dietary patterns and glycaemic control (HbA1c < 7.0%) in a middle-aged Chinese population.

Subjects and methods

Study population

We used the data set from the Hangzhou Nutrition and Health Survey in China, collected from June 2015 to December 2016. A detailed description of this study has been published previously⁽¹⁶⁾. In brief, 2437 participants aged 45–59 years were recruited when having their annual health examinations at the Medical Center for Physical Examination, 3A Hospital, where participants were interviewed face-to-face by well-trained interviewers using written questionnaire. A flow chart detailing the process of study participants' selection is presented in Fig. 1. Among the eligible participants, we excluded 193 participants with self-reported having history of CVD, cancer or stroke (because of these possible disease-related changes in dietary habits), thirty-six participants with incomplete anthropometric information, 138 participants with the missing or incomplete dietary information in their questionnaires, 275 participants with self-reported having

diabetes or taking medications (insulin or oral hypoglycaemic agents), forty-two participants who did not provide the HbA1c measurements and fourteen participants with implausible energy intakes (< 600 or > 4000 kcal/d). After the mentioned exclusions, in total, 1739 participants (914 men, 825 women) were included in the final analysis. Written informed consent was obtained from all subjects.

Dietary assessment

We used a semi-quantitative FFQ to assess habitual food intake from the preceding year. This FFQ composing of 138 food items was designed to investigate the dietary intake in the middle-aged Chinese population⁽¹⁶⁾. To evaluate the validity and reliability of this FFQ, a pilot survey in the same population by comparison with three 24-h dietary recalls before the study was performed. The results showed that the correlation coefficients between this FFQ and averages from 24-h dietary recalls ranged between 0.43 and 0.67 for major food groups. In short, these data revealed that this FFQ provided reasonably valid measurements of dietary intake. Participants were asked to recall the frequency of each food item during the previous a year and the estimated portion size, using local weight units (1 Jin = 500 g, 1 Liang = 50 g) or natural units (cups). For instance, the frequency response for food consumption included nine categories, as follows: never, < 1 time a month, 1 to 3 times a month, 1 to 2 times a week, 3 to 4 times a week, 5 to 6 times a week, 1 time a day, 2 times a day and 3 times a day. Standard tableware and food models were provided to help the participants to quantify food consumption accurately. Then, they were converted into an average daily consumption in our analyses. Nutrients intakes (i.e. carbohydrate, protein and fat) based on the food consumption data were computed using the Chinese Food Composition Table⁽²⁷⁾.

Identification of dietary patterns

We classified 138 food items from this FFQ into thirty predefined food groups (see online supplemental Table S1) based on culinary usage and common nutrient composition in a middle-aged Chinese population⁽¹⁶⁾. Before beginning the factor analysis, the Bartlett's test of sphericity and Kaiser–Meyer–Olkin (KMO) measure of sample adequacy were used to identify the sample adequacy for factor analysis. Then, we generated dietary patterns by factor analysis (principal components) on the basis of thirty predefined food groups by means of orthogonal transformation (varimax rotation), maintaining the uncorrelated nature of the factors and greater interpretability. The eigenvalue, scree plot and factor interpretability were applied to decide which factors remained⁽²⁸⁾. After evaluating the eigenvalues, the scree plot test and interpretability, factors with an eigenvalues ≥ 2.0 were retained. The factor score for each pattern was calculated by summing intakes of food groups weighted by their factor loadings, and each




Fig. 1 Flow chart of article screening and selection process

participant received a factor score for each identified pattern. In our study, the predefined factor groups with an absolute value factor loading of 0.4 were considered the main factor affecting the pattern. Labelling of dietary patterns was based on the interpretation of foods with high factor loadings for each dietary pattern. Lastly, factor scores were categorised into four equal groups by using quartile cutoffs. Quartile 1 (Q1) represents the lowest adherence, and Quartile 4 (Q4) represents the highest adherence to a certain dietary pattern.

In our previous studies^(16,29), three major dietary patterns were extracted and labelled as the ‘traditional southern Chinese’ (high consumption of refined grains, vegetables, fruits, pickled vegetables, fish and shrimp, bacon and salted fish, salted and preserved eggs, milk, soya bean and its products, miscellaneous bean, fats and drinks), the ‘Western’ pattern (high consumption of red meats, poultry and organs, processed and cooked meat, eggs,

seafood, cheese, fast foods, snacks, chocolates, alcoholic beverages and coffee) and the ‘grains-vegetables’ pattern (high consumption of whole grains, tubers, vegetables, mushrooms, vegetable oil, nuts, honey and tea).

Assessment of blood pressure

The well-trained nurse measured the participant’s blood pressure using an automatic device (Omron) with the participants in the sitting position after a 5–10 min rest. Then, three blood pressure readings taken 5 min apart were recorded, and the mean of the three readings was considered as the final blood pressure.

Assessment of biomarker

Blood samples were collected from all eligible participants after an overnight fast (at least 8 h) and were stored temporarily at -20°C until assessment assays were performed. The

participants underwent an oral glucose tolerance test using 75 g of glucose. All blood samples were analysed in the Medical Center for Physical Examination, Zhejiang Hospital for fasting plasma glucose, glycosylated Hb (HbA1c) using the Hitachi 7180 automatic biochemical analyser (Hitachi, Tokyo, Japan).

Anthropometric measurements

Weight in light clothes and without shoes was measured with a digital scale to the nearest 0.1 kg, and height was measured without shoes to the nearest 0.1 cm. BMI was calculated as weight (kg) divided by the square of height in meters (m²). Waist circumference was measured to the nearest 0.1 cm at the midpoint between the lower rib and the upper iliac crest, while the participants were standing and hip circumference was measured at the maximum level over light clothing by using an inelastic plastic tape⁽³⁰⁾. All anthropometric measurements were performed by well-trained nurses according to standard procedures.

Assessment of other variables

Data on physical activity were collected using the short form of International Physical Activity Questionnaire and was expressed as metabolic equivalent hours per week (MET-h/week)⁽¹⁶⁾. Subsequently, physical activity level was categorised as light, moderate and heavy⁽³¹⁾. Additional information including smoking status (never, current and former smokers) and education level (primary school or below, middle and high school, junior college or above) was collected with a written questionnaire. Current smokers were defined as those who smoked at least one cigarette/d. Energy intake was assessed through this semi-quantitative FFQ, and results were expressed in kilocalorie per day (kcal/d) and categorised according to quartile.

Definition

Poor glycaemic control was defined as HbA1c \geq 7.0⁽¹⁶⁾.

Statistical analyses

Quartiles were made of three dietary patterns of consumption, and the first quartile (the lowest consumption of the pattern) was compared with the fourth quartile (the highest consumption of the pattern). Data were presented as counts (percentages) for categorical variables and as mean \pm SD for continuous variables. Before data analysis, Shapiro–Wilk test, Q–Q plots and histograms were used to test the normality of distribution. If data are normal distributed variables, we used independent-samples *t* test to compare the significant differences in continuous variables. Otherwise, the Mann–Whitney test was used. Besides, we used the χ^2 test to compare the difference in categorical variables. Multivariate logistic regression analyses were applied to determine the associations of dietary patterns with glycaemic control in crude and adjusted

Table 1 Demographic and clinical characteristics of study participants

Variables	Median (n 1739)	Interquartile range (P25–P75)
Age (years)	50	47–55
Gender, male		
<i>n</i>	914	
%	52.6	
BMI (kg/m ²)	24.3	22.4–26.2
Obesity		
<i>n</i>	324	
%	18.6	
WC (cm)	84	78–90
Men: > 85 cm, women > 80 cm		
<i>n</i>	1046	
%	60.1	
HbA1c	6.3	5.8–7.4
>7.0		
<i>n</i>	970	
%	55.8	
SBP (mmHg)	129	118–138
DBP (mmHg)	80	71–80
Hypertension		
<i>n</i>	498	
%	28.6	
Physical activity		
Light		
<i>n</i>	1376	
%	79.1	
Moderate		
<i>n</i>	275	
%	15.8	
Heavy		
<i>n</i>	88	
%	5.1	
Total energy (kcal/d)	1916	1856–2163

WC, waist circumference; SBP, systolic blood pressure; DBP, diastolic blood pressure.

models. For the multivariate analysis, the model 1 was adjusted for age (years) and sex (male/female); model 2 was further adjusted for educational level (primary school or below, middle and high school, junior college or above), smoking (never, current, former), physical activity (light, moderate and heavy) and BMI (continuous); model 3 was additionally adjusted for total energy intake (kcal/d). All analyses were performed using IBM Statistical Package SPSS (version 23.0, SPSS Inc.), and a 2-sided *P* < 0.05 was set as the threshold for statistical significance.

Results

The demographic and clinical characteristics of study participants are shown in Table 1. This present sample was composed mainly of male (52.6%), with a mean age of 50 years (interquartile range, 47–55). Of the 1739 eligible participants, 18.6% were obesity, 60.1% had abnormal waist circumference, 55.8% lacked glycaemic control (HbA1c > 7%) and 28.6% had hypertension. Total energy consumption was 1916 kcal/d (interquartile range:

Table 2 The characteristics of study participants across quartiles of the main dietary pattern scores

	Traditional southern Chinese pattern score				<i>P</i>	Western pattern score				<i>P</i>	Grains-vegetables pattern score				<i>P</i>
	Q1 (<i>n</i> 434)		Q4 (<i>n</i> 435)			Q1 (<i>n</i> 435)		Q4 (<i>n</i> 435)			Q1 (<i>n</i> 435)		Q4 (<i>n</i> 434)		
	<i>n</i>	%	<i>n</i>	%		<i>n</i>	%	<i>n</i>	%		<i>n</i>	%	<i>n</i>	%	
Age (years)															
Mean	51.3		50.6		0.355	51.6		50.3		<0.05	50.4		51.8		<0.05
SD	4.5		4.7		4.7		4.3		4.6		4.7				
BMI (kg/m ²)					0.108	24.26		25.02		<0.05	24.99		24.08		<0.05
Mean	24.42		24.86		2.68		2.92		2.72		2.64				
SD	2.97		3.08												
WC (cm)					0.120	84.29		87.32		<0.01	86.98		82.80		<0.01
Mean	84.34		85.18		8.75		8.68		9.08		9.00				
SD	9.03		9.11												
WHR					0.602	0.87		0.89		<0.05	0.88		0.86		<0.05
Mean	0.87		0.88		0.07		0.06		0.06		0.08				
SD	0.06		0.06												
FPG (mmol/l)					0.080	5.10		5.84		<0.01	5.77		5.35		<0.01
Mean	5.25		5.34		0.79		0.73		0.81		0.70				
SD	0.63		0.71												
Obesity	71	16.4	62	14.3	0.388	50	11.5	80	18.4	<0.01	72	16.6	47	10.8	<0.001
Hypertension	125	28.8	153	35.2	<0.05	102	23.4	142	32.6	<0.01	130	29.9	106	24.4	0.070
Diabetes	87	20.0	94	21.6	0.570	72	16.6	104	23.9	<0.01	92	21.1	65	15.0	<0.05
Gender					<0.001					<0.05					<0.001
Female	136	31.3	215	49.4		199	45.7	166	38.2		174	40.0	251	57.8	
Male	298	68.7	220	50.6		236	54.3	269	61.8		261	60.0	183	42.2	
Smoking status					0.001					<0.01					<0.01
Never	353	81.3	307	70.6		296	68.1	268	61.6		332	76.3	366	84.3	
Current	66	15.2	107	24.6		94	21.6	134	30.8		76	17.5	44	10.2	
Former	15	3.5	21	4.8		45	10.3	33	7.6		27	6.2	24	5.5	
Education					0.395					<0.05					0.268
Primary school or below	151	34.8	165	37.9		137	31.5	156	35.9		166	38.2	164	37.8	
Middle and high school	237	54.6	234	53.8		267	61.4	230	52.9		224	51.5	238	54.8	
Junior college or above	46	10.6	36	8.3		31	7.1	49	11.3		45	10.3	32	7.4	
The average monthly income/person					0.167					0.093					<0.05
< 2000 (RMB)	183	42.2	203	46.7		194	44.6	177	40.7		182	41.8	211	48.7	
2000–4000 (RMB)	205	47.2	178	40.9		190	43.7	185	42.5		184	42.3	176	40.5	
> 4000 (RMB)	46	10.6	54	12.4		51	11.7	73	16.8		69	15.9	47	10.8	
Physical activity					0.422					0.193					0.096
Light	338	77.9	348	80.0		340	78.2	357	82.1		339	77.9	333	76.7	
Moderate	78	18.0	65	14.9		62	14.3	57	13.1		79	18.2	70	16.1	
Heavy	18	4.1	22	5.1		33	7.6	21	4.8		17	3.9	31	7.2	
Total energy (kcal/d)					<0.01					<0.001					<0.01
Mean	2163.7		1916.5		1930.8		2286.4		2122.5		1902.9				
SD	149.6		162.1		160.7		173.7		156.8		180.6				

WC, waist circumference; WHR, waist:hip ratio; FPG, fasting plasma glucose.

P values for continuous variables (ANOVA) and for categorical variables (χ^2 test).

 Categorical variables are presented as sum and percentages, and continuous variables are presented as mean \pm sd.

1856–2163). The Bartlett's test ($P < 0.0001$) and KMO index (0.798) indicated that the correlation between variables was sufficiently robust to perform factor analysis²¹. Herein, three main dietary patterns were identified, naming the traditional southern Chinese, western and grains-vegetables patterns, which accounted for 10.3%, 8.5% and 6.8% of the dietary intake variance, respectively. The factor-loading matrixes for these dietary patterns are shown in Supplemental Table S2.

The characteristics of the study participants across quartile categories of the main dietary pattern scores are shown in Table 2. Compared with the study participants in the lowest quartile, those in the highest quartile of the traditional southern Chinese pattern were females,

smokers and had higher prevalence of hypertension, lower total energy intake. In contrast, participants who belonged to the highest quartile of the Western pattern were more likely to be younger, males, smokers, with higher BMI, waist circumference, WHR, FBG, educational level, total energy intake and higher prevalence of obesity, hypertension and diabetes than those in the lowest quartile. Besides, in comparison with the participants from the lowest quartile of the grains-vegetables pattern, those in the highest quartile were more likely to be older, females, never-smokers and had lower BMI, waist circumference, WHR, FBG, income and total energy intake and had lower prevalence of obesity, hypertension and diabetes ($P < 0.05$).

Table 3 Multivariable adjusted OR and 95 % CI for glycaemic control (HbA1c < 7.0 %) across the quartile (Q) categories of dietary pattern scores in Zhejiang Province, China

	Traditional southern Chinese pattern score				Western pattern score				Grains-vegetables pattern score						
	Q1	Q4			Q1	Q4			Q1	Q4					
		OR	95 % CI			P	OR	95 % CI		P	OR	95 % CI		P	
Model 1	1.00	1.15	1.014, 1.309		0.030	1.00	1.70	1.323, 2.174		0.000	1.00	0.51	0.320, 0.817		0.000
Model 2	1.00	1.09	0.902, 1.323		0.365	1.00	1.39	1.069, 1.842		0.017	1.00	0.68	0.522, 0.899		0.009
Model 3	1.00	1.08	0.889, 1.322		0.425	1.00	1.05	1.000, 1.095		0.048	1.00	0.82	0.720, 0.947		0.038

Model 1: adjusted for sex and age; model 2: further adjusted for educational level (< high school, high school and > high school), physical activity (MET-h/week), smoking status (never, current and former), hypertension (yes/no) and BMI; model 3: additionally adjusted for total energy intake. Q4: the highest quartile of dietary patterns, Q1: the lowest quartile of dietary patterns (reference).

The association between dietary patterns and glycaemic control by multivariate logistic regression analysis is shown in Table 3. After adjusting for potential confounding variables, study participants in the fourth quartile (Q4, highest consumption) of the Western pattern scores had greater OR for glycaemic control (OR = 1.05; (95 % CI 1.000, 1.095); $P = 0.048$) than did those in the first quartile (Q1, lowest consumption). Compared with those in the first quartile, participants in the fourth quartile of the grains-vegetables pattern scores had a lower OR for glycaemic control (OR = 0.82; (95 % CI 0.720, 0.947); $P = 0.038$). Besides, no significant association was established between the traditional southern Chinese pattern and glycaemic control ($P > 0.05$).

Discussion

In the past three decades, T2DM and its complications have reached epidemic levels, particularly in China. However, reports in the literature on the association of dietary patterns with glycaemic control are scarce. To the authors' knowledge, this is one of the few studies to identify the association of dietary patterns with glycaemic control in a middle-aged Chinese population. In the present study, three main dietary patterns were identified by factor analysis and named as 'traditional Chinese', 'Western' and 'grains-vegetables', respectively. Results from this study showed that the Western pattern was associated with a greater risk for glycaemic control, and the grains-vegetables pattern was associated with a lower risk for glycaemic control. Moreover, no significant association was observed between traditional Chinese and poor glycaemic control.

Driven by rapid economic growth and nutrition transition, China has experienced an increased burden of nutrition-related diseases, e.g. diabetes and obesity⁽³²⁾. Zhejiang province, located in East China, where people's dietary patterns have also undergone the rapid westernisation that changes the traditional pattern to the western pattern⁽³³⁾. Besides, the specifics of the eating habits differ in China depending on the region and cultural practices. People in Zhejiang tend to have a high amount of sweet food

and seafood. Meanwhile, with the rapid development of local online-to-offline food delivery, this dieting trend makes foods more convenient and accessible, influencing the local people's dietary behaviours⁽³²⁾. Compared with the home-made foods, take-outs foods often contain higher oil, salt, unhealthy fat or sugars⁽³⁴⁾. Thus, studying the associations between dietary patterns and glycaemic control in middle-aged adults from Zhejiang Province can be of value.

In the present study, no significant association was observed for the traditional southern Chinese pattern in relation to improved glycaemic control. This finding is inconsistent with a prior research performed in Brazil, which showed that the healthy eating pattern with a high intake of whole carbohydrates, dairy, white meat, fish, fruits and vegetables was significantly associated with reduced fasting plasma glucose, HbA1c and LDL-cholesterol levels in patients with type 2 diabetes⁽³⁵⁾. There are several possible explanations concerning the apparent null finding for this pattern and glycaemic control. First of all, in East China, rice is the staple food of three meals a day. Previous studies showed that higher consumption of rice was associated with an increased risk of diabetes⁽³⁶⁾. Second, pickled vegetables with high salt content have also been associated with higher risk of hypertension, an important risk for diabetes⁽³⁷⁾. On the other hand, constituents of fruits and vegetables, including dietary fibre, antioxidants (e.g. vitamins C, E and folate) and Mg, have been independently associated with reduced risk of diabetes⁽³⁸⁾. Some constituents such as dietary fibre, antioxidants (e.g. vitamins C, E and folate) and magnesium found in fruits and vegetables have been shown to be associated with reduced risk of diabetes⁽³⁸⁾. Jiang *et al.* have found that increasing dietary fibre intake may be an effective approach to reduce the level of HbA1c, improving glycaemic control in Chinese diabetic patients⁽³⁹⁾. Moreover, an emerging body of evidence has demonstrated that antioxidants, such as vitamin C and other carotenoids compounds abundant in fruits and vegetable, are associated with decreased risk of obesity and hypertension, which are the established risk factor for T2DM^(40,41). Thus, these possibilities cannot be excluded in the analysis.



In our study, higher adherence to the Western pattern was positively associated with poor glycaemic control. Similarly, in the study by Ruiz Martinez *et al.*, the significant association between consuming a Western-style diet and HbA1c $\geq 7.0\%$ was observed in Mexican population⁽²⁵⁾. The detrimental effect of this pattern on glycaemic control may be related to the unhealthy foods, e.g. red meat, processed and cooked meat, fast foods, snacks and chocolates. First, a previous study from our group in the sample found that higher consumption of red meat was positively associated with T2DM⁽¹⁶⁾. Alternatively, red meat is rich in haem Fe. Previous study has reported that body Fe overload may promote insulin resistance and increase the risk of T2DM⁽⁴²⁾. Second, processed and cooked meats contain a lot of salt. The aforementioned findings suggest that high salt intake can increase the risk of diabetes⁽³⁷⁾. Third, snack and chocolates with high glycaemic index could lead to an increase in weight gain, which can affect the metabolism of glucose and insulin sensitivity, thus increasing the level of blood glucose⁽⁴³⁾. Likewise, a high consumption of energy and energy/kg of ideal weight has been shown to affect glycaemic control⁽⁴⁴⁾. Finally, fast foods including in the Western pattern has been reported to be associated with an increased risk of diabetes⁽⁴⁵⁾. A cross-sectional study in the United Arab Emirates found that fast-food consumption (>1 time/week) had an OR of 1.83 (95% CI 1.02, 3.3) for poor glycaemic control among diabetic patients⁽⁴⁶⁾.

Higher adherence to the grains-vegetables pattern, composed mainly of whole grains, vegetables and beans, was positively associated with improving glycaemic control. Our results are in agreement with recent a study, which suggest that consuming a healthy pattern was associated with improved glycaemic control⁽²⁵⁾. This favourable association of grains-vegetables pattern with improving glycaemic control may be attributed to this pattern's healthy constituents, for example whole grains, tuber, vegetable and beans. First, dietary fibre intake has been found to be associated with reduced risk of insulin resistance⁽³⁸⁾. Second, most foods (such as whole grains, tubers, vegetables and bean) in this pattern have a low glycaemic load, which has been reported to be associated with a lower risk of T2DM⁽⁴⁶⁾. Third, our finding also indicated that participants who belonged to the fourth quartile of grains-vegetables pattern had a higher level of physical activity, compare with those in the lowest quartile. A recent meta-analysis of observational studies revealed that higher level of physical activity was associated with decreased risk of obesity, a major factor for diabetes⁽⁴⁷⁾.

Notably, in the present work, we used the data set from the Hangzhou Nutrition and Health Survey, designed to investigate the relationship between diet and health consequences. Two previous studies by our groups have also examined the effects of dietary patterns on the risks of T2DM and pre-diabetes, respectively^(16,29). Shu *et al.*⁽¹⁶⁾

identified three major dietary patterns in 1918 participants aged 45–59 years in Hangzhou city, Zhejiang province, in East China. Participants in the highest quartile of the grains-vegetables dietary pattern scores had a lower OR for T2DM (OR = 0.72 (95% CI 0.596, 0.952), $P = 0.04$), compared with those in the lowest quartile. Subsequently, in the same study sample, Shen *et al.*⁽²⁹⁾ found that Western pattern with a high content in meats, fast foods, snacks and chocolates was associated with higher risk of prediabetes. However, in contrast to the mentioned two studies^(16,29), this study possessed its unique features. In the selection process of study participants, we excluded those with diabetes or using insulin or oral hypoglycaemic agents, who may have changed their dietary behaviours by the time when the survey was carried out. The exclusion of study participants helps confirm our hypotheses in a healthy middle-aged population-based setting and minimises the chance of reverse causation. Moreover, our research used HbA1c ≥ 7.0 as poor glycaemic control. In this study, these included participants have not been diagnosed with diabetes or impaired glucose tolerance. Therefore, our findings reinforced the importance of optimising dietary behaviours for a healthy middle-aged population.

Strengths and limitations

Our study has several strengths. First, to our knowledge, this is one of the few studies to assess the association between dietary patterns and glycaemic control in a middle-aged Chinese population. Our findings may be helpful to develop dietary recommendations for the early prevention of hyperglycaemia. Second, our study used the dietary pattern analysis, which considered the complexity of overall diet, facilitating dietary recommendations. Third, data on dietary intake were collected by face-to-face interview using a validated semi-quantitative FFQ. This FFQ enabled us to obtain more reliable information on dietary intake of participants in the past 12 months. Meanwhile, the validity and reproducibility of this FFQ has also been confirmed elsewhere⁽¹⁶⁾. Fourth, we have also adjusted for potential known confounders in multivariate regression models. Nevertheless, some limitations also exist in this study. First, the cross-sectional design of this study does not allow for a causal interpretation between dietary patterns and glycaemic control. Thus, future studies with longitudinal designs are needed to confirm the our findings. Second, though we adjusted for multiple confounding variables, the possibility of residual confounding remains in our study. Third, the dietary pattern analysis is somewhat subjective in decisions on the number of factors extracted, food group methods and the labelling of dietary patterns⁽¹⁷⁾. Finally, it is important to note that the generalizability of our findings may be limited because the sample of this study consisted of middle-aged adults, which were recruited from the municipal area of Hangzhou city, China.

Conclusions

To conclude, the results of our study showed that the Western pattern was associated with an increased risk, whereas grains-vegetables pattern was associated with a decreased risk of poor glycaemic control. Our findings reinforced the importance of optimising dietary behaviours and could be helpful to develop nutrition guide for early IGT prevention in a healthy middle-aged population in China. Given the cross-sectional nature of this study, additional prospective studies are necessary to confirm these findings.

Acknowledgements

Acknowledgements: The authors thank all participants and staffs, in particular from Department of Nutrition, Zhejiang Hospital and Department of Endocrinology, the No.1 People's Hospital of Pinghu for their active cooperation and assistance. We also acknowledge the Medical Center for Physical Examination, Zhejiang Hospital for their important contributions to collection of data in the present study.

Financial support: This study was supported by National Natural Science Foundation of China (grant number: 82004040), Natural Science Foundation of Zhejiang (grant number: LY17H030008), the joint construction of projects by provinces and the ministry of education (grant number: 2014PYA002), Traditional Chinese Medicine Research Project of Zhejiang (grant number: 2020ZB009) and Medical and Health research fund project of Zhejiang Province (grant number: 2017KY190). **Conflicts of interest:** The authors declare that there is no conflict of interest associated with this manuscript. **Authorship:** The authors' responsibilities were as follows: Z.Q. and Z.P.F. designed and conducted this research; S.L. and H.Y.Q. analysed data; Z.Q. and S.X.M. wrote this manuscript; Z.Q. had primary responsibility for the final content of the manuscript and all authors read and approved the final manuscript. **Ethics of human subject participation:** Written informed consent was obtained from all study participants prior to their participation in the survey. This study was conducted according to the guidelines laid down in the Declaration of Helsinki, and all procedures involving research study participants were approved by the Institutional Review Board of the Zhejiang Hospital (2014-KA-1).

Supplementary material

For supplementary material accompanying this paper visit <https://doi.org/10.1017/S1368980021003931>

References

- International Diabetes Federation (2017) *IDF Diabetes Atlas*, 8th ed. Belgium: International Diabetes Federation.
- Lozano R, Naghavi M, Foreman K *et al.* (2012) Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* **380**, 2095–2128.
- Xu Y, Wang L, He J *et al.* (2013) Prevalence and control of diabetes in Chinese adults. *JAMA* **310**, 948–959.
- Wang Z, Li X & Chen M (2018) Socioeconomic factors and inequality in the prevalence and treatment of diabetes among middle-aged and elderly adults in China. *J Diabetes Res* **2018**, 1471808.
- Zhang N, Du SM & Ma GS (2017) Current lifestyle factors that increase risk of T2DM in China. *Eur J Clin Nutr* **71**, 832–838.
- Salas-Salvadó J, Bulló M, Estruch R *et al.* (2014) Prevention of diabetes with Mediterranean diets: a subgroup analysis of a randomized trial. *Ann Intern Med* **160**, 1–10.
- Zheng Y, Ley SH & Hu FB (2018) Global aetiology and epidemiology of type 2 diabetes mellitus and its complications. *Nat Rev Endocrinol* **14**, 88–98.
- Cooper AJ, Sharp SJ, Lentjes MA *et al.* (2012) A prospective study of the association between quantity and variety of fruit and vegetable intake and incident type 2 diabetes. *Diabetes Care* **35**, 1293–1300.
- Geiker NRW, Larsen ML, Dyerberg J *et al.* (2018) Egg consumption, cardiovascular diseases and type 2 diabetes. *Eur J Clin Nutr* **72**, 44–56.
- Kurotani K, Nanri A, Goto A *et al.* (2013) Red meat consumption is associated with the risk of type 2 diabetes in men but not in women: a Japan public health center-based prospective study. *Br J Nutr* **110**, 1910–1918.
- Djoussé L, Petrone AB, Hickson DA *et al.* (2016) Egg consumption and risk of type 2 diabetes among African Americans: the Jackson Heart Study. *Clin Nutr* **35**, 679–684.
- Forouhi NG, Koulman A, Sharp SJ *et al.* (2014) Differences in the prospective association between individual plasma phospholipid saturated fatty acids and incident type 2 diabetes: the EPIC-InterAct case-cohort study. *Lancet Diabetes Endocrinol* **2**, 810–818.
- Soriguer F, Colomo N, Olveira G *et al.* (2013) White rice consumption and risk of type 2 diabetes. *Clin Nutr* **32**, 481–484.
- Mohan V, Radhika G, Sathya RM *et al.* (2009) Dietary carbohydrates, glycaemic load, food groups and newly detected type 2 diabetes among urban Asian Indian population in Chennai, India (Chennai urban rural epidemiology study 59). *Br J Nutr* **102**, 1498–1506.
- Ley SH, Hamdy O, Mohan V *et al.* (2014) Prevention and management of type 2 diabetes: dietary components and nutritional strategies. *Lancet* **383**, 1999–2007.
- Shu L, Shen XM, Li C *et al.* (2017) Dietary patterns are associated with type 2 diabetes mellitus among middle-aged adults in Zhejiang Province, China. *Nutr J* **16**, 81.
- Xu SS, Hua J, Huang YQ *et al.* (2020) Association between dietary patterns and chronic kidney disease in a middle-aged Chinese population. *Public Health Nutr* **23**, 1058–1066.
- Hu FB (2002) Dietary pattern analysis: a new direction in nutritional epidemiology. *Curr Opin Lipidol* **13**, 3–9.
- Huang L, Wang Z, Wang H *et al.* (2021) Nutrition transition and related health challenges over decades in China. *Eur J Clin Nutr* **75**, 247–252.
- Yuan YQ, Li F, Meng P *et al.* (2016) Gender difference on the association between dietary patterns and obesity in Chinese middle-aged and elderly populations. *Nutrients* **8**, 448.
- Zheng PF, Shu L, Zhang XY *et al.* (2016) Association between dietary patterns and the risk of hypertension among Chinese: a cross-sectional study. *Nutrients* **8**, 239.
- Zhu WL, Guan Y, Xu CZ *et al.* (2020) Influence of dietary patterns on type 2 diabetes mellitus in local residents aged 40 years and above in Songjiang district, Shanghai. *Zhonghua Liu Xing Bing Xue Za Zhi* **41**, 508–513.



23. Wei ZY, Liu JJ, Zhan XM *et al.* (2018) Dietary patterns and the risk of metabolic syndrome in Chinese adults: a population-based cross-sectional study. *Public Health Nutr* **21**, 2409–2416.
24. Hakeem R, Ahmedani MY, Alvi SF *et al.* (2014) Dietary patterns and glycemic control and compliance to dietary advice among fasting patients with diabetes during Ramadan. *Diabetes Care* **37**, e47–e48.
25. Martínez MLR, Gómez-Díaz RA, González ALV *et al.* (2020) Association between glycemic control and dietary patterns in patients with type 2 diabetes in a Mexican institute. *Nutrition* **78**, 10901.
26. Jaacks LM, Crandell J, Mendez MA *et al.* (2015) Dietary patterns associated with HbA1c and LDL cholesterol among individuals with type 1 diabetes in China. *J Diabetes Complications* **29**, 343–349.
27. Yang Y (2005) *Chinese Food Composition Table 2004*. Beijing: Peking University Medical Press.
28. Wang CJ, Yang TF, Wang GS *et al.* (2018) Association between dietary patterns and depressive symptoms among middle-aged adults in China in 2016–2017. *Psychiatry Res* **260**, 123–129.
29. Shen XM, Huang YQ, Zhang XY *et al.* (2020) Association between dietary patterns and prediabetes risk in a middle-aged Chinese population. *Nutr J* **19**, 77.
30. Yang CQ, Shu L, Wang S *et al.* (2015) Dietary patterns modulate the risk of non-alcoholic fatty liver disease in Chinese adults. *Nutrients* **7**, 4778–4791.
31. Craig CL, Marshall AL, Sjöström M *et al.* (2003) International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* **35**, 1381–1395.
32. Zhu S (2021) Development of nutritional studies in China. *Eur J Clin Nutr* **75**, 230–231.
33. Shu L, Zheng PF, Zhang XY *et al.* (2015) Association between dietary patterns and the indicators of obesity among Chinese: a cross-sectional study. *Nutrients* **7**, 7995–8009.
34. Jaworowska A, Blackham T, Davies IG *et al.* (2013) Nutritional challenges and health implications of takeaway and fast food. *Nutr Rev* **71**, 310–318.
35. Sarmento RA, Antonio JP, de Miranda IL *et al.* (2017) Eating patterns and health outcomes in patients with type 2 diabetes. *J Endocr Soc* **2**, 42–52.
36. Hu EA, Pan A, Malik V *et al.* (2012) White rice consumption and risk of type 2 diabetes: meta-analysis and systematic review. *BMJ* **344**, e1454.
37. Cryer MJ, Horani T & DiPette DJ (2016) Diabetes and hypertension: a comparative review of current guidelines. *J Clin Hypertens* **18**, 95–100.
38. Greenhill C (2020) Dietary factors in the risk of T2DM. *Nat Rev Endocrinol* **16**, 537.
39. Jiang J, Qiu H, Zhao G *et al.* (2012) Dietary fiber intake is associated with HbA1c level among prevalent patients with type 2 diabetes in Pudong new area of Shanghai, China. *PLoS One* **7**, e46552.
40. Garcia-Diaz DF, Lopez-Legarrea P, Quintero P *et al.* (2014) Vitamin C in the treatment and/or prevention of obesity. *J Nutr Sci Vitaminol* **60**, 367–379.
41. Nuñez-Cordoba JM, Alonso A, Beunza JJ *et al.* (2009) Role of vegetables and fruits in Mediterranean diets to prevent hypertension. *Eur J Clin Nutr* **63**, 605–612.
42. Jiang R, Ma J, Ascherio A *et al.* (2004) Dietary iron intake and blood donations in relation to risk of type 2 diabetes in men: a prospective cohort study. *Am J Clin Nutr* **79**, 70–75.
43. Al-Khudairy L, Stranges S, Kumar S *et al.* (2013) Dietary factors and type 2 diabetes in the Middle East: what is the evidence for an association? A systematic review. *Nutrients* **5**, 3871–3897.
44. Evert AB, Boucher JL, Cypress M *et al.* (2014) Nutrition therapy recommendations for the management of adults with diabetes. *Diabetes Care* **37**, Suppl. 1, S120–S143.
45. Pereira MA, Kartashov AI, Ebbeling CB *et al.* (2005) Fast-food habits, weight gain, and insulin resistance (the CARDIA study): 15-year prospective analysis. *Lancet* **365**, 36–42.
46. Sugiyama M, Tang AC, Wakaki Y *et al.* (2003) Glycemic index of single and mixed meal foods among common Japanese foods with white rice as a reference food. *Eur J Clin Nutr* **57**, 743–752.
47. Sharkey T, Whatnall MC, Hutchesson MJ *et al.* (2020) Effectiveness of gender-targeted *v.* gender-neutral interventions aimed at improving dietary intake, physical activity and/or overweight/obesity in young adults (aged 17–35 years): a systematic review and meta-analysis. *Nutr J* **19**, 78.