

# The role of rumination in illness trajectories in youth: linking trans-diagnostic processes with clinical staging models

A. B. Grierson<sup>1</sup>, I. B. Hickie<sup>1</sup>, S. L. Naismith<sup>2</sup> and J. Scott<sup>3\*</sup>

<sup>1</sup>Brain and Mind Centre, The University of Sydney, Sydney, Australia

<sup>2</sup>Charles Perkins Centre & Brain and Mind Centre, The University of Sydney, Sydney, Australia

<sup>3</sup>Academic Psychiatry, Institute of Neuroscience, Newcastle University, Newcastle upon Tyne, UK

Research in developmental psychopathology and clinical staging models has increasingly sought to identify trans-diagnostic biomarkers or neurocognitive deficits that may play a role in the onset and trajectory of mental disorders and could represent modifiable treatment targets. Less attention has been directed at the potential role of cognitive-emotional regulation processes such as ruminative response style. Maladaptive rumination (toxic brooding) is a known mediator of the association between gender and internalizing disorders in adolescents and is increased in individuals with a history of early adversity. Furthermore, rumination shows moderate levels of genetic heritability and is linked to abnormalities in neural networks associated with emotional regulation and executive functioning. This review explores the potential role of rumination in exacerbating the symptoms of alcohol and substance misuse, and bipolar and psychotic disorders during the peak age range for illness onset. Evidence shows that rumination not only amplifies levels of distress and suicidal ideation, but also extends physiological responses to stress, which may partly explain the high prevalence of physical and mental co-morbidity in youth presenting to mental health services. In summary, the normative developmental trajectory of rumination and its role in the evolution of mental disorders and physical illness demonstrates that rumination presents a detectable, modifiable trans-diagnostic risk factor in youth.

Received 28 December 2015; Revised 4 May 2016; Accepted 12 May 2016; First published online 29 June 2016

**Key words:** Brooding, clinical staging, mental disorders, rumination, trans-diagnostic, youth.

## Introduction

Developmental psychopathology explores illness trajectories from a life-course perspective, and as such it may be especially helpful for understanding the emergence of mental disorders in youth (Scott *et al.* 2013). In young people, a particular challenge is to identify deviations in basic psychological and/or biological processes that can explain why the initial clinical presentation may demonstrate concurrent mental, physical, and alcohol/substance misuse disorders and/or the longitudinal phenomenology may show heterotypic continuity over time (McGorry *et al.* 2006; Hickie *et al.* 2013; Scott *et al.* 2013).

Attempts to understand evolving patterns of signs and symptoms have increasingly focused on the application of dimensional approaches such as *clinical staging* and the *Research Domain Criteria (RDoC)* initiative (Insel *et al.* 2010; Sanislow *et al.* 2010; Hickie *et al.* 2013; McGorry *et al.* 2014). Both approaches highlight

the need to consider psychopathology on a continuum (from normal experiences through to established clinical disorders) and emphasize that our understanding of liability and co-morbidity is more likely to be improved by examining putative underlying mechanisms that cut across current heterogeneous disorder categories (Baskin-Sommers & Foti, 2015). Implicit in these frameworks is the notion that identifying trans-diagnostic processes that translate into ‘modifiable factors’ is a critical step in the development of precision or personalized medicine for mental disorders (Insel *et al.* 2010; McGorry *et al.* 2014), and moreover, that clinical advances will only really be achieved by greater integration of behavioural neuroscience into the study of psychopathology (Sanislow *et al.* 2010). However, research to date has predominantly focused on ‘biomarkers’ (measurable biological characteristics) and/or neurocognition (‘cold circuits’) (McGorry *et al.* 2014). There have been fewer reviews that synthesize data on ‘hot cognitions’ (Simon *et al.* 2015), such as cognitive processing styles involved in emotional regulation.

Given the lack of reviews that focus on cognitive-emotional processes that may act as a trans-diagnostic risk factor in young people with emerging severe mental disorders, we explore whether a particular ‘coping

\* Address for correspondence. Professor J. Scott, Academic Psychiatry, Institute of Neuroscience, Wolfson Unit, Newcastle University, Newcastle upon Tyne, NE4 6BE, UK.  
(Email: jan.scott@newcastle.ac.uk)

This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

style', namely rumination (and particularly 'toxic brooding'), can be considered as a specific age- and stage-related mechanism for psychopathology in youth. Whilst rumination has been studied extensively, we particularly focus on its evolution and its role in shaping emotional responses to stress in childhood, adolescence and early adulthood, including the peak age of onset period for psychosis, mood disorders and alcohol/substance misuse (between the ages of about 15 and 25 years) (Kessler *et al.* 2005; Jones, 2013) and whether rumination is a mechanism that can also help explain why many disorders in young people show heterotypic continuity over time and related phenomena such as multi-finality and pluri-potentiality (Ehring & Watkins, 2008).

The paper begins by briefly discussing cognitive-emotional regulation and the concept of coping response style, and then reviews rumination from the perspective of theories about its evolution, its normal developmental trajectory and its role in the evolution of high prevalence disorders (such as anxiety and depression) that are often the early stage clinical phenotypes that co-occur with severe mental disorders in adolescence and early adulthood. Next, the paper explores the emerging evidence for links between rumination and alcohol and substance misuse, and bipolar and psychotic disorders. Finally, we briefly highlight the neurobiological underpinnings of rumination and how its putative role in physiological responses to stress may offer insights into the high prevalence of physical and mental co-morbidities in young people.

### Brief overview of cognitive-emotional regulation and rumination

Cognitive-emotional regulation (CER) encompasses any attempt to understand and shape the emotions experienced and how these are expressed (Gross & Munoz, 1995). This regulation can be implicit (automatic, unconscious) or explicit (deliberate, requiring conscious effort) (Gyurak *et al.* 2011). Conscious, effortful cognitive processes can be adaptive or maladaptive and include strategies such as reappraisal and rumination (Garnefski & Kraaij, 2007).

The most widely researched example of CER is the Response Styles Theory (RST; Nolen-Hoeksema, 1987). RST focuses on individual styles of processing emotion that emerge in reaction to stressors and identifies four main coping responses: distraction, problem solving, risk taking, and rumination (see Table 1). Problem solving and distraction are considered to be adaptive response styles as they can positively impact on outcomes and provide opportunities for positive reinforcement, which increases future positive emotions and decreases negative affect (Abela *et al.* 2007).

Risk-taking is considered maladaptive as, although it may temporarily offer distraction from or avoidance of difficult situations or feelings, the 'downstream' effects are more often negative (Nolen-Hoeksema, 1987; Nolen-Hoeksema *et al.* 2008).

Rumination can be viewed as a stable individual trait (Smith & Alloy, 2009) characterized by 'the tendency to repetitively analyse one's problems, concerns and feelings of distress without taking actions to make positive changes' (Nolen-Hoeksema, 1991, p. 569). Rumination appears to be a multifaceted construct, having adaptive and maladaptive components, often referred to as 'reflective pondering' and 'toxic brooding', respectively (Treyner *et al.* 2003). Treyner *et al.* (2003) define reflection as a purposeful turning inward to engage in cognitive problem solving to alleviate one's distress whilst brooding is defined as 'passive comparison of one's current situation with some unachieved standard. Brooding is associated with the maintenance and exacerbation of negative mood states such as depression and dysphoria (Verhaeghan *et al.* 2014), and negative cognitive biases such as attentional engagement with negatively valenced stimuli (Joormann, 2004).

Several models of rumination confirm that it acts to amplify and prolong a prevailing mood state (see Smith & Alloy, 2009), increasing the likelihood that dysphoria will progress to depression, and can increase anxiety in individuals faced with threatening situations (Nolen-Hoeksema & Watkins, 2011). Negative cycles of rumination and dysphoria make it increasingly likely that an individual will focus on prior negative thoughts and experiences, which further impairs instrumental behaviour and effective problem solving (Brosschot *et al.* 2006; Nolen-Hoeksema & Watkins, 2011). In contrast, there is emerging evidence that positive self-focused attention (positive reflection or 'basking') can be associated with lower levels of anxiety or negative affect (Seegerstrom *et al.* 2003; Feldman *et al.* 2008; Raes *et al.* 2012; Hou & Ng, 2014).

How ruminative thought processes, such as toxic brooding, emerge and the underpinning mechanisms have been addressed by several research groups. The most prominent models of rumination, like RST, propose that rumination emerges in reaction to emotions such as negative affect and sadness, and that it is the styles of processing and reacting to emotions that lead to adverse psychological and physical health consequences (Nolen-Hoeksema, 1991; Conway *et al.* 2000). Other models, such as the 'goal progress theory' or 'goal conflict theory' propose that rumination is a response to behavioural inhibition and failure to progress satisfactorily towards a goal, or inaction and increased negative cognitive rehearsal associated with conflictual strivings, rather than a reaction to mood state *per se* (Emmons & King, 1988; Martin *et al.*

**Table 1.** Key elements of cognitive emotion regulation and response styles theory

Cognitive emotional regulation (Gross & Munoz, 1995; Gyurak <i>et al.</i> 2011)		
Emotion regulation	Process	Examples
Explicit	Deliberate and conscious, and demands some level of monitoring during implementation. It is performed with insight and awareness	A situation is selected, modified, attended to, and appraised to yield a certain set of emotional responses. This is conducted in accordance with immediate and long term goals
Implicit	Automatic and unconscious, and thus requires little effort. It is performed without monitoring, insight, or comprehension	Like explicit emotion regulation, a situation is selected, modified, attended to, and appraised to produce a set of emotional responses. However this is done without cognitive engagement in the emotion regulation process
Response style theory (Nolen-Hoeksema, 1987, 1991; Abela <i>et al.</i> 2007)		
Response style	Cognitions/behaviours	
Rumination	Behaviours and thoughts that focus attention on one's depressive symptoms and on the implication of these symptoms	<ul style="list-style-type: none"> <li>• Symptom focused cognitions</li> <li>• Isolating behaviours</li> </ul>
Problem solving	Movement toward a desired end state by active engagement in strategizing to overcome an issue	<ul style="list-style-type: none"> <li>• Deliberate cognitions</li> <li>• Suppression of distracting thoughts</li> <li>• Goal-directed behaviours</li> </ul>
Distraction	Engaging in pleasant, engrossing, and positively reinforcing activities to divert attention from one's depressive symptoms	<ul style="list-style-type: none"> <li>• Suppression of self-contemplation</li> <li>• Active and pleasurable behaviours</li> </ul>
Risk taking	Behaviours that have the potential to be harmful or dangerous, but that may allow the individual to avoid engaging with painful or intense emotions	<ul style="list-style-type: none"> <li>• Cognitive reappraisal of fear</li> <li>• Positive outcome expectancies</li> <li>• Active and instantly gratifying behaviours</li> </ul>

1993). It is suggested that failure to progress toward a goal and act on competing desires inaugurate ruminative thinking, which then leads to aversive psychological consequences such as mental disorders (Jones *et al.* 2009; Einstein, 2014). There is evidence that individuals who fail to progress toward goal states have lower wellbeing (Emmons & King, 1988), that high levels of goal failure combined with maladaptive rumination (brooding) are associated with more depressive symptoms (Jones *et al.* 2009), and that rumination mediates the link between goal attainment and positive affect (McIntosh *et al.* 1995).

Intolerance of uncertainty (IU), defined as a cognitive bias that affects how a person interprets and responds to uncertain situations (Dugas *et al.* 2004) is another mechanism proposed to underpin ruminative thinking (de Jong-Meyer *et al.* 2009; Yook *et al.* 2010; Einstein, 2014). It is suggested that individuals with IU are more sensitive to stress and use maladaptive coping strategies, such as dysfunctional cognitive processes or safety behaviours such as rumination, in order to reduce uncertainty, which may amplify

affective states (McIntosh & Martin, 1992). Studies of IU and rumination demonstrate that rumination completely mediates the relationship between IU and depression (Yook *et al.* 2010; Liao & Wei, 2011), but other cognitive biases such as worry did not show a mediation effect (Yook *et al.* 2010).

### Developmental trajectory of rumination

This section briefly examines evidence for the development of rumination over the life span, selecting three key topics to discuss: age, gender and early environment (including exposure to adversity).

#### Age and gender

Rumination involves a process of cognitive appraisal and so requires some capacity for basic abstract thinking and formal operational thought, which usually become apparent during the transition from mid-childhood to early adolescence (Piaget, 1977; Cole *et al.* 2006). Population-based studies confirm that

children and pre-adolescents show significantly lower levels of rumination than adolescents (Hampel & Petermann, 2005; Thompson *et al.* 2010). Interestingly, Sütterlin *et al.* (2012) showed a decline in rumination from early adulthood onwards, where the highest levels of 'brooding' occurred in individuals aged <25 years. Explanations of the age trajectory of rumination concentrate on the potential long-term impact of childhood coping skills development and experiences. For example, some investigators have suggested that a peak in toxic brooding during adolescence is linked to difficulties in coping with increased levels of normative stressors that occur in this age range especially in those who failed to acquire sufficient adaptive problem-solving skills at an earlier age (Compas, 1987).

Gender differences in rumination also emerge during adolescence, with levels increasing in a linear fashion for girls from 12 to 15 years of age onwards, but with limited increases in rumination in males during the post-pubertal period (Tamres Janicki & Helgeson, 2002; Jose & Ratcliffe, 2004; Jose & Brown, 2008). There are a number of theories as to why females may ruminate more than males, including gender-stereotyped coping style (Cox *et al.* 2010) and differences in self-perceptions of emotionality and the stressfulness of the environment (Rudolph & Hammen, 1999; Gohm, 2003; Hyde *et al.* 2008; Nolen-Hoeksema, 2012). Against the latter, a meta-analysis by Tamres *et al.* (2002) found that gender differences in rumination were independent of the appraisal of stressors. Interestingly, work on 'pubertal timing' suggests that early puberty predicted higher levels of interpersonal dependent events and enhanced the risk of dysphoric symptoms in boys and girls who had a more negative cognitive style and a lower level of emotional clarity (Hamilton *et al.* 2014a, b).

### *Early environment*

Some research indicates that future risk of becoming a 'ruminator' may be associated with early adversity, and rumination can be a learnt response style developed via parent modelling and communication (Hankin *et al.* 2009; Cox *et al.* 2010). Nolen-Hoeksema (1998) proposed that children who have little perceived control over their environment might be especially prone to becoming ruminators in adolescence. For example, pre-school exposure to 'negative-submissive family expressivity' has been shown to predict rumination in adolescence (Hilt *et al.* 2012), whilst exposure to 'over-controlling' parenting can undermine the child's sense of self-efficacy or mastery and contribute to the development of maladaptive coping strategies (Parker, 1983; Blatt & Homann, 1992) and a greater tendency to ruminate (Spasojevic & Alloy, 2002; Manfredi *et al.* 2011).


Neglect or abuse in childhood may shape a child's response to their environment, e.g. being vigilant to signs of threat and becoming passive to avoid aggravation in such situations, paying greater attention to negative stimuli, etc. (Macleod *et al.* 2002). In such circumstances, a child who feels helpless and socially and emotionally isolated may show reduced reliance on externally-oriented problem solving and an increased likelihood of engaging in solitary coping styles such as rumination (Browne & Finkelhor, 1986; Nolen-Hoeksema *et al.* 1994). Moreover, Pearson *et al.* suggested that childhood abuse increases the development of anxious attachments, which have been found to prospectively predict rumination (Pearson *et al.* 2011).

Several studies offer support for the association between early adversity and rumination. For example, in non-clinical samples of college students, individuals with a history of childhood sexual abuse showed higher levels of rumination about sadness and dysphoria than those without such a history (Conway *et al.* 2004). Likewise, clinical studies confirm an association between childhood sexual and emotional abuse and adult ruminative response style (e.g. Spasojevic & Alloy, 2002); however, trends were statistically significant for females but not males (although the latter might be due to limited statistical power). A recent study by O'Mahen *et al.* (2015) examined different experiences of childhood maltreatment in depressed and non-depressed pregnant women and suggested that childhood emotional neglect was related to behavioural avoidance whilst childhood emotional abuse was associated with rumination. Importantly, repeated assessment of nearly 1000 adolescents demonstrated that emotional abuse by parents or peers was an antecedent of negative coping and cognitive styles, and, as elaborated below, brooding mediated the association between abuse and future depressive symptoms (Padilla & Calvete, 2014).

It is important to note that all these adverse experiences may be linked by an overarching issue, namely that they reduce the capacity of a child or adolescent to engage in effective and balanced control of their multiple needs; this in turn may exacerbate feelings of hopelessness, especially if they are living in unsupportive environments (Mansell & Carey, 2013). Furthermore, these elements may increase the long-term risk of developing psychological problems and mental disorders in the future (Powers, 1973).

### **Rumination and psychopathology**

The increased interest in early intervention in youth with severe mental disorders has led to greater awareness of illness trajectories, in particular that preschool temperament or personality traits can be risk markers



**Fig. 1.** Representation of clinical stages of mental disorders beginning with early childhood temperament through to anxiety, depression, and then severe mental disorders (with peak age of onset in late adolescence/early adulthood).

for psychopathology such as internalizing or anxiety problems in childhood, which may precede the onset of depressive symptoms or disorders and be the antecedents of later psychopathology (see Fig. 1). Not all individuals show progression from high prevalence disorders (e.g. anxiety, depression) to the lower prevalence disorders (e.g. psychosis, bipolar disorders), which is a developmental pattern that mimics a clinical staging model (McGorry *et al.* 2006). Clinical staging involves a detailed assessment of where an individual exists on a continuum of disorder progression from stage 0 (an at-risk but asymptomatic state) through to stage IV (late or end-stage disease) (McGorry *et al.* 2006; Hickie *et al.* 2013; McGorry *et al.* 2014), and it is used routinely to describe disease progression in general medicine, e.g. for heart disease or diabetes. If this is a plausible approach for mental disorders, then trans-diagnostic mechanisms should be identifiable across stages and over time both within and between individuals.

### *Temperament, rumination and future psychopathology*

Temperament can be broadly defined as biologically based, relatively stable individual differences in emotions, behaviours, and cognitions (Rothbart & Bates, 2006). Negative emotionality (NE) is an infant temperamental style that is widely associated with psychopathology in adolescence, with high NE individuals experiencing negative emotions more frequently and intensely than their peers (Belsky *et al.* 1996), and are more likely to exhibit depression (Mezulis *et al.* 2011), anxiety (Hudson & Rapee, 2004), and/or substance abuse (Allen & Gabbay, 2013).

A prospective longitudinal study from birth to adolescence, found that rumination significantly mediated the association between NE in infancy and depressive

symptoms at age 15 years (Mezulis *et al.* 2011). Consistent with other studies (e.g. Verstraeten *et al.* 2009), Mezulis *et al.* (2011) noted that the association was stronger among adolescent girls, although a further study by the same group highlighted that high levels of depressive symptoms at age 11 were associated with both early puberty and high levels of NE in boys (Mezulis *et al.* 2014).

Neuroticism is a similar but narrower construct than NE (Eysenck & Eysenck, 1985). Like NE, ruminative response style has been shown to fully or partially mediate the association between neuroticism and anxiety and depression in adolescents and young adults (Muris *et al.* 2005; Kuyken *et al.* 2006; Roelofs *et al.* 2008).

### *Anxiety and depression*

About 50% of individuals with anxiety disorders experience their first symptoms by the age of 11 (Kessler *et al.* 2005). A number of studies demonstrate that childhood anxiety and/or internalizing symptoms may predict adolescent depressive symptoms/episodes (Last *et al.* 1997; Horn & Wuyek, 2010), whilst anxiety disorders in early adolescence significantly predict first onset of a major depressive episode (MDE; Bittner *et al.* 2004). Rates of depressive disorders increase markedly during mid to late adolescence with about 20% of adolescents experiencing a MDE (Lewinsohn *et al.* 1998), and concurrent depression and anxiety occurs significantly more frequently in adolescence compared to childhood (Lamers *et al.* 2011).

Studies suggest that there is a robust relationship between rumination and the development of internalizing problems (such as withdrawal, anxiety and depression) but not externalizing problems (such as aggression, hyperactivity, etc.) in childhood (Garnefski *et al.* 2005). Rumination is also significantly associated

with the development and maintenance of anxiety symptoms in adolescents (e.g. Garnefski *et al.* 2001, 2002; Tan *et al.* 2012; Jose *et al.* 2012; Jose & Weir, 2013), and increases the risk of developing anxiety problems in early adolescence (Tan *et al.* 2012). Furthermore, in a mid-teen sample, social anxiety (a recognized antecedent of severe mental disorders) directly predicted higher levels of rumination over time, especially in females (Jose *et al.* 2012).

Stressful life events and experiences such as bullying are potent risk factors for the development of internalizing symptoms (e.g. McLaughlin *et al.* 2009). In cross-sectional and prospective studies, McLaughlin and colleagues (e.g. McLaughlin & Hatzenbuehler, 2009; McLaughlin & Nolen-Hoeksema, 2012) demonstrated that rumination fully mediated the relationship between stressful life events and anxiety and depression among adolescents, but only partially mediated the association in adults. A large-scale study by Michl *et al.* (2013) confirmed that rumination is a key mediator between stress and anxiety and depressive phenomena in adolescents.

Independently for depression, there is a vast literature that indicates that rumination is significantly associated with depression onset, maintenance, and relapse, especially during adolescence (e.g. Wood *et al.* 1990; Nolen-Hoeksema, 2000; Michalak *et al.* 2012). A meta-analysis demonstrated that the higher levels of rumination in females compared to males significantly mediated the increased rates of depression that becomes apparent post-puberty (Aldao & Nolen-Hoeksema, 2012). Broderick (1998) suggests that adolescents use rumination to deal with family, academic, and peer group problems, which become increasingly more stressful and prominent during the adolescent period. However, as ruminative response style exacerbates rather than reduces the negative outcomes of stressful life events, it increases the likelihood of depressive outcomes (Schwartz & Koenig, 1996; Broderick, 1998; Kraaij *et al.* 2003). Furthermore, a study of an ethnically diverse sample of over 1000 college students demonstrated that individuals who brood in response to negative life events may be vulnerable to thinking about suicide, and that whilst level of suicidal ideation was partly explained by severity of symptoms of depression, it was also directly related to brooding itself (Chan *et al.* 2009).

### *Alcohol and substance misuse*

The prevalence of alcohol use disorders (AUD) and substance use disorders (SUD) in adolescents range from about 5–10% (Merikangas *et al.* 2010; Merikangas & McClair, 2012). Approximately 40% of individuals with an AUD and just below 50% of those with SUD

experienced their first symptoms of harmful use before 19 years (Helzer *et al.* 1991; Dennis *et al.* 2002).

In clinical settings, 70–80% of adolescents with AUD and SUD have a co-morbid mental disorder (Kaminer & Bukstein, 2008), including anxiety, depression, psychosis or bipolar disorders. Studies indicate that rumination increases the risk for harmful alcohol and/or substance use (e.g. Nolen-Hoeksema & Harrell, 2002; Caselli *et al.* 2008; Caselli *et al.* 2010), particularly in female adolescents where rumination predicted the onset of substance abuse (and of bulimia), and predicted a future increase in substance abuse symptoms over a 4-year period (Nolen-Hoeksema *et al.* 2007).

The association between rumination and alcohol and substance misuse may be direct or indirect. For example, some studies of adolescents suggest that depression increases rumination, which in turn leads to increased substance use, as demonstrated by the association between brooding and marijuana use (Adrian *et al.* 2014). Skitch & Abela (2008) showed that ruminative brooding predicted substance abuse for up to 18 weeks following negative events, and that this effect was exacerbated in older adolescents with an existing MDE. Other research supports a direct association (e.g. Willem *et al.* 2011), reporting that, independent of depression, adolescents with lower reflective pondering demonstrated higher drug consumption whilst higher brooding was associated with more problematic substance use.

### *Bipolar disorders (BD)*

The peak age of onset for BD is late adolescence and early adulthood, although most individuals experience one or more MDE, brief hypomania or sub-threshold mania before syndromal episodes of hypomania or mania occur (Duffy *et al.* 2009; Douglas & Scott, 2014). Those individuals with depression who may be at highest risk of transition to BD often show cyclothymic or hypomanic personality traits or have a family history of bipolar disorders (Duffy *et al.* 2009; Geoffroy *et al.* 2013; Bechdolf *et al.* 2014).

There is limited research on rumination and mania in adolescents and young adults, but studies in middle-aged adults or mixed samples of younger and older adults show that rumination is present in both the depressed and (hypo)manic phases of BD, is increased in unaffected first degree relatives compared to healthy controls, and that increased rumination predicts relapse and/or worse course of illness (Johnson *et al.* 2008; Van der Gucht *et al.* 2009; Green *et al.* 2011; Ghaznavi & Deckersbach, 2012).

A study of university students (mean age 22 years) (Thomas & Bentall, 2002), reported that ruminative

response style was positively correlated with scores on the Hypomanic Personality Scale (Eckblad & Chapman, 1986). A second study of undergraduate students (mean age 19 years) by the same group found that rumination was independently associated with both depression and hypomania symptom scores (Knowles *et al.* 2005), and that risk-taking behaviours were also increased in association with rumination, especially in males. Knowles *et al.* (2005) hypothesized that rumination may exacerbate depressive symptoms whilst some individuals make active attempts to avoid negative or intense mood states by engaging in high-risk activities, which may be linked to hypomanic or manic symptoms (Van der Gucht *et al.* 2009; Weiss *et al.* 2015).

Studies of individuals at high risk of BD or the offspring of BD parents demonstrate early evidence of deficits in CER (e.g. Nijjar *et al.* 2014; Van Rheene *et al.* 2015). In a study of response styles, Jones *et al.* (2006) found that offspring with current or past evidence of psychopathology (affected offspring) and unaffected offspring had higher levels of rumination than age- and gender-matched controls. Pavlickova *et al.* (2014) demonstrated that, compared to offspring of healthy parents, affected offspring of bipolar parents showed significantly increased levels of rumination and of hypomanic cognitions compared to unaffected offspring or healthy controls. An experiential sampling study by the same group examined the interrelationships between mood, self-esteem and response styles over 6 days (Pavlickova *et al.* 2015). Interestingly, increased negative as well as positive mood resulted in greater rumination in offspring and controls, and low self-esteem triggered greater risk-taking in the bipolar offspring group, while negative affect instigated increased active coping in the control group. Additionally, as in studies of depression, CER impairments such as increased ruminative response style are associated with suicidal ideation in young people with emerging bipolar disorders (Stange *et al.* 2015).

### Psychosis

The typical age of onset for psychotic disorders is late adolescence or early twenties, being slightly later in females than males (Hafner *et al.* 1994; Gogtay *et al.* 2011). Several studies report that anxiety or depression in childhood or adolescence may precede the onset of psychosis (Baynes *et al.* 2000; Mulholland & Cooper, 2000; Maggini & Raballo, 2006). Research in groups at high risk of psychosis indicate that transition is common in those who have experienced brief periods of psychosis or psychotic-like symptoms (Yung *et al.* 2006), whilst the presence of depression and/or anxiety is associated with increased

suicidality, self-harm, disorganized behaviour, disorganized speech, and anhedonia (Fusar-Poli *et al.* 2014).

Rumination has been shown to be associated with hallucination-proneness and a range of mild anomalous experiences including feelings of unreality, perceptual alterations, and temporal disintegration (Jones & Fernyhough, 2009; Freeman *et al.* 2013). Experiential sampling in young adults with psychosis demonstrates that antecedent rumination and worry predict persecutory delusions and auditory hallucinations, and that rumination predicted the level of distress associated with these psychotic experiences (Hartley *et al.* 2014). Also, Halari *et al.* (2009) demonstrated that rumination can be associated with negative symptoms such as stereotyped thinking and emotional withdrawal.

As ruminative response style overlaps with perseverative thinking, it could indicate that rumination in psychosis is associated with underlying psychotic or dysphoric (anxiety/depressive) dimensions (Cernis *et al.* 2015). Theoretically, rumination might also help explain why some young people who experience psychotic symptoms become distressed and seek help, whilst others do not (Van Nierop *et al.* 2012; Rapado-Castro *et al.* 2015).

Lastly, in a cross-sectional study of psychotic inpatients with and without suicidality ( $n=2383$ ; mean age at onset 25 years), ruminative thinking, social withdrawal, and lack of activity were all associated with increased suicidality (Ahrens & Linden, 1996).

### Neurobiological underpinnings of rumination

Neurobiological factors play a key role in the activation and maintenance of cognitive-emotional processes. This section briefly examines a selection of the developing research on the associations between rumination in young people and genetics, neuropsychology, physiology, sleep and physical health.

#### Genes

There are reasons to consider the role of genetics in the development of rumination. As noted, rumination may mediate the association between temperamental styles that are heritable (e.g. NE, neuroticism) and depression. Furthermore, children at high risk of depression (e.g. offspring of mothers with a history of MDE) show higher levels of rumination than controls (Gibb *et al.* 2012). A small number of twin studies have examined genetic and environmental influences (shared and non-shared elements) on rumination, brooding and mood disorders. In a Chinese study of adolescent twins, the heritability of depression (30–42%) was modest, but genetic effects accounted for 24% of the variation in adolescent rumination, and genetics mediated

the relationship between rumination and depression (Chen & Li, 2013). In a US study of adolescent twins aged 12–14 years, heritability accounted for 21% of the variation in adolescent brooding, highlighting again that although the heritable influences were modest, they accounted for the majority of the relationship between brooding and depression ( $h^2 = 0.62$ ) (Moore *et al.* 2013). Preliminary data from a UK study of BD in twins and siblings showed that ruminative response style had a low to moderate heritability with 23% of the total variance accounted for by additive genetic effects (Beards *et al.* 2012).

Certain polymorphisms of the BDNF gene (a protein involved in the growth of new and existing neurons and synapses) have also been found to be linked to rumination (Beevers *et al.* 2009). Results suggest that individuals with the Val<sup>66</sup>Met BDNF gene polymorphism are significantly more likely to ruminate than individuals without the gene. This is consistent with recent research that has found an association between the BDNF gene Val<sup>66</sup>Met polymorphism, rumination, and depression (Hilt *et al.* 2007), where analyses suggested that rumination mediated the relationship between the BDNF polymorphism and depressive symptoms.

### Neuropsychology

A number of cognitive processes are activated when engaging in rumination, in particular attention, memory, and self-referential processing (Lyubomirsky *et al.* 1998). Investigators have reported that rumination generates difficulties in controlling the entry of irrelevant information into working memory (WM) and short-term memory (STM) (Joormann, 2004; Goeleven *et al.* 2006; Joormann *et al.* 2010), or the removal of negative self-relevant information from WM and STM storage (Joormann & Gotlib, 2008). Ruminators also show general deficits in the ability to shift attention from unhelpful to helpful strategies (Davis & Nolen-Hoeksema, 2000). Furthermore, a study of over 200 adolescents demonstrated that engaging in ruminative thoughts consumes cognitive resources that would otherwise be allocated towards difficult tests of executive functioning (the resource allocation hypothesis). In contrast, no evidence was found to support the notion that lower levels of executive functioning at baseline predicted levels of rumination or depressive symptoms at follow-up (Connolly *et al.* 2014).

Functional magnetic resonance imaging (fMRI) shows that rumination is positively correlated with an increased activation of the medial prefrontal cortex and amygdala (Ray *et al.* 2005), and adults who ruminate regularly (e.g. dysphoric or depressed patients)

show high connectivity between these structures (Gotlib & Hamilton, 2008). Ruminators also engage in higher rates of self-referential processing compared to their non-ruminative counterparts (Gotlib & Joormann, 2010); a process that can be examined via neuroimaging of the default mode network (DMN) (Ochsner & Gross, 2005; Schmitz & Johnson, 2006). In non-ruminators, activation of DMN brain regions is reduced after performing non-self-referential (i.e. goal-directed) activities during off-task or rest periods, suggesting an ability to lose one's self in the work (Sheline *et al.* 2009). In contrast, high ruminators show high connectivity between DMN regions during these rest periods, reflecting an inability to suppress self-referential thoughts during off-task times. Berman *et al.* (2011b) noted that in young adults (mean age 22 years) these connectivities were only found when participants engaged in brooding and not reflective pondering.

Marchetti *et al.* (2012) propose that an imbalance in the task positive (TP) and task negative (TN) elements of the DMN is the overarching neural mechanism involved in rumination. Studies of rumination in youth (Berman *et al.* 2011a) and of adult depressed cases *v.* controls (Hamilton *et al.* 2011) support the proposal that a TN-TP imbalance is associated with a failure to attenuate TN activity in the transition from rest to task periods and with toxic brooding.

A recent fMRI study compared healthy controls with unmedicated adolescents in remission from MDE (Jacobs *et al.* 2014) and showed that the remitted MDE adolescents exhibited hyper-connectivities within the DMN and between the DMN and salience networks (posterior cingulate cortex, subgenual anterior cingulate, and amygdala) and regions of the cognitive control network, which were related to rumination and sustained attention. However, it was not possible to conclude that the hyper-connectivities represented brain-based markers of traits (e.g. rumination), as it is plausible that the hyper-connectivities represent compensatory mechanisms in individuals with emerging mood disorders. Fewer studies examine CER in individuals aged 15–25 years with emerging bipolar disorders, although in mixed samples of younger and older adults it has been shown that individuals at increased risk of developing bipolar disorders are less successful at down-regulating amygdala activity and demonstrate inefficient use of reappraisal and distraction strategies (e.g. Heissler *et al.* 2014).

### Hypothalamic-pituitary-adrenal (HPA) axis and physiological studies

Several physiological studies have examined the relationship between rumination and heart rate variability.

Key *et al.* (2008) demonstrated that heart rate variability in response to stress is more strongly associated with trait rather than state rumination in female college students.

A review of the relationship between rumination and cortisol identified that 13 of 17 studies were undertaken in children, students or young adults (Zoccola & Dickerson, 2012). Whilst higher levels of state rumination were consistently linked to increased cortisol concentrations, findings were less consistent for the physiological effects of stressors (basal cortisol levels or cortisol awakening response to naturally occurring or laboratory-induced stress). However, those studies that utilized social-evaluative stressor tasks generally showed that rumination predicted greater cortisol reactivity or delayed recovery. These findings are supported by a study using pupillary response to perceived rejection in depressed adolescents (Stone *et al.* 2016).

Gianferante *et al.* (2014) also showed that rumination in response to repeated stress predicted non-habituation of the HPA axis, which the authors suggest may offer a pathway linking rumination to negative health outcomes. Interestingly, this association may be bidirectional or attenuated by baseline physical status (Puterman *et al.* 2011). For example, Puterman *et al.* (2011) demonstrated that although ruminators experienced a more rapid initial increase in cortisol levels, greater HPA axis reactivity, and slower HPA axis recovery from stress, this profile was only significant in sedentary participants. In active participants, cortisol trajectories were equivalent in high and low ruminators. These findings are noteworthy given the evidence of increased sedentary lifestyles and lower physical activity in youth with mental disorders (even when medication-free) compared to their peers (Gehue *et al.* 2015; Vallarino *et al.* 2015).

### *Sleep and circadian rhythms*

Rumination may indirectly influence circadian rhythms through its association with increased cortisol production (Rea *et al.* 2010) and alterations in cortisol secretion patterns (for a review see Chan & Debono, 2010). Furthermore, studies of sleep in adolescents show rumination to be directly associated with poorer general sleep quality, longer time to fall asleep and more awakenings after sleep onset (Thomsen *et al.* 2003). Dysregulated sleep is associated with increased rumination compared to good sleepers with evidence that brooding is associated with fatigue, poor concentration and low mood (Carney *et al.* 2006). Zoccola *et al.* (2009) found that ruminating about a past stressor before bedtime predicted longer sleep onset latencies. This suggestion that post-stressor ruminative thought


may predict delayed sleep onset is of particular interest as delayed sleep phase syndrome is a recognized marker of circadian disturbance that occurs more often in young compared to older adults (Robillard *et al.* 2014; Alloy *et al.* 2015; Steinar *et al.* 2015). Batterham *et al.* (2012) found that rumination and neuroticism mediated the relationship between self-reported sleep disturbance and new onset depressions in younger adults, but not the onset of generalized anxiety of panic disorders.

The above studies mainly relied on self-ratings rather than objective recordings of sleep. An exception is the study of university students (mean age about 20 years) by Pillai *et al.* (2014), which showed that nightly variations in pre-sleep rumination were predictive of significantly longer sleep onset latency (SOL) as recorded by actigraphy and by a self-report diary. It was estimated that, after controlling for baseline sleep disturbance and depressive symptoms, a one standard deviation increase on the pre-sleep rumination scale was associated with an approximately 7-min increase in actigraphy-based SOL.

### *Physical health and immune system*

Rumination has been found to be negatively associated with self-reported physical health, including higher levels of somatic complaints and lower general health (Lok & Bishop, 1999; Rector & Roger, 1996). In a longitudinal study of younger compared to older adults, Thomsen *et al.* (2004a) demonstrated that higher levels of rumination predicted poorer self-reported physical health and more somatic complaints in those aged 20–35 years only. High levels of rumination have been linked with immune functioning, but to date these associations have only been reported for some of the immune markers measured (numbers of leucocytes, lymphocytes, and polyclonal activation), and the findings were significant in older but not younger adults (Thomsen *et al.* 2004b).

Brosschot (2010) suggests that sustained cognitive representations of events (namely rumination and worry) can cause prolonged physiological activity, which may lead to intermediate pathogenic states (such as increased allostatic load) and finally to somatic disease. Glynn *et al.* (2002) found that ruminating about an emotional task resulted in increased blood pressure and delayed blood pressure recovery, which parallel findings reported in students with high levels of rumination and in adults (Bermúdez & Perez-Garcia, 1996; Neumann *et al.* 2001). Finally, although the co-morbidity between mood disorders and cardiovascular disease reported in young adults may be linked to medications or lifestyle factors (Goldstein *et al.* 2015), it is hypothesized that trait rumination



**Fig. 2.** Model of rumination as a trans-diagnostic process impacting psychopathology and physical health, underpinned by genes and neural networks, age, gender, and temperament, and early environment. AUD, Alcohol use disorder; HPA, hypothalamic-pituitary-adrenal; SUD, substance use disorder.

may delay physiological recovery from acute stress and could act as a mechanism (Larsen & Christenfeld, 2009) (Fig. 2).

### Conclusions

This paper highlights several key aspects of rumination. First, we note that, of the two core components of rumination, it is the maladaptive (brooding) rather than the adaptive element (reflection or positive basking) that is consistently associated with the development of psychopathology. Also, the normative pattern of development of rumination parallels age and gender profiles associated with the typical evolution of internalizing disorders. Importantly, we highlight the association between the development of a ruminative response style and traumatic and abusive experiences in childhood, which are risk factors linked to a range of psychological problems in adolescence including mood and psychotic disorders. The evidence suggests that rumination is an underlying mechanism that contributes to a significant proportion of the explained variance between early adversity and later mental health problems. We do not claim that rumination is the only explanatory model, as other cognitive structures (e.g. beliefs and schemata) and processes have been implicated in the links between early trauma, CER and the development of psychopathology

and acts of self-harm. Several studies demonstrate that these models and constructs may overlap and that there are links between early maladaptive schema, rumination and future symptoms of anxiety and depression in adolescents (Orue *et al.* 2014; Black & Pössel, 2015) or between early childhood trauma, over-general memory and high levels of ruminative thinking (Watkins & Teasdale, 2001; Williams *et al.* 2007). However, more research is needed to clarify the interactions between these different CER elements.

As reported in reviews of older adult populations, we confirm the importance of rumination in the evolution and maintenance of depression and highlight that in adolescence (but not always older adults) rumination may fully or partially mediate the relationship between childhood temperament and/or stressful events and the later onset of anxiety disorders. This sequence offers support for the notion of rumination being important in the longitudinal trajectory of the development of mental disorders and the notion that rumination plays a role in the transition between clinical stages. Our review differs from many previous reviews of rumination as a trans-diagnostic process operating across age groups and populations (Harvey *et al.* 2004; Ehrling & Watkins, 2008; Watkins, 2009; Watkins & Nolen-Hoeksema, 2014), as we also examine the possible role of rumination in evolution of a problem from its sub-threshold phenomenology

through to levels of harmful alcohol and substance use, bipolar and psychotic disorders that lead to help-seeking in an individual. Furthermore, we explore its putative influence on the trajectories of illness observed in this younger population and how similar processes may operate across physical and mental disorders. There is emerging evidence that rumination can be directly linked to the onset or maintenance of these problems in young adults, as well as existing evidence of indirect associations, with rumination exacerbating levels of anxiety or depressive symptoms, which in turn increase the distress that accompanies the symptoms or impede the ability to cope with those symptoms. It was also notable that rumination is frequently associated with suicidal ideation in depression, bipolar and psychotic disorders. We suggest that this is an important area for research, as the role of rumination in amplifying mood states and reducing flexibility in thinking styles and CER is under-explored in the age groups at the highest risk for onset of severe mental disorders or deliberate self-harm.

Evidence of the heritability of and neuropsychological pathways implicated in rumination is important as, for example, changes in the neural networks associated with CER may precede behavioural manifestations in unaffected high risk populations (Heissler *et al.* 2014). Critically, this review suggests ruminative response style is a putative shared mechanism for the development of sleep and physiological dysregulation and health problems in young adults with evolving mental disorders. This is relevant as attempts to explain physical and mental co-morbidities on the basis of factors such as the adverse physical effects of treatment or reduced daytime activity (post-onset of mental disorder) have proven to be overly simplistic and fail to take into account findings of the high levels of physical and mental co-morbidity in untreated, early stage cases. It appears that a ruminative response style may be associated with circadian and HPA disruptions, and its relationship to physical disease is not driven by greater reactivity in systems, but instead through extending activation and increasing allostatic load, which may increase the risks for cardiovascular or other damage (Larsen & Christenfeld, 2009). Indeed, Ottaviani *et al.* (2009) suggest that rumination represents an 'autonomic phenotype' because the associated autonomic dysregulation plays a role in the relationships between temperament, anxiety, depression and cardiovascular health.

In conclusion, this selective review of the association between rumination and the early clinical stages of mental disorders suggests that it is an important underlying trans-diagnostic process that operates in adolescents and young adults. Further, rumination shows a predictable developmental trajectory that is

both detectable and modifiable. From the perspective of primary or early secondary prevention, modifying the maladaptive 'toxic brooding' component of rumination and substituting a more flexible response style that engenders a greater sense of self-control has the potential to reduce the onset or maintenance of problems such as depression and anxiety (Cook & Watkins, 2016), and may reduce the distress or severity of other disorders, such as psychosis and bipolar disorders (Vallarino *et al.* 2015; Scott, 2016). This review indicates that examination of rumination and other CER is an important, underexplored issue in trans-diagnostic and dimensional approaches to mental disorders, which is especially important given the apparent links to physical and well as mental health, and the prospects for modification of this risk factor.

### Acknowledgements

A.G. has received grant funding from the National Health and Medical Research Council in Australia. I.B.H. is a Commissioner in Australia's National Mental Health Commission; a Member of the Medical Advisory Panel for Medibank; a Board Member of Psychosis Australia Trust. He has received honoraria for presentations of his own work at educational seminars supported by a number of non-government organizations and by the pharmaceutical industry (including Servier, Pfizer, AstraZeneca, and Eli Lilly) and funding from Servier for a study of major depression and sleep disturbance in primary-care settings. Other relevant funding for Professor Ian Hickie is in relation to this study includes 'Testing and delivering early interventions for young people with depression' (APP ID: 1046899). S.N. has received grant funding from the National Health and Medical Research Council including for research on sleep and actigraphy. J.S. is a visiting professor at the Brain & Mind Centre at The University of Sydney. J.S. has received UK grant funding from the Medical Research Council (including for projects on actigraphy and bipolar disorders) and from the Research for Patient Benefit programme (PB-PG-0609-16166: Early identification and intervention in young people at risk of mood disorders).

### Declaration of Interest

None.

### References

- Abela JR, Aydin CM, Auerbach RP (2007). Responses to depression in children: reconceptualizing the relation among response styles. *Journal of Abnormal Child Psychology* 35, 913–927.

- Adrian M, McCarty C, King K, McCauley E, Vander Stoep A** (2014). The internalizing pathway to adolescent substance use disorders: mediation by ruminative reflection and ruminative brooding. *Journal of Adolescence* **37**, 983–991.
- Ahrens B, Linden M** (1996). Is there a suicidality syndrome independent of specific major psychiatric disorder? Results of a split half multiple regression analysis. *Acta Psychiatrica Scandinavica* **94**, 79–86.
- Aldao A, Nolen-Hoeksema S** (2012). When are adaptive strategies most predictive of psychopathology? *Journal of Abnormal Psychology*, **121**, 276.
- Allen KJ, Gabbay FH** (2013). The amphetamine response moderates the relationship between negative emotionality and alcohol use. *Alcoholism: Clinical and Experimental Research* **37**, 348–360.
- Alloy LB, Nusslock R, Boland EM** (2015). The development and course of bipolar spectrum disorders: an integrated reward and circadian rhythm dysregulation model. *Annual Review of Clinical Psychology* **11**, 213.
- Baskin-Sommers AR, Foti D** (2015). Abnormal reward functioning across substance use disorders and major depressive disorder: considering reward as a transdiagnostic mechanism. *International Journal of Psychophysiology* **98**, 227–239.
- Batterham PJ, Glozier N, Christensen H** (2012). Sleep disturbance, personality and the onset of depression and anxiety: prospective cohort study. *Australian and New Zealand Journal of Psychiatry* **46**, 1089–1098.
- Baynes D, Mulholland C, Cooper SJ, Montgomery RC, MacFlynn G, Lynch G, Kelly C, King DJ** (2000). Depressive symptoms in stable chronic schizophrenia: prevalence and relationship to psychopathology and treatment. *Schizophrenia Research* **45**, 47–56.
- Beards S, Georgiades A, Sharma M, Kane F, Kalidindi S, Schulze K, Walshe M, Scott J, Murray R, Rijsdijk F, Kraviriti E** (2012). *Psychological endophenotypes for bipolar disorders*. Abstracts of the 3rd Schizophrenia International Research Society Conference, Florence, Italy.
- Bechdolf A, Ratheesh A, Cotton SM, Nelson B, Chanen AM, Betts J, Bingmann T, Yung AR, Berk M, McGorry PD** (2014). The predictive validity of bipolar at-risk (prodromal) criteria in help-seeking adolescents and young adults: a prospective study. *Bipolar Disorders* **16**, 493–504.
- Beevers CG, Wells TT, McGeeary JE** (2009). The BDNF Val66Met polymorphism is associated with rumination in healthy adults. *Emotion* **9**, 579.
- Belsky J, Hsieh KH, Crnic K** (1996). Infant positive and negative emotionality: one dimension or two? *Developmental Psychology*, **32**, 289.
- Berman MG, Nee DE, Casement M, Kim HS, Deldin P, Kross E, Gonzalez R, Demiralp E, Gotlib IH, Hamilton P, Joermann J, Waugh C, Jonides J** (2011a). Neural and behavioral effects of interference resolution in depression and rumination. *Cognitive, Affective, & Behavioral Neuroscience* **11**, 85–96.
- Berman MG, Peltier S, Nee DE, Kross E, Deldin PJ, Jonides J** (2011b). Depression, rumination and the default network. *Social, Cognitive, and Affective Neuroscience* **6**, 548–555.
- Bermúdez J, Perez-García AM** (1996). Cardiovascular reactivity, affective responses and performance related to the risk dimensions of coronary-prone behaviour. *Personality and Individual Differences* **21**, 919–927.
- Bittner A, Goodwin RD, Wittchen HU, Beesdo K, Höfler M, Lieb R** (2004). What characteristics of primary anxiety disorders predict subsequent major depressive disorder? *Journal of Clinical Psychiatry* **65**, 618–26.
- Black SW, Pössel P** (2015). Integrating Beck's cognitive model and the response style theory in an adolescent sample. *Journal of Youth and Adolescence* **44**, 195–210.
- Blatt SJ, Homann E** (1992). Parent-child interaction in the aetiology of dependent and self-critical depression. *Clinical Psychology Review* **12**, 47–91.
- Broderick PC** (1998). Early adolescent gender differences in the use of ruminative and distracting coping strategies. *Journal of Early Adolescence* **18**, 173–191.
- Brosschot JF** (2010). Markers of chronic stress: prolonged physiological activation and (un)conscious perseverative cognition. *Neuroscience & Biobehavioral Reviews* **35**, 46–50.
- Brosschot JF, Gerin W, Thayer JF** (2006). The perseverative cognition hypothesis: a review of worry, prolonged stress-related physiological activation, and health. *Journal of Psychosomatic Research* **60**, 113–124.
- Browne A, Finkelhor D** (1986). Impact of child sexual abuse: a review of the research. *Psychological Bulletin* **99**, 66.
- Carney CE, Edinger JD, Meyer B, Lindman L, Istre T** (2006). Symptom-focused rumination and sleep disturbance. *Behavioral Sleep Medicine* **4**, 228–241.
- Caselli G, Bortolai C, Leoni M, Rovetto F, Spada MM** (2008). Rumination in problem drinkers. *Addiction Research & Theory* **16**, 564–571.
- Caselli G, Ferretti C, Leoni M, Rebecchi D, Rovetto F, Spada MM** (2010). Rumination as a predictor of drinking behaviour in alcohol abusers: a prospective study. *Addiction* **105**, 1041–1048.
- Cernis E, Dunn G, Startup H, Kingdon D, Wingham G, Evans N, Lister R, Pugh K, Cordwell J, Mander H, Freeman D** (2015). The perseverative thinking questionnaire in patients with persecutory delusions. *Behavioural and Cognitive Psychotherapy* **44**, 472–481.
- Chan S, Debono M** (2010). Review: replication of cortisol circadian rhythm: new advances in hydrocortisone replacement therapy. *Therapeutic Advances in Endocrinology and Metabolism* **1**, 129–138.
- Chan S, Miranda R, Surrence K** (2009). Subtypes of rumination in the relationship between negative life events and suicidal ideation. *Archives of Suicide Research* **13**, 123–135.
- Chen J, Li X** (2013). Genetic and environmental influences on adolescent rumination and its association with depressive symptoms. *Journal of Abnormal Child Psychology* **41**, 1289–1298.
- Cole DA, Nolen-Hoeksema S, Girgus J, Paul G** (2006). Stress exposure and stress generation in child and adolescent depression: a latent trait-state-error approach to longitudinal analyses. *Journal of Abnormal Psychology* **115**, 40.
- Compas BE** (1987). Coping with stress during childhood and adolescence. *Psychological Bulletin* **101**, 393.
- Connolly SL, Wagner CA, Shapero BG, Pendergast LL, Abramson LY, Alloy LB** (2014). Rumination prospectively

- predicts executive functioning impairments in adolescents. *Journal of Behavior Therapy and Experimental Psychiatry* **45**, 46–56.
- Conway M, Csank PAR, Holm SL, Blake CK** (2000). On assessing individual differences in rumination on sadness. *Journal of Personality Assessment* **75**, 404–425.
- Conway M, Mendelson M, Giannopoulos C, Csank PA, Holm SL** (2004). Childhood and adult sexual abuse, rumination on sadness, and dysphoria. *Child Abuse & Neglect* **28**, 393–410.
- Cook L, Watkins ER** (2016). Guided, internet-based, rumination-focused cognitive behavioural therapy (i-RFCBT) versus a no-intervention control to prevent depression in high-ruminating young adults, along with an adjunct assessment of the feasibility of unguided i-RFCBT, in the REducing Stress and Preventing Depression trial (RESPOND): study protocol for a phase III randomised controlled trial. *Trials* **17**, 1.
- Cox SJ, Mezulis AH, Hyde JS** (2010). The influence of child gender role and maternal feedback to child stress on the emergence of the gender difference in depressive rumination in adolescence. *Developmental Psychology* **46**, 842.
- Davis RN, Nolen-Hoeksema S** (2000). Cognitive inflexibility among ruminators and non-ruminators. *Cognitive Therapy and Research* **24**, 699–711.
- de Jong-Meyer R, Beck B, Riede K** (2009). Relationships between rumination, worry, intolerance of uncertainty and metacognitive beliefs. *Personality and Individual Differences* **46**, 547–551.
- Dennis M, Babor TF, Roebuck C, and Donaldson J** (2002). Changing the focus: the case for recognizing and treating cannabis use disorders. *Addiction* **97**, 4–15.
- Douglas J, Scott J** (2014). A systematic review of gender-specific rates of unipolar and bipolar disorders in community studies of pre-pubertal children. *Bipolar Disorders* **16**, 5–15.
- Duffy A, Alda M, Hajek T, Grof P** (2009). Early course of bipolar disorder in high-risk offspring: prospective study. *British Journal of Psychiatry* **195**, 457–458.
- Dugas MJ, Schwartz A, Francis K** (2004). Intolerance of uncertainty, worry, and depression. *Cognitive Therapy and Research* **28**, 835–842.
- Eckblad M, Chapman LJ** (1986). Development and validation of a scale for hypomanic personality. *Journal of Abnormal Psychology* **95**, 214.
- Ehring T, Watkins ER** (2008). Repetitive negative thinking as a transdiagnostic process. *International Journal of Cognitive Therapy* **1**, 192–205.
- Einstein DA** (2014). Extension of the transdiagnostic model to focus on intolerance of uncertainty: a review of the literature and implications for treatment. *Clinical Psychology: Science and Practice* **21**, 280–300.
- Emmons RA, King LA** (1988). Conflict among personal strivings: immediate and long-term implications for psychological and physical well-being. *Journal of Personality and Social Psychology* **54**, 1040.
- Eysenck HJ, Eysenck MW** (1985). *Personality and Individual Differences*. Plenum: New York.
- Feldman GC, Joormann J, Johnson SL** (2008). Responses to positive affect: a self-report measure of rumination and dampening. *Cognitive Therapy and Research* **32**, 507–525.
- Freeman D, Startup H, Dunn G, Černis E, Wingham G, Pugh K, Cordwell J, Kingdon D** (2013). The interaction of affective with psychotic processes: a test of the effects of worrying on working memory, jumping to conclusions, and anomalies of experience in patients with persecutory delusions. *Journal of Psychiatric Research* **47**, 1837–1842.
- Fusar-Poli P, Nelson B, Valmaggia L, Yung AR, McGuire PK** (2014). Comorbid depressive and anxiety disorders in 509 individuals with an at-risk mental state: impact on psychopathology and transition to psychosis. *Schizophrenia Bulletin* **40**, 120–131.
- Garnefski N, Kraaij V, Spinhoven P** (2001). Negative life events, cognitive emotion regulation and emotional problems. *Personality and Individual Differences* **30**, 1311–1327.
- Garnefski N, Legerstee J, Kraaij V, Van Den Kommer T, Teerds JAN** (2002). Cognitive coping strategies and symptoms of depression and anxiety: a comparison between adolescents and adults. *Journal of Adolescence* **25**, 603–611.
- Garnefski N, Kraaij V** (2007). The cognitive emotion regulation questionnaire. *European Journal of Psychological Assessment* **23**, 141–149.
- Garnefski N, Kraaij V, van Etten M** (2005). Specificity of relations between adolescents' cognitive emotion regulation strategies and internalizing and externalizing psychopathology. *Journal of Adolescence* **28**, 619–631.
- Gehue LJ, Scott E, Hermens DF, Scott J, Hickie I** (2015). Youth Early-intervention Study (YES) – group interventions targeting social participation and physical well-being as an adjunct to treatment as usual: study protocol for a randomized controlled trial. *Trials* **16**, 333.
- Geoffroy PA, Leboyer M, Scott J** (2013). Predicting bipolar disorder: what can we learn from prospective cohort studies? *Encephale* **41**, 10–16.
- Ghaznavi S, Deckersbach T** (2012). Rumination in bipolar disorder: evidence for an unquiet mind. *Biology of Mood and Anxiety Disorders* **2**, 2.
- Gianferante D, Thoma MV, Hanlin L, Chen X, Breines JG, Zoccola PM, Rohleder N** (2014). Post-stress rumination predicts HPA axis responses to repeated acute stress. *Psychoneuroendocrinology* **49**, 244–252.
- Gibb BE, Grassia M, Stone LB, Uhrlass DJ, McGeary JE** (2012). Brooding rumination and risk for depressive disorders in children of depressed mothers. *Journal of Abnormal Child Psychology* **40**, 317–326.
- Glynn LM, Christenfeld N, Gerin W** (2002). The role of rumination in recovery from reactivity: cardiovascular consequences of emotional states. *Psychosomatic Medicine* **64**, 714–726.
- Goeleven E, De Raedt R, Baert S, Koster EH** (2006). Deficient inhibition of emotional information in depression. *Journal of Affective Disorders* **93**, 149–157.
- Gogtay N, Vyas NS, Testa R, Wood SJ, Pantelis C** (2011). Age of onset of schizophrenia: perspectives from structural neuroimaging studies. *Schizophrenia Bulletin* **37**, 504–513.
- Gohm CL** (2003). Mood regulation and emotional intelligence: individual differences. *Journal of Personality and Social Psychology* **84**, 594.

- Goldstein BI, Carnethon MR, Matthews KA, McIntyre RS, Miller GE, Raghuvveer G, Stoney CM, Wasiaik H, McCrindle BW** (2015). Major depressive disorder and bipolar disorder predispose youth to accelerated atherosclerosis and early cardiovascular disease: a scientific statement from the American Heart Association. *Circulation* **132**, 965–986.
- Gotlib IH, Hamilton JP** (2008). Neuroimaging and depression current status and unresolved issues. *Current Directions in Psychological Science* **17**, 159–163.
- Gotlib IH, Joormann J** (2010). Cognition and depression: current status and future directions. *Annual Review of Clinical Psychology* **6**, 285.
- Green MJ, Lino BJ, Hwang EJ, Sparks A, James C, Mitchell PB** (2011). Cognitive regulation of emotion in bipolar I disorder and unaffected biological relatives. *Acta Psychiatrica Scandinavica* **124**, 307–316.
- Gross JJ, Munoz RF** (1995). Emotion regulation and mental health. *Clinical Psychology: Science and Practice* **2**, 151–164.
- Gyurak A, Gross JJ, Etkin A** (2011). Explicit and implicit emotion regulation: a dual-process framework. *Cognition and Emotion* **25**, 400–412.
- Hafner H, Maurer K, Löffler W, Fatkenheuer B** (1994). The epidemiology of early schizophrenia influence of age and gender on onset and early course. *British Journal of Psychiatry* **164**, 29–38.
- Halari R, Premkumar P, Farquharson L, Fannon D, Kuipers E, Kumari V** (2009). Rumination and negative symptoms in schizophrenia. *Journal of Nervous and Mental Disease* **197**, 703–706.
- Hamilton JL, Hamlat EJ, Stange JP, Abramson LY, Alloy LB** (2014a). Pubertal timing and vulnerabilities to depression in early adolescence: differential pathways to depressive symptoms by sex. *Journal of Adolescence* **37**, 165–174.
- Hamilton JL, Stange JP, Kleiman EM, Hamlat EJ, Abramson LY, Alloy LB** (2014b). Cognitive vulnerabilities amplify the effect of early pubertal timing on interpersonal stress generation during adolescence. *Journal of Youth and Adolescence* **43**, 824–833.
- Hamilton JP, Furman DJ, Chang C, Thomason ME, Dennis E, Gotlib IH** (2011). Default-mode and task-positive network activity in major depressive disorder: implications for adaptive and maladaptive rumination. *Biological Psychiatry* **70**, 327–333.
- Hampel P, Petermann F** (2005). Age and gender effects on coping in children and adolescents. *Journal of Youth and Adolescence* **34**, 73–83.
- Hankin BL, Oppenheimer C, Jenness J, Barrocas A, Shapero BG, Goldband J** (2009). Developmental origins of cognitive vulnerabilities to depression: review of processes contributing to stability and change across time. *Journal of Clinical Psychology* **65**, 1327–1338.
- Hartley S, Haddock G, Vasconcelos e Sa D, Emsley R, Barrowclough C** (2014). An experience sampling study of worry and rumination in psychosis. *Psychological Medicine* **44**, 1605–1614.
- Harvey AG, Watkins ER, Mansell W, Shafran R** (2004). *Cognitive Behavioural Processes Across Psychological Disorders: A Transdiagnostic Approach to Research and Treatment*. Oxford University Press: Oxford.
- Heissler J, Kanske P, Schonfelder S, Wessa M** (2014). Inefficiency of emotion regulation as vulnerability marker for bipolar disorder: evidence from healthy individuals with hypomanic personality. *Journal of Affective Disorders* **152**, 83–90.
- Helzer JE, Burnam A, McEvoy LT** (1991). Alcohol abuse and dependence. In *Psychiatric Disorders in America: The Epidemiologic Catchment Area Study*, (ed. L. Robins and D. Regier), pp. 81–115. MacMillan: New York.
- Hickie IB, Naismith SL, Robillard R, Scott EM, Hermens DF** (2013). Manipulating the sleep-wake cycle and circadian rhythms to improve clinical management of major depression. *BMC Medicine* **11**, 79.
- Hilt LM, Armstrong JM, Essex MJ** (2012). Early family context and development of adolescent ruminative style: moderation by temperament. *Cognition & Emotion* **26**, 916–926.
- Hilt LM, Sander LC, Nolen-Hoeksema S, Simen AA** (2007). The BDNF Val66Met polymorphism predicts rumination and depression differently in young adolescent girls and their mothers. *Neuroscience Letters* **429**, 12–16.
- Horn PJ, Wuyek LA** (2010). Anxiety disorders as a risk factor for subsequent depression. *International Journal of Psychiatry in Clinical Practice* **14**, 244–247.
- Hou WK, Ng SM** (2014). Emotion-focused positive rumination and relationship satisfaction as the underlying mechanisms between resilience and psychiatric symptoms. *Personality and Individual Differences* **71**, 159–164.
- Hudson JL, Rapee RM** (2004). From anxious temperament to disorder: an etiological model of generalized anxiety disorder. In *Generalized Anxiety Disorder: Advances in Research and Practice* (ed. R. G. Heimberg, C. L. Turk and D. S. Mennin), pp. 51–76. Guilford Press: New York.
- Hyde JS, Mezulis AH, Abramson LY** (2008). The ABCs of depression: integrating affective, biological, and cognitive models to explain the emergence of the gender difference in depression. *Psychological Review* **115**, 291.
- Insel T, Cuthbert B, Garvey M, Heinszen R, Pine DS, Quinn K, Sanislow C, Wang P** (2010). Research domain criteria (RDoC): toward a new classification framework for research on mental disorders. *American Journal of Psychiatry* **167**, 748–751.
- Jacobs RH, Jenkins LM, Gabriel LB, Barba A, Ryan KA, Weisenbach SL, Verges A, Baker AM, Peters AT, Crane NA, Gotlib IH, Zubieta JK, Phan KL, Langenecker SA, Welsh RC** (2014). Increased coupling of intrinsic networks in remitted depressed youth predicts rumination and cognitive control. *PLoS ONE* **9**, e104366.
- Johnson SL, McKenzie G, McMurrich S** (2008). Ruminative responses to negative and positive affect among students diagnosed with bipolar disorder and major depressive disorder. *Cognitive Therapy and Research* **32**, 702–713.
- Jones NP, Papadakis AA, Hogan CM, Strauman TJ** (2009). Over and over again: rumination, reflection, and promotion goal failure and their interactive effects on depressive symptoms. *Behaviour Research and Therapy* **47**, 254–259.
- Jones PB** (2013). Adult mental health disorders and their age at onset. *British Journal of Psychiatry* **202**(s54), s5–s10.
- Jones SH, Tai S, Evershed K, Knowles R, Bentall R** (2006). Early detection of bipolar disorder: a pilot familial high-risk study of parents with bipolar disorder and their adolescent children. *Bipolar Disorders* **8**, 362–372.

- Jones SR, Fernyhough C (2009). Rumination, reflection, intrusive thoughts, and hallucination-proneness: towards a new model. *Behaviour Research and Therapy* **47**, 54–59.
- Joormann J (2004). Attentional bias in dysphoria: the role of inhibitory processes. *Cognition and Emotion* **18**, 125–147.
- Joormann J, Gotlib IH (2008). Updating the contents of working memory in depression: interference from irrelevant negative material. *Journal of Abnormal Psychology* **117**, 182.
- Joormann J, Nee DE, Berman MG, Jonides J, Gotlib IH (2010). Interference resolution in major depression. *Cognitive, Affective & Behavioral Neuroscience* **10**, 21–33.
- Jose PE, Brown I (2008). When does the gender difference in rumination begin? Gender and age differences in the use of rumination by adolescents. *Journal of Youth and Adolescence* **37**, 180–192.
- Jose PE, Ratcliffe V (2004). Stressor frequency and perceived intensity as predictors of internalizing symptoms: gender and age differences in adolescence. *New Zealand Journal of Psychology* **33**, 145–154.
- Jose PE, Weir KF (2013). How is anxiety involved in the longitudinal relationship between brooding rumination and depressive symptoms in adolescents? *Journal of Youth and Adolescence* **42**, 1210–1222.
- Jose PE, Wilkins H, Spendelov JS (2012). Does social anxiety predict rumination and co-rumination among adolescents?. *Journal of Clinical Child & Adolescent Psychology* **41**, 86–91.
- Kaminer Y, Bukstein OG (2008). *Adolescent Substance Abuse: Psychiatric Comorbidity and High-Risk Behaviors*. Taylor & Francis: New York, NY.
- Kessler RC, Chiu WT, Demler O, Walters EE (2005). Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Archives of General Psychiatry* **62**, 617–627.
- Key BL, Campbell TS, Bacon SL, Gerin W (2008). The influence of trait and state rumination on cardiovascular recovery from a negative emotional stressor. *Journal of Behavioral Medicine* **31**, 237–248.
- Knowles R, Tai S, Christensen I, Bentall R (2005). Coping with depression and vulnerability to mania: a factor analytic study of the Nolen-Hoeksema (1991). Response Styles Questionnaire. *British Journal of Clinical Psychology* **44**, 99–112.
- Kraaij V, Garnefski N, de Wilde EJ, Dijkstra A, Gebhardt W, Maes S, ter Doest L (2003). Negative life events and depressive symptoms in late adolescence: bonding and cognitive coping as vulnerability factors?. *Journal of Youth and Adolescence* **32**, 185–193.
- Kuyken W, Watkins ER, Holden E, Cook W (2006). Rumination in adolescents at risk for depression. *Journal of Affective Disorders* **96**, 39–47.
- Lamers F, van Oppen P, Comijs HC, Smit JH, Spinhoven P, Van Balkom AJ, Penninx BW (2011). Comorbidity patterns of anxiety and depressive disorders in a large cohort study: the Netherlands Study of Depression and Anxiety (NESDA). *Journal of Clinical Psychiatry* **72**, 341–348.
- Larsen BA, Christenfeld NJ (2009). Cardiovascular disease and psychiatric comorbidity: the potential role of perseverative cognition. *Cardiovascular Psychiatry and Neurology*, vol. **2009**, doi:10.1155/2009/791017.
- Last CG, Hansen C, Franco N (1997). Anxious children in adulthood: a prospective study of adjustment. *Journal of the American Academy of Child & Adolescent Psychiatry* **36**, 645–652.
- Lewinsohn PM, Rohde P, Seeley JR (1998). Major depressive disorder in older adolescents: prevalence, risk factors, and clinical implications. *Clinical Psychology Review* **18**, 765–794.
- Liao KYH, Wei M (2011). Intolerance of uncertainty, depression, and anxiety: the moderating and mediating roles of rumination. *Journal of Clinical Psychology* **67**, 1220–1239.
- Lok CF, Bishop GD (1999). Emotion control, stress, and health. *Psychology and Health* **14**, 813–827.
- Lyubomirsky S, Caldwell ND, Nolen-Hoeksema S (1998). Effects of ruminative and distracting responses to depressed mood on retrieval of autobiographical memories. *Journal of Personality and Social Psychology* **75**, 166.
- MacLeod C, Rutherford E, Campbell L, Ebsworthy G, Holker L (2002). Selective attention and emotional vulnerability: assessing the causal basis of their association through the experimental manipulation of attentional bias. *Journal of Abnormal Psychology* **111**, 107.
- Maggini C, Raballo A (2006). Exploring depression in schizophrenia. *European Psychiatry* **21**, 227–232.
- Manfredi C, Caselli G, Rovetto F, Rebecchi D, Ruggiero GM, Sassaroli S, Spada MM (2011). Temperament and parental styles as predictors of ruminative brooding and worry. *Personality and Individual Differences* **50**, 186–191.
- Marchetti I, Koster EH, Sonuga-Barke EJ, De Raedt R (2012). The default mode network and recurrent depression: a neurobiological model of cognitive risk factors. *Neuropsychology Review* **22**, 229–251.
- Mansell W, Carey TA (2013). Perceptual control theory as an integrative framework for clinical interventions. In *Cognitive Behavioural Approaches to the Understanding and Treatment of Dissociation* (ed. F. Kennedy, H. Kennerley and D. Pearson), pp. 221–235. Routledge: London.
- Martin LL, Tesser A, McIntosh WD (1993). Wanting but not having: The effects of unattained goals on thoughts and feelings. In *Handbook of Mental Control* (ed. D. M. Wegner and J. W. Pennybaker), pp. 552–572. Prentice-Hall: Englewood Cliffs, US.
- McGorry P, Keshavan M, Goldstone S, Amminger P, Allott K, Berk M, Lavoie S, Pantelis C, Yung A, Wood S, Hickie I (2014). Biomarkers and clinical staging in psychiatry. *World Psychiatry* **13**, 211–223.
- McGorry PD, Hickie IB, Yung AR, Pantelis C, Jackson HJ (2006). Clinical staging of psychiatric disorders: a heuristic framework for choosing earlier, safer and more effective interventions. *Australian and New Zealand Journal of Psychiatry* **40**, 616–622.
- McIntosh WD, Harlow TE, Martin LL (1995). Linkers and non-linkers: goal beliefs as a moderator of the effects of everyday hassles on rumination, depression, and physical complaints. *Journal of Applied Social Psychology* **25**, 1231–1244.
- McIntosh WD, Martin LL (1992). The cybernetics of happiness: the relation of goal attainment, rumination, and affect. *Review of Personality and Social Psychology* **14**, 222–246.

- McLaughlin KA, Hatzenbuehler ML** (2009). Stressful life events, anxiety sensitivity, and internalizing symptoms in adolescents. *Journal of Abnormal Psychology* **118**, 659.
- McLaughlin KA, Hatzenbuehler ML, Hilt LM** (2009). Emotion dysregulation as a mechanism linking peer victimization to internalizing symptoms in adolescents. *Journal of Consulting and Clinical Psychology* **77**, 894.
- McLaughlin KA, Nolen-Hoeksema S** (2012). Interpersonal stress generation as a mechanism linking rumination to internalizing symptoms in early adolescents. *Journal of Clinical Child & Adolescent Psychology* **41**, 584–597.
- Merikangas KR, He JP, Burstein M, Swanson SA, Avenevoli S, Cui L, Benjet C, Georgiades K, Swendsen J** (2010). Lifetime prevalence of mental disorders in US adolescents: results from the National Comorbidity Survey Replication–Adolescent Supplement (NCS-A). *Journal of the American Academy of Child & Adolescent Psychiatry* **49**, 980–989.
- Merikangas KR, McClair VL** (2012). Epidemiology of substance use disorders. *Human Genetics* **131**, 779–789.
- Mezulis A, Salk RH, Hyde JS, Priess-Groben HA, Simonson JL** (2014). Affective, biological, and cognitive predictors of depressive symptom trajectories in adolescence. *Journal of Abnormal Child Psychology* **42**, 539–550.
- Mezulis AH, Priess HA, Hyde JS** (2011). Rumination mediates the relationship between infant temperament and adolescent depressive symptoms. *Depression Research and Treatment*, vol. **2011**, doi:10.1155/2011/487873.
- Michalak EE, Hole R, Livingston JD, Murray G, Parikh SV, Lapsley S, McBride S** (2012). Improving care and wellness in bipolar disorder: origins, evolution and future directions of a collaborative knowledge exchange network. *International Journal of Mental Health Systems* **6**, 1–13.
- Michl LC, McLaughlin KA, Shepherd K, Nolen-Hoeksema S** (2013). Rumination as a mechanism linking stressful life events to symptoms of depression and anxiety: longitudinal evidence in early adolescents and adults. *Journal of Abnormal Psychology* **122**, 339.
- Moore MN, Salk RH, Van Hulle CA, Abramson LY, Hyde JS, Lemery-Chalfant K, Goldsmith HH** (2013). Genetic and environmental influences on rumination, distraction, and depressed mood in adolescence. *Clinical Psychological Science* **1**, 316–322.
- Mulholland C, Cooper S** (2000). The symptom of depression in schizophrenia and its management. *Advances in Psychiatric Treatment* **6**, 169–177.
- Muris P, Meesters C, Diederer R** (2005). Psychometric properties of the Big Five Questionnaire for Children (BFQ-C) in a Dutch sample of young adolescents. *Personality and Individual Differences* **38**, 1757–1769.
- Neumann S, Waldstein SR, Sollers III JJ, Thayer JF, Sorkin JD** (2001). The relation of hostility, rumination, and distraction to cardiovascular reactivity and recovery responses to anger. *Annals of Behavioral Medicine* **23**, 631–640.
- Nijjar R, Ellenbogen MA, Hodgins S** (2014). Personality, coping, risky behavior, and mental disorders in the offspring of parents with bipolar disorder: a comprehensive psychosocial assessment. *Journal of Affective Disorders* **166**, 315–323.
- Nolen-Hoeksema S** (1987). Sex differences in unipolar depression: evidence and theory. *Psychological Bulletin* **101**, 259–282.
- Nolen-Hoeksema S** (1991). Responses to depression and their effects on the duration of depressive episodes. *Journal of Abnormal Psychology* **100**, 569–582.
- Nolen-Hoeksema S** (1998). Contributors to the gender difference in rumination. In *Annual Meeting of the American Psychological Association*, San Francisco, CA.
- Nolen-Hoeksema S** (2000). The role of rumination in depressive disorders and mixed anxiety/depressive symptoms. *Journal of Abnormal Psychology* **109**, 504–511.
- Nolen-Hoeksema S** (2012). Emotion regulation and psychopathology: the role of gender. *Annual Review of Clinical Psychology* **8**, 161–187.
- Nolen-Hoeksema S, Harrell ZA** (2002). Rumination, depression, and alcohol use: tests of gender differences. *Journal of Cognitive Psychotherapy* **16**, 391–403.
- Nolen-Hoeksema S, Parker LE, Larson J** (1994). Ruminative coping with depressed mood following loss. *Journal of Personality and Social Psychology* **67**, 92–104.
- Nolen-Hoeksema S, Stice E, Wade E, Bohon C** (2007). Reciprocal relations between rumination and bulimic, substance abuse, and depressive symptoms in female adolescents. *Journal of Abnormal Psychology* **116**, 198.
- Nolen-Hoeksema S, Watkins ER** (2011). A heuristic for developing transdiagnostic models of psychopathology explaining multifinality and divergent trajectories. *Perspectives on Psychological Science* **6**, 589–609.
- Nolen-Hoeksema S, Wisco BE, Lyubomirsky S** (2008). Rethinking rumination. *Perspectives on Psychological Science* **3**, 400–424.
- Ochsner KN, Gross JJ** (2005). The cognitive control of emotion. *Trends in Cognitive Sciences* **9**, 242–249.
- O'Mahen HA, Karl A, Moberly N, Fedock G** (2015). The association between childhood maltreatment and emotion regulation: two different mechanisms contributing to depression? *Journal of Affective Disorders* **174**, 287–295.
- Orue I, Calvete E, Padilla P** (2014). Brooding rumination as a mediator in the relation between early maladaptive schemas and symptoms of depression and social anxiety in adolescents. *Journal of Adolescence* **37**, 1281–1291.
- Ottaviani C, Shapiro D, Davydov DM, Goldstein IB, Mills PJ** (2009). The autonomic phenotype of rumination. *International Journal of Psychophysiology* **72**, 267–275.
- Padilla Paredes P & Calvete E** (2014). Cognitive vulnerabilities as mediators between emotional abuse and depressive symptoms. *Journal of Abnormal Child Psychology* **42**, 743–753.
- Parker G** (1983). *Parental Overprotection: A Risk Factor in Psychosocial Development*. Grune & Stratton: New York.
- Pavlickova H, Turnbull O, Bentall RP** (2014). Cognitive vulnerability to bipolar disorder in offspring of parents with bipolar disorder. *British Journal of Clinical Psychology* **53**, 386–401.
- Pavlickova H, Turnbull OH, Myin-Germeys I, Bentall RP** (2015). The inter-relationship between mood, self-esteem and response styles in adolescent offspring of bipolar parents: an experience sampling study. *Psychiatry Research* **225**, 563–570.

- Pearson KA, Watkins ER, Mullan EG** (2011). Rejection sensitivity prospectively predicts increased rumination. *Behaviour Research and Therapy* **49**, 597–605.
- Piaget J** (1977). The role of action in the development of thinking. In *Knowledge and Development* (ed. W. F. Overton and J. M. Gallagher), pp. 17–42. Plenum Press: New York.
- Pillai V, Steenburg LA, Ciesla JA, Roth T, Drake CL** (2014). A seven day actigraphy-based study of rumination and sleep disturbance among young adults with depressive symptoms. *Journal of Psychosomatic Research* **77**, 70–75.
- Powers WT** (1973). *Behavior: The Control of Perception*. Aldine: Chicago, pp. 264–265.
- Puterman E, O'Donovan A, Adler NE, Tomiyama AJ, Kemeny M, Wolkowitz OM, Epel E** (2011). Physical activity moderates effects of stressor-induced rumination on cortisol reactivity. *Psychosomatic Medicine* **73**, 604–611.
- Raes F, Smets J, Nelis S, Schoofs H** (2012). Dampening of positive affect prospectively predicts depressive symptoms in non-clinical samples. *Cognition & Emotion* **26**, 75–82.
- Rapado-Castro M, McGorry PD, Yung A, Calvo A, Nelson B** (2015). Sources of clinical distress in young people at ultra-high risk of psychosis. *Schizophrenia Research* **165**, 15–21.
- Ray RD, Ochsner KN, Cooper JC, Robertson ER, Gabrieli JD, Gross JJ** (2005). Individual differences in trait rumination and the neural systems supporting cognitive reappraisal. *Cognitive, Affective, & Behavioral Neuroscience* **5**, 156–168.
- Rea M, Smith A, Bierman A, Figueiro M** (2010). *The Potential of Outdoor Lighting for Stimulating the Human Circadian System*. Lighting Research Center, Rensselaer Polytechnic Institute: New York.
- Rector NA, Roger D** (1996). Cognitive style and well-being: a prospective examination. *Personality and Individual Differences* **21**, 663–674.
- Robillard R, Naismith SL, Smith KL, Rogers NL, White D, Terpening Z, Ip TKC, Hermens DF, Whitwell B, Scott EM, Hickie IB** (2014). Sleep-wake cycle in young and older persons with a lifetime history of mood disorders. *PLoS ONE* **9**, e87763.
- Roelofs J, Huibers M, Peeters F, Arntz A** (2008). Effects of neuroticism on depression and anxiety: rumination as a possible mediator. *Personality and Individual Differences* **44**, 576–586.
- Rothbart M, Bates J** (2006). Temperament. In *Handbook of Child Psychology: vol. 3, Social, Emotional, and Personality Development*, 6th edn (ed. N. Eisenberg, W. Damon and L. M. Richard), pp. 99–166. John Wiley & Sons Inc.: New Jersey.
- Rudolph KD, Hammen C** (1999). Age and gender as determinants of stress exposure, generation, and reactions in youngsters: a transactional perspective. *Child Development* **70**, 660–677.
- Sanislow CA, Pine DS, Quinn KJ, Kozak MJ, Garvey MA, Heisssen RK, Wang PS, Cuthbert BN** (2010). Developing constructs for psychopathology research: research domain criteria. *Journal of Abnormal Psychology* **119**, 631.
- Schmitz TW, Johnson SC** (2006). Self-appraisal decisions evoke dissociated dorsal-ventral aMPFC networks. *Neuroimage* **30**, 1050–1058.
- Schwartz JA, Koenig LJ** (1996). Response styles and negative affect among adolescents. *Cognitive Therapy and Research* **20**, 13–36.
- Scott J** (2016). Psychological interventions for early stage bipolar disorder. In *The Treatment of Bipolar Disorders* (ed. E. Vieta), OUP: Oxford.
- Scott J, Leboyer M, Hickie I, Berk M, Kapczinski F, Frank E, Kupfer D, McGorry P** (2013). Clinical staging in psychiatry: a cross-cutting model of diagnosis with heuristic and practical value. *British Journal of Psychiatry* **202**, 243–245.
- Segerstrom SC, Stanton AL, Alden LE, Shortridge BE** (2003). A multidimensional structure for repetitive thought: what's on your mind, and how, and how much? *Journal of Personality and Social Psychology* **85**, 909.
- Sheline YI, Barch DM, Price JL, Rundle MM, Vaishnavi SN, Snyder AZ, Mintun MA, Wang S, Coalson RS, Raichle ME** (2009). The default mode network and self-referential processes in depression. *Proceedings of the National Academy of Sciences USA* **106**, 1942–1947.
- Simon D, Stenstrom DM, Read SJ** (2015). The coherence effect: blending cold and hot cognitions. *Journal of Personality and Social Psychology* **109**, 369–394.
- Skitch SA, Abela JR** (2008). Rumination in response to stress as a common vulnerability factor to depression and substance misuse in adolescence. *Journal of Abnormal Child Psychology* **36**, 1029–1045.
- Smith JM, Alloy LB** (2009). A roadmap to rumination: a review of the definition, assessment, and conceptualization of this multifaceted construct. *Clinical Psychology Review* **29**, 116–128.
- Spasojevic J, Alloy LB** (2002). Who becomes a depressive ruminator? Developmental antecedents of ruminative response style. *Journal of Cognitive Psychotherapy* **16**, 405–419.
- Stange JP, Hamilton JL, Burke TA, Kleiman EM, O'Garro-Moore JK, Seligman ND, Abramson LY, Alloy LB** (2015). Negative cognitive styles synergistically predict suicidal ideation in bipolar spectrum disorders: a 3-year prospective study. *Psychiatry Research* **226**, 162–168.
- Steinan M, Lagerberg T, Melle I, Andersson O, Morken G, Scott J** (2015). Delayed sleep phase: an important circadian subtype of sleep disturbance in bipolar disorders. *Journal of Affective Disorders* **191**, 156–163.
- Stone LB, Silk JS, Siegle GJ, Lee KH, Stroud LR, Nelson EE, Dahl RE, Jones NP** (2016). Depressed adolescents' pupillary response to peer acceptance and rejection: the role of rumination. *Child Psychiatry & Human Development* **47**, 397–406.
- Sütterlin S, Paap M, Babic S, Kübler A, Vögele C** (2012). Rumination and age: some things get better. *Journal of Aging Research* **2012**, 1–10.
- Tamres LK, Janicki D, Helgeson VS** (2002). Sex differences in coping behavior: a meta-analytic review and an examination of relative coping. *Personality and Social Psychology Review* **6**, 2–30.
- Tan PZ, Forbes EE, Dahl RE, Ryan ND, Siegle GJ, Ladouceur CD, Silk JS** (2012). Emotional reactivity and regulation in anxious and non-anxious youth: a cell-phone ecological momentary assessment study. *Journal of Child Psychology and Psychiatry* **53**, 197–206.

- Thomas J, Bentall RP** (2002). Hypomanic traits and response styles to depression. *British Journal of Clinical Psychology* **41**, 309–313.
- Thompson RJ, Mata J, Jaeggi SM, Buschkuhl M, Jonides J, Gotlib IH** (2010). Maladaptive coping, adaptive coping, and depressive symptoms: variations across age and depressive state. *Behaviour Research and Therapy* **48**, 459–466.
- Thomsen DK, Mehlsen MY, Hokland M, Viidik A, Olesen F, Avlund K, Munk K, Zachariae R** (2004a). Negative thoughts and health: associations among rumination, immunity, and health care utilization in a young and elderly sample. *Psychosomatic Medicine* **66**, 363–371.
- Thomsen DK, Mehlsen MY, Olesen F, Hokland M, Viidik A, Avlund K, Zachariae R** (2004b). Is there an association between rumination and self-reported physical health? A one-year follow-up in a young and an elderly sample. *Journal of Behavioral Medicine* **27**, 215–231.
- Thomsen DK, Yung Mehlsen M, Christensen S, Zachariae R** (2003). Rumination: relationship with negative mood and sleep quality. *Personality and Individual Differences* **34**, 1293–1301.
- Treynor W, Gonzalez R, Nolen-Hoeksema S** (2003). Rumination reconsidered: a psychometric analysis. *Cognitive Therapy and Research* **27**, 247–259.
- Vallarino M, Henry C, Etain B, Gehue LJ, Macneil C, Scott EM, Barbato A, Conus P, Hlastala SA, Fristad M, Miklowitz DJ, Scott J** (2015). An evidence map of psychosocial interventions for the earliest stages of bipolar disorder. *The Lancet Psychiatry* **2**, 548–563.
- Van der Gucht E, Morriss R, Lancaster G, Kinderman P, Bentall RP** (2009). Psychological processes in bipolar affective disorder: negative cognitive style and reward processing. *British Journal of Psychiatry* **194**, 146–151.
- Van Nierop M, van Os J, Gunther N, Myin-Germeys I, de Graaf R, ten Have M, van Dorsseleer S, Bak M, van Winkel R** (2012). Phenotypically continuous with clinical psychosis, discontinuous in need for care: evidence for an extended psychosis phenotype. *Schizophrenia Bulletin* **38**, 231–238.
- Van Rheenen TE, Murray G, Rossell SL** (2015). Emotion regulation in bipolar disorder: profile and utility in predicting trait mania and depression propensity. *Psychiatry Research* **225**, 425–432.
- Verhaeghen P, Joormann J, Aikman SN** (2014). Creativity, mood, and the examined life: self-reflective rumination boosts creativity, brooding breeds dysphoria. *Psychology of Aesthetics, Creativity, and the Arts* **8**, 211.
- Verstraeten K, Vasey MW, Raes F, Bijttebier P** (2009). Temperament and risk for depressive symptoms in adolescence: mediation by rumination and moderation by effortful control. *Journal of Abnormal Child Psychology* **37**, 349–361.
- Watkins ER** (2009). Depressive rumination and comorbidity: evidence for brooding as a trans-diagnostic process. *Journal of Rational-Emotive & Cognitive-Behavior Therapy* **27**, 160–175.
- Watkins ER, Nolen-Hoeksema S** (2014). A habit-goal framework of depressive rumination. *Journal of Abnormal Psychology* **123**, 24.
- Watkins ER, Teasdale JD** (2001). Rumination and overgeneral memory in depression: effects of self-focus and analytic thinking. *Journal of Abnormal Psychology* **110**, 353.
- Weiss NH, Sullivan TP, Tull MT** (2015). Explicating the role of emotion dysregulation in risky behaviors: a review and synthesis of the literature with directions for future research and clinical practice. *Current Opinion in Psychology* **3**, 22–29.
- Willem L, Bijttebier P, Claes L, Raes F** (2011). Rumination subtypes in relation to problematic substance use in adolescence. *Personality and Individual Differences* **50**, 695–699.
- Williams JMG, Barnhofer T, Crane C, Herman D, Raes F, Watkins ER, Dalgleish T** (2007). Autobiographical memory specificity and emotional disorder. *Psychological Bulletin* **133**, 122.
- Wood JV, Saltberg JA, Neale JM, Stone AA, Rachmiel TB** (1990). Self-focused attention, coping responses, and distressed mood in everyday life. *Journal of Personality and Social Psychology* **58**, 1027.
- Yook K, Kim KH, Suh SY, Lee KS** (2010). Intolerance of uncertainty, worry, and rumination in major depressive disorder and generalized anxiety disorder. *Journal of Anxiety Disorders* **24**, 623–628.
- Yung AR, Stanford C, Cosgrave E, Killackey E, Phillips L, Nelson B, McGorry PD** (2006). Testing the ultra high risk (prodromal) criteria for the prediction of psychosis in a clinical sample of young people. *Schizophrenia Research* **84**, 57–66.
- Zoccola PM, Dickerson SS** (2012). Assessing the relationship between rumination and cortisol: a review. *Journal of Psychosomatic Research* **73**, 1–9.
- Zoccola PM, Dickerson SS, Lam S** (2009). Rumination predicts longer sleep onset latency after an acute psychosocial stressor. *Psychosomatic Medicine* **71**, 771–775.