INSTRUCTION TO CONTRIBUTORS

The Journal of Helminthology publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance, but only in exceptional circumstances will systematic or taxonomic studies be acceptable.

Manuscripts, which must be in English or French (with an English summary), should be addressed to:

The Editor, Journal of Helminthology, London School of Hygiene and Tropical Medicine, Winches Farm Field Station, 395 Hatfield Road, St Albans, Herts AL4 0XQ, England.

Two copies of a typescript, on size A4 paper with double spacing, should be submitted Papers should be preceded by a short abstract and will normally have the followin sections: brief Introduction; Materials and Methods; Results; Discussior Acknowledgements; References. However, the form of the paper may vary, dependin on its subject matter; recent past issues should be consulted for a suitable form Research Notes should also be preceded by a brief abstract. Illustrations should b drawn in Indian ink, preferably not more than double the final size. Care should b taken that all illustrations fit into the format of the Journal. The maximum size a illustration will be printed is 12.0×20.0 cm. Where many separate drawings are made some indication of how they may be grouped to make a corporate plate without undu wastage of space, should be indicated. Some indication of scale (preferably a scale bar should normally be given on the figure. Photocopies of illustrations should be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in th Journal (maximum size 12.0×20.0 cm). Composite prints must be mounted and ca have the separate photographs abutting; they will then have a separating line inserte by the printers. All figures and letters on photographs must be inserted by the author

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to b understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. Th following are examples:

DUKE, B. O. L. (1971) The ecology of onchocerciasis in man and animals. In: *Ecolog* and *Physiology of parasites* (editor, A. M. Fallis) pp. 213–222. Adam Hilger Ltd London.

JAMES, C. & WEBBE, G. (1973) A comparison of Egyptian and East African strains c Schistosoma haematobium. Journal of Helminthology, 47, 49–59.

25 offprints are provided free of charge; additional copies may be ordered at the proc stage.

Journal of Helminthology volume 56 • no. 2 • jun

Contents

Pages

Filaricidal properties of a wild herb, Andrographis paniculata. ANITA DUTTA and NIRMAL C. SUKUL	81-84
The effect of controlled release copper sulphate and tributyltin fluoride on the mortality and infectivity of <i>Schistosoma mansoni</i> cercariae. VITHOON VIYANANT, SANAM THIRACHANTRA and SANTASIRI SORNMANI	85–92
Brugia-type microfilariae in the Madras tree shrew Anathana ellioti (Waterhouse). (Research Note). N. C. NANDI	93-94
Scanning electron microscopic observations on the tegumental surfaces of two rumen flukes (Trematoda: Paramphistomata). VEENA TANDON and S. C. MAITRA	95–104
Evaluation of modified Baermann apparatus for the recovery of infective trichostrongylid larvae from herbage samples. (Research Note) S. N. CHIEJINA	105–109
Histochemical and electrophoretic studies on phosphatases of some Indian trematodes. MASOODUL HAQUE and ATHER H. SIDDIQI	111–116
Techniques for examining paramphistomes. (Research Note) S. L. EDUARDO	117-119
Habitat specificity of the acanthocephalan Acanthocephalus clavula (Dujardin, 1845) in eels Anguilla anguilla (L): C. R. KENNEDY and D. LORD	121–129
Recovery of <i>Taenia crassiceps</i> cysticerci (ERS Tol derived strain) from sub-zero temperatures. (Research Note) P. J. HAM	131–133
The pathogenesis of a single challenge dose of <i>Haemonchus contortus</i> in lambs under six months of age. A. R. HUNTER and GRACE MACKENZIE	135–144
Pathophysiology of <i>Mesocestoides corti</i> infection in the mouse. T. R. WHITE, R. C. A. THOMPSON, W. J. PENHALE, D. A. PASS and J. N. MILLS	145-153
The infectivity of microfilariae of <i>Brugia pahangi</i> of different ages to <i>Aedes aegypti</i> . J. C. de HOLLANDA, D. A. DENHAM and R. R. SUSWILLO	155–157
The morphology of <i>Dentostomella kuntzi</i> (Nematoda: Oxyuroidea) from Egyptian rodents. A. A. ASHOUR and J. W. LEWIS	159–168

https://doi.org/10.1017/S0022149X00034258 Published online by Cambridge University Press Printed by Taylor & Francis (Printers) Ltd, Hankine Road, Basingstoke, Hampshire RG24 OPR and published by the London School of Hygiene and Tropical Medicine