



Disclosure: REFERENCES [1] Drăgan-Chirilă, Diana.(24-26.05 2018), Associate Professor Ph.D., University of Art and Design Cluj-Napoca, Romania, visual artist, Coordinator of the multimedia installation “Diagnostique” new media and multimedia performance instal

Keywords: dying is inevitable; emotional disorders; non-verbal communication; art therapy an resilience in per

EPV0226

Negative impact of family religious and spiritual beliefs in schizophrenia – a 2 year follow-up case

J. Mauricio^{1*} and M. Albuquerque²

¹Mental Health, ULSAM, VIANA DO CASTELO, Portugal and

²Mental Health, Hospital de Cascais, Alcabideche, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1810

Introduction: For individuals with mental disorders and their families, religion and spirituality may have a significant influence over how these conditions are understood, managed and treated. Family can act as a moderator in which psychotic patients interpret and explain internalized events. However, they can have a negative impact when discouraging diagnosis and treatment adherence.

Objectives: Explore the impact of family religious and spiritual beliefs on clinical outcomes among a schizophrenic patient. Investigate the psychiatrist’s role in addressing barriers to treatment adherence.

Methods: Data retrieved from clinical interview. Subsequent non-systematic review of the most relevant literature on the topic.

Results: We report a case of a 30-year-old single catholic woman, living with her parents. She had a past history of psychotic symptoms that were interpreted in a context of a depressive episode. After some months she fulfilled the criteria for Schizophrenia and anti-psychotic was prescribed. Family always demonstrated doubts about the disease and negatively influenced the treatment adherence. They believed she was possessed by demons and she was submitted to exorcisms and spiritual therapies. After a 2-year follow-up with erratic treatment regimens and worsening symptoms they accepted her hospitalisation. The majority of symptoms were controlled allowing complete adherence to the same treatment proposed before.

Conclusions: The disease acceptance is a complex process, influenced by multiple beliefs that play different roles in each patient and family, that can adversely influence clinical management. It is essential to understand the family sociocultural environment, by gauging the most influential elements aiming to enhance their compliance with treatment.

Disclosure: No significant relationships.

Keywords: religious beliefs; treatment adherence; schizophrenia; cultural

EPV0227

Transcultural validation of the “regulation of emotions questionnaire” among Tunisian university students

E. Bouguira, U. Ouali*, M. Ben Alaya, Y. Zgueb and F. Nacef

Department Psychiatry A, Razi Hospital, Manouba, Tunisia

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1811

Introduction: The regulation of emotions is a psychological process covering the capacity of inhibiting, maintaining or modulating emotions to create a coherent psychological functioning and to

enable the individual to live in adequacy with his or her environment.

Objectives: The aim of the study was to develop an Arabic version of the “Regulation of Emotions Questionnaire-2” (REQ-2T) from Phillips & Power, 2007, and to validate it in a sample of Tunisian university students.

Methods: This is a validation study conducted in a sample of 384 Tunisian university students to whom we administered the REQ-2T, the DASS-21 questionnaire and a sociodemographic questionnaire. We tested face and content validity, reliability, and construct validity of the Arabic version of the REQ-2.

Results: Face and content validity were satisfying. The internal consistency was average, with Cronbach’s alpha coefficient ranging from 0.44 to 0.65. The inter-dimensional correlation reflected statistically significant and logical correlations within the REQ-2T. Temporal stability was satisfying with a variable concordance of 0.65. Exploratory analysis revealed four factors similarly to the original version of the questionnaire. Statistically significant correlations were found between the REQ-2T and its external validator (DASS-21).

Conclusions: The REQ-2T demonstrated good psychometric properties, thus this scale can be reliably used as a measure of emotion regulation in the Tunisian population.

Disclosure: No significant relationships.

Keywords: emotion regulation; validation; questionnaire; Depression

EPV0228

Why does transcultural consultation matter?

A. Quintão^{1*}, C. Santos¹, M.D. Urzal¹, I. Donas-Boto¹, F. Azevedo¹, M.M. Lemos², F. Coelho² and J. Vian¹

¹Psychiatry Department, Ocidental Lisbon Hospital Center, Lisboa, Portugal and ²Psychiatry Department, North Lisbon Hospital Center, Lisboa, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1812

Introduction: Psychiatrists should be aware of the new challenges and needs that globalization poses.

Objectives: To highlight the need for a culturally sensitive approach to mental health.

Methods: Non-systematic review of “cultural consultation” on PubMed.

Results: Most people assume a direct connection between pathophysiology and clinical symptoms. However, evidence shows that the translation of pathophysiology and psychopathology into specific symptoms is mediated by cognitive processes and social interactions, which reflect models/practices specific to our culture. Patients focus on specific aspects of being sick, reinforced by cultural narratives or to fit expectations. Thus, people from different cultural backgrounds might have trouble communicating; cultural idioms of distress can be misinterpreted. The role of structural violence bestowed upon cultural minorities, which leads to discrimination and social exclusion, has extensively been studied as a risk factor for mental illness. Furthermore, ignoring cultural differences and diversity has been shown to contribute to healthcare

disparities, hampering access to care and diminishing the quality of care received. In Canada, the Cultural Consultation Model provides cultural expertise, either by evaluating patients (preferably accompanied by the referring doctor, a translator and cultural mediator) for 1-3 sessions, providing recommendations to the referring doctor; or providing consulting to a referring doctor or organisation, through general guidance or discussion of specific cases.

Conclusions: In an evergrowing globalization process, we will inevitably have more contact with patients from culturally distinct backgrounds. To provide the best care, we must be aware of the ways in which culture can shape symptom expression, and take into account cultural explanations and preferences.

Disclosure: No significant relationships.

Keywords: transcultural; cultural consultation

EPV0231

Cross-cultural research of coping: Individualization and modernization in Post-Soviet societies

V. Lianguzova^{1*}, D. Khoroshilov² and E. Belinskaya¹

¹Psychology, Moscow State University, Moscow, Russian Federation and ²Faculty Of Psychology, Russian State University for the Humanities, Moscow, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1813

Introduction: The article is devoted to the main methodological problems of qualitative research in the psychology of culture. Methods of ethnographic (field) observation were borrowed by psychology from social and cultural anthropologists in the first half of the twentieth century, and in modern research practices there is a special ethnographic direction that claims to analyze behavior and lifestyle in various subcultures and communities.

Objectives: The purpose of this study was to identify the key cross-cultural difference in coping behavior of respondents from Moscow and Tashkent.

Methods: The methodological basis of the research was the thematic analysis of narratives and free-form interviews. The sample was N = 60 (residents of Russia and Uzbekistan age 17-39).

Results: The results of thematic analysis of narratives and free-form interviews of respondents from Moscow and Tashkent allow us to conclude that the key cross-cultural difference in coping behavior is the degree of its individualization: representatives of Uzbek culture are focused on receiving support and care from significant Others, and not on independent internal work (unlike Russian respondents). At the same time, they are not satisfied with the traditional prescriptions that come from the family environment, which forces them to coping practices that go beyond the boundaries of normative social (often religious) ideas.

Conclusions: This can be interpreted from the point of view of the process of modernization of Uzbek culture, which is gradually becoming individualistic, and the latter circumstance requires the construction of flexible coping strategies in the situation of social and cultural changes.

Disclosure: No significant relationships.

Keywords: Qualitative methods; Culture; Ethnopsychology; coping