uted substantially to the body of scientific data on which our current understanding of droplet nuclei transmission and UV air disinfection is based, can hardly be considered radical. His proposal should be considered very seriously.

REFERENCES

- 1. Centers for Disease Control and Prevention. *Guidelines far Preventing Transmission of Tuberculosis* in Health-Care Settings. 2nd ed. Notice of comment period. *Federal Register* October 12, 1993;58:52810-52854.
- 2. Catanzaro A. Nosocomial tuberculosis. Am Rev Respir Dis 1982; 125:559.
- Hutton MD, Stead WW, Cauthen GM, Bloch AB, Ewing WM. Nosocomial transmission of tuberculosis associated with a draining abscess. *J Infect Dis* 1990;161:286-295.
- Edlin BR, Tokars JI, Grieco MH, et al. An outbreak of multi-drug resistant tuberculosis among hospitalized patients with the acquired immunodeficiency syndrome. N Engl J Med 1992;326:1514.
- Cousell SR, Tan JS, Dittus RS. Unsuspected pulmonary tuberculosis in a community teaching hospital. Arch Intern Med 1989;149:1274-1278.
- Mathur P, Sacks L, Auten G, Sall R, Levy C, Gordin F. Delayed diagnoses of pulmonary tuberculosis in city hospitals. Arch Intern Med 1994;154:306-310.
- 7. OSHA enforcement policy and procedures for occupational exposure to tuberculosis. *Infect Control Hosp Epidemiol* 1993;14:694-699.
- 8. McGowan JE. Implementing administrative controls, risk assessment analysis of healthcare workers' skin test data, and transmission among patients and health care workers. Presented at the American Society for Hospital Engineering of the American Hospital Association: TB and the Health Care Environment;

January 31-February 1, 1994; Dallas, Texas.

- 9. Scott B, Schmid M, Nettleman MD. Early identification and evaluation of inpatients at high risk for tuberculosis. *Arch Intern Med* 1994;154:326-330.
- Iseman MD. A leap of faith: what can we do to curtail intrainstitutional transmission of tuberculosis? Ann Intern Med 1992;117:251. Editorial.
- 11. Turner M, McGowan K, Cuchural GJ, et al. Environmental controls and PPD conversion rates in an inpatient tuberculosis unit. Accepted for presentation at the 1994 American Lung Association/American Thoracic Society International Meeting; May 22:25, 1994; Boston, Massachusetts. Abstract.
- Riley RL, Nardell EA. Clearing the air: the theory and application of ultraviolet air disinfection. Am Rev *Respir Dis* 1989;139:1286.
- Macher JM. The use of germicidal lamps to control tuberculosis in healthcare facilities. Infect Control Hosp Epidemiol 1993;14:723-729.
- 14. Nardell EA. Fans, filters, or rays? Pros and cons of the current environmental tuberculosis control technologies. Infect Control *Hosp Epidemiol* 1993;14:681-685.
- International Agency for Research on Cancer. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans: Solar and Ultraviolet Light, vol. 55. Lyons, France: World Health Organization; 1992:227-228.
- Nardell EA, Riley RL. A new ultraviolet germicidal irradiation (UVGI) fixture design for upper room air disinfection with low ceilings. In: *Program and Abstracts of the World Congress on Tuberculosis.* Bethesda, MD: National Institutes of Health; 1992:38.
- 17. Permutt S. The impact of ultraviolet light on transmission of infection. Presented at the American College of Chest Physicians/ American Thoracic Society/National Heart, Lung, and Blood Institute Consensus Conference: Institutional Infection Control Measures for Tuberculosis in the Era of Multidrug Resistance; November 13, 1993; Chicago, Illinois.

Computer-Based Epidemiologic Case Study Available from CDC

by Gina Pugliese, RN, MS Medical News Editor

The Centers for Disease Control and Prevention has developed an IBM-compatible, computerbased epidemiologic case study, "Investigating an Outbreak: Pharyngitis in Louisiana," to teach outbreak investigation skills to public health workers. The case study simulates a real outbreak investigation in which the student plays the role of the lead investigator. Through the case study, students learn to apply appropriate outbreak investigation practices, perform a variety of epidemiologic tasks (eg, creating a line list, developing a case definition, and calculating attack rates), recognize the role of the laboratory in an outbreak investigation, and understand the applications of EpiInfo computer software. Additional information is available by calling (800) 418-7246.