
INSTRUCTIONS TO CONTRIBUTORS

The *Journal of Helminthology* publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance, but only in exceptional circumstances will systematic or taxonomic studies be acceptable.

Manuscripts, which must be in English or French (with an English summary) should be accompanied by a letter signed by *all* the authors and should be addressed to:

The Editors, Journal of Helminthology,
London School of Hygiene and Tropical Medicine,
395A Hatfield Road,
St Albans, Herts AL4 0XQ.
England.

Three copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief Introduction; Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Research Notes should also be preceded by a brief abstract. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration will be printed is 12·0×20·0 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space, should be indicated. Some indication of scale (preferably a scale bar) should normally be given on the figure. Photocopies of illustrations should be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size 12·0×20·0 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

ASHFORD, R. W. & BARNISH, G. (1989) *Strongyloides fuelleborni* and similar parasites in animals and man. In: *Strongyloidiasis: a major roundworm infection of man* (editor, D. I. Grove) pp. 271–286. Taylor & Francis: London.

GARSDIE, P. & BEHNKE, J. M. (1989) Primary infection of hamsters with *Ancylostoma ceylanicum*. *Parasitology*, **98**, 283–289.

Contents

	Pages
The daily cercarial emission rhythm of <i>Schistosoma margrebowiei</i> with particular reference to dark period stimuli. K. RAYMOND and A. J. PROBERT	159-168
Studies on the infectivity of <i>Diplostomum spathaceum</i> in rainbow trout (<i>Oncorhynchus mykiss</i>). S. K. WHYTE, C. J. SECOMBES and L. H. CHAPPELL	169-178
Effect of water temperature on the ability of <i>Diplostomum spathaceum</i> miracidia to establish in lymnaeid snails. G. D. B. WAADU and L. H. CHAPPELL	179-185
Some enzymes of carbohydrate metabolism in <i>Mesocestoides corti</i> and <i>Heterakis spumosa</i> . P. DUBINSKÝ, B. RUŠCINOVÁ, S. L. HETMANSKI, C. ARME, L. TURČEKOVÁ and M. RYBOŠ	187-192
<i>In vitro</i> stress selection of nematophagous fungi for biocontrol of parasitic nematodes in ruminants. M. LARSEN, J. WOLSTRUP, S. A. HENRIKSEN, C. DACKMAN, J. GRØNVOLD and P. NANSEN	193-200
Survival of sheep and goat first stage protostrongylid larvae in experimental conditions: influence of humidity and temperature. J. CABARET, S. RISYE RISEANI and E. BAEZA	201-207
Chromosome numbers in <i>Onchocerca dukei</i> and <i>O. tarsicola</i> . R. J. POST, O. BAIN, and S. KLÄGER	208-210
<i>Acanthocheilonema viteae</i> (<i>Dipetalonema viteae</i>) in mice: differences in the relative binding of microfilarial surface-specific antibody may explain the contrasting response phenotypes of BALB/c and C57BL/10. A. J. GATRILL, J. KEE, J. M. BEHNKE and D. WAKELIN	211-218
Recovery of adult stages and microfilaraemia after low dose inoculation of third stage larvae of <i>Litomosoides carinii</i> in <i>Sigmodon hispidus</i> . P. WENK and J. MÖSSINGER	219-225
The effectiveness of combined control measures on the prevalence of guinea worm disease in Anambra State, Nigeria. O. C. NWAORGU	226-231
First record of the larval parasitic nematode <i>Rhabditis orbitalis</i> from Japanese wood mice (<i>Apodemus</i> spp.) W. SUDHAUS and M. ASAKAWA (Research Note)	232-233
Use of acetic orcein as a fast method to relax, fix, clear and stain cercariae for light microscopy. S. C. WEBB (Research Note)	234-238
Book review	186