

## AN APPLICATION OF RITT'S LOW POWER THEOREM

MICHIHIKO MATSUDA

### Abstract

Consider an algebraic differential equation  $F = 0$  of the first order. A rigorous definition will be given to the classical concept of "particular solutions" of  $F = 0$ . By Ritt's low power theorem we shall prove that a singular solution of  $F = 0$  belongs to the general solution of  $F$  if and only if it is a particular solution of  $F = 0$ .

### §0. Introduction

Let  $k\{y\}$  be the differential polynomial algebra in a single indeterminate  $y$  over an algebraically closed differential field  $k$  of characteristic zero, and  $F$  be an algebraically irreducible element of  $k\{y\}$  of the first order. The totality  $\Pi$  of those elements  $A$  of  $k\{y\}$  such that the remainder of  $A$  with respect to  $F$  is zero is an essential prime divisor of the perfect ideal  $\{F\}$  in  $k\{y\}$  generated by  $F$ . Let  $\Pi, \Sigma_1, \dots, \Sigma_s$  be the essential prime divisors of  $\{F\}$ . Then, each of the  $\Sigma_i$  contains the separant  $S$  of  $F$  (Cf. [5, pp. 30-32]). Take and fix a universal extension  $\Omega$  of  $k$ , the existence of which was proved by Kolchin [3, p. 771]. The manifold of  $\Pi$  in  $\Omega$  is called the general solution of  $F$ . A zero of  $F$  in  $\Omega$  is called a singular solution of  $F = 0$  if it is a zero of  $S$ . The manifold of  $\Sigma_i$  in  $\Omega$  consists of a single point for each  $i$  (Cf. [5, p. 63]). A singular solution of  $F = 0$  is an element of  $k$ , because it is either a zero of the discriminant of  $F$  with respect to  $y'$  or a zero of the initial of  $F$ .

Take a generic point  $w$  of the general solution of  $F$ . Then,  $w$  is transcendental over  $k$ . Hence,  $k(w, w')$  is a one-dimensional algebraic function field over  $k$ , which will be denoted by  $K$ . We shall give a rigorous definition to the classical concept of "particular solutions" of  $F = 0$  as follows (Cf. [1, p. 257]):

**DEFINITION.** A singular solution  $\eta$  of  $F = 0$  will be called a partic-

Received September 6, 1976.

ular solution of  $F = 0$  if there exists a prime divisor  $P$  of  $K$  such that

$$(1) \quad \nu_P(w' - \eta') \geq \nu_P(w - \eta) > 0,$$

where  $\nu_P$  is the normalized valuation belonging to  $P$ .

This definition is independent of the choice of a generic point  $w$  of the general solution of  $F$ .

By Ritt's low power theorem we shall prove the following:

**THEOREM.** *A singular solution  $\eta$  of  $F = 0$  belongs to the general solution of  $F$  if and only if  $\eta$  is a particular solution of  $F = 0$ .*

### §1. Proof of Theorem

Suppose that  $\eta$  is a singular solution of  $F = 0$ . Then,  $\eta$  is an element of  $k$ . Let  $G$  denote the polynomial in  $u, v$  obtained from  $F$  by the replacement of  $y = u + \eta$ ,  $y' = v + \eta'$ . Suppose that

$$G = a_0(u)v^n + a_1(u)v^{n-1} + \cdots + a_n(u),$$

where the  $a_i$  are elements of  $k[u]$ . Unless  $a_i = 0$ , we define  $s_i$  as the least exponent of  $u$  in  $a_i$ . If  $a_i = 0$ , we do not define  $s_i$ . For  $i = n$ ,  $s_n$  can be defined, and  $s_n > 0$ . The following lemma is a corollary of Ritt's low power theorem (Cf. [5, p. 65]):

**LEMMA.** *The singular solution  $\eta$  belongs to the general solution of  $F$  if and only if we have the inequality*

$$(2) \quad s_n \geq s_i + n - i$$

for some  $i$  different from  $n$  ( $0 \leq i < n$ ).

Let us make Puiseux diagram in  $G$ . Then, we have rational numbers  $\mu_1, \dots, \mu_m$  and subscripts  $i_0, i_1, \dots, i_m$  of the  $a$  such that they satisfy the following four conditions:

- (i)  $0 \leq i_0 < i_1 < \cdots < i_m = n$ ;
- (ii)  $0 < \mu_1 < \cdots < \mu_m$ ;
- (iii) for each  $j$  ( $1 \leq j \leq m$ ),

$$(3) \quad s_p + \mu_j(n - p) = s_q + \mu_j(n - q), \quad p = i_{j-1}, \quad q = i_j;$$

- (iv)  $s_i + \mu_j(n - i) \geq \tau_j$

for all  $i, j$  ( $0 \leq i \leq n, 1 \leq j \leq m$ ), where  $\tau_j$  is the number given by the equality (3).

Let  $P$  be a prime divisor of  $K$  such that

$$(4) \quad \nu_P(w - \eta) > 0, \quad \nu_P(w' - \eta') > 0.$$

Then, we have

$$(5) \quad \nu_P(w' - \eta') = \mu_h \nu_P(w - \eta)$$

for some  $h$ . Conversely, for each  $h$  ( $1 \leq h \leq m$ ), there exists some prime divisor  $P$  of  $K$  which satisfies (4) and (5) (Cf. [2, Chap. 2], [4, Chap. 13]).

Because of (ii), there exists a prime divisor  $P$  of  $K$  satisfying (1) if and only if  $\mu_m \geq 1$ . The inequality (2) holds for some  $i$  different from  $n$  ( $0 \leq i < n$ ) if and only if  $\mu_m \geq 1$ . Hence, we have our Theorem by Definition and Lemma.

## §2. An example

Let  $k_0$  be an algebraically closed field of characteristic zero, and  $k_0(x)$  be the one-dimensional rational function field over  $k_0$ . We set  $x' = 1$ , and  $a' = 0$  for all elements  $a$  of  $k_0$ . Suppose that  $k$  is the algebraic closure of  $k_0(x)$ , and that

$$F = x^2(y')^2 + (2x + y)yy' + y^2.$$

Then, the singular solutions of  $F = 0$  are 0 and  $-4x$ . The former is a particular solution of  $F = 0$ , and the latter is not.

Let  $t$  denote  $x + w/w'$ . Then,  $t$  is a constant. We have  $w = t^2(x - t)^{-1}$  and  $w' = -t^2(x - t)^{-2}$ . Hence,  $k(w, w') = k(t)$  with  $t' = 0$ .

## BIBLIOGRAPHY

- [ 1 ] A. R. Forsyth, *Theory of differential equations, Part II, Ordinary equations, not linear*, Vol. II, Cambridge Univ. Press, London, 1906.
- [ 2 ] K. Iwasawa, *Theory of algebraic function fields (in Japanese)*, Iwanami Shoten, Tokyo, 1952.
- [ 3 ] E. R. Kolchin, *Galois theory of differential fields*, *Amer. J. Math.*, **75** (1953), 753–824.
- [ 4 ] É. Picard, *Traité d'analyse, Tome II, 2<sup>e</sup> Édition*, Gauthier-Villars, Paris, 1905.
- [ 5 ] J. F. Ritt, *Differential algebra*, *Amer. Math. Soc. Colloq. Publ. Vol. 33*, New York, 1950.

*Department of Mathematics  
Osaka University*