

Brucellosis as a neglected disease in a neglected population: a seroepidemiological study of migratory nomads in the Fars province of Iran

B. HONARVAR¹, M. MOGHADAMI², K. B. LANKARANI¹,
M. A. DAVARPANAH^{2*}, M. ATAOLAH², A. FARBOD³, E. ESKANDARI³,
M. PANAHI³, A. GHORBANI³, Z. ZAHIRI³, R. TABRIZI¹, M. POURJAFAR⁴
AND S. M. M. HEIDARI⁴

¹Health Policy Research Centre, Shiraz University of Medical Sciences, Shiraz, Iran

²Shiraz HIV/AIDS Research Centre, Shiraz University of Medical Sciences, Shiraz, Iran

³Eghlid Health Network, Shiraz University of Medical Sciences, Shiraz, Iran

⁴Department of Clinical Science, School of Veterinary Medicine, Shiraz University, Shiraz, Iran

Received 13 March 2016; Final revision 20 September 2016; Accepted 15 October 2015;
first published online 21 November 2016

SUMMARY

This study assessed the seroprevalence of brucellosis and its risk factors in migratory nomads in the Fars province of Iran. Active brucellosis was defined as the combination of clinical symptoms, including fever, chills, night sweats, headache, low back pain, arthralgia, or myalgia, and positive laboratory testing, including either a serum agglutination test (SAT) $\geq 1:80$ with a 2-mercaptoethanol (2-ME) test $\geq 1:40$, or a SAT $< 1:80$ combined with a positive Coombs Wright test (CWT) at a titre of at least threefold higher than SAT titre results. For the 536 participants, the female (316, 59%) to male (220, 41%) ratio was 1.4 and the participants' mean age was 32.4 \pm 18.9 (range 1–96) years. Of all participants, 325 (60.6%) showed clinical symptoms; in symptomatic participants, the Rose Bengal plate test was positive in 33 (6.1%) cases, the SAT was positive in 18 (3.3%) cases, and the 2-ME test was positive in 30 (5.5%) cases. Positive SAT and 2-ME results were seen in 18 (3.3%) cases, but a negative SAT and a positive CWT were found in 36 (6.7%) cases. As a result, active brucellosis was detected in 54 cases, indicating a prevalence of 10% (95% confidence interval 8–12). In conclusion, we determined that brucellosis is a prevalent yet neglected disease in this nomadic population. Brucellosis control is not possible as long as these high-risk populations remain neglected.

Key words: Brucellosis, migratory, nomad, neglected, serology, study.

INTRODUCTION

Brucellosis is one of the most common zoonotic infections worldwide and is a public health and economic burden in most developing countries [1]. Globally,

more than 500 000 new cases of brucellosis are reported annually [2–5]. However, the incidence rate of this disease varies from < 0.01 to > 200 cases/100 000 people, depending on the population [6], and the incidence is far higher in occupationally at-risk groups, such as agro-pastoral communities [6].

In Iran, around 16 000 cases of brucellosis are reported annually; this imposes a huge burden on the country [7, 8]. This burden is augmented by under-detection and underestimation of the disease due to

* Author for correspondence: Dr M. A. Davarpanah, Shiraz HIV/AIDS Research Centre, Shiraz University of Medical Sciences, 7134853185 Shiraz, Iran.
(Email: davarpanah@sums.ac.ir)

inefficient surveillance systems for both humans and animals [6]. The migratory nomadic population numbers 1·2 million in Iran, most of whom endure poor living conditions; such as poor access to clean water. In addition, their pastoralist lifestyle entails regular contact with livestock, as they are involved in importing and exporting herds to other regions, and these individuals play an important role in producing, processing, and handling dairy products. In this study, we aimed to learn the prevalence and risk factors for brucellosis in this nomadic population in order to provide data for improved, evidence-based planning and interventions in the fight against this disease.

MATERIALS AND METHODS

General protocol

A 4-month (June–September 2015) cross-sectional sero-study was conducted to determine the prevalence of brucellosis in Iranian nomadic pastoralists. Eghlid, located in the north of the Fars province of Iran, is one of the most important nomadic poles and hosts a large seasonal migratory nomadic population of 10 000 people. This population lives in 2000 tents, occupying 19 regions that cover 7000 km² while cohabitating with half a million livestock, mainly sheep and goats. Their pattern of migration takes them from warm areas in the south of Fars province to Eghlid city in spring, and eventually to hot areas in autumn. Each of the 19 regions in Eghlid maintains a healthcare centre that gives each nomadic family a unique code.

The required sample size for this study was calculated to be 578 participants, assuming an estimated 50% prevalence of brucellosis, a 95% confidence interval (CI), 5% error, and, due to the sampling method, a design effect of 1·5. To achieve a proportional-based systematic randomized sampling, we defined the sample size for each of the 19 regions based on their population size; we selected 18 individuals from the region with the smallest population up to 95 individuals from the region with the largest population. In order to find eligible individuals, families were first selected through a systematic randomized approach using the unique number that was given to each family by the regional healthcare centres. Then, one person from each family was selected by simple randomization and was interviewed in his/her tent.

Eleven teams were organized to conduct this study. Each team consisted of two endemic health sector

staff members: an interviewer and a phlebotomist. The central laboratory of Eghlid city was also prepared to conduct laboratory tests according to the defined protocols. All teams were trained regarding the aims of this study, how to correctly complete the data collection forms, and regarding blood sampling and proper sample handling. The coded data collection form consisted of 31 questions designed to assess demographic characteristics, symptoms related to brucellosis (including fever, chills, sweating, headache, low back pain, arthralgia, myalgia), previous history of brucellosis or its treatment, number of livestock owned and their brucellosis vaccination status, history of animal abortion in the year prior to this study, participation of interviewees in the production, processing, and handling of dairy products, and the accessibility and use of personal protective equipment. The form also included the results of the following laboratory tests: the Rose Bengal plate test (RBPT), the serum agglutination test (SAT) test, the 2-mercaptoethanol (2-ME) test, and the Coombs Wright test (CWT). The forms were completed via face-to-face interviews in each tent after explaining the purpose of the research and obtaining consent. In the case of children, their mothers were interviewed. A 3-ml blood sample was taken from each participant and placed in blood tubes that were labelled with codes that matched the data collection forms. Blood samples were transported to the laboratory maintaining cold chain standards.

Laboratory protocol

A total of 536 blood samples were collected. The sera were separated by centrifuging the blood samples at 2000 *g* and kept at –20 °C until testing. The RBPT, SAT, 2-ME, and CWT were performed on all samples using whole-cell antigen produced by the Pasteur Institute for Health Sciences Research Company (Iran) based on their established protocol [9, 10]. For the RBPT, any degree of agglutination in bright light was interpreted as positive, and no agglutination was interpreted as negative [9]. The SAT was performed in eight sequential serum dilutions of 1:20, 1:40, 1:80, 1:160, 1:320, 1:640, 1:1280, and 1:2560, and the highest dilution with 50% agglutination was reported. An agglutination titre \geq 1:80 was considered positive [9, 10]. In the 2-ME test, eight sequential serum dilutions of 1:20 to 1:2560 were assessed, and titres \geq 1:40 were considered positive [9]. In the presence of SAT < 1:80, CWT was considered positive if it

yielded a 50% agglutination at a titre at least threefold higher than the SAT titre [9].

Diagnosis protocol

Taking into account the pastoral lifestyle of the nomad participants, symptomatic persons with positive SAT and positive 2-ME tests, or persons with a negative SAT and a positive CWT, were diagnosed with active brucellosis [9, 11].

Data entry and analysis

The data from completed forms were extracted and entered into SPSS software v. 20 (IBM Corp., USA). The accuracy of data entry was ensured by checking completed forms against their corresponding data entered in the SPSS files. χ^2 and Mann-Whitney *U* tests were used for univariable analysis. The variables that met inclusion criteria for the regression model ($P \leq 0.2$) were entered into the logistic regression model. In the logistic regression (enter method), *P* values < 0.05 were considered significant. Agreement ratios between serological tests were also calculated using a kappa formula.

Ethical statement

Strict ethical standards were maintained, including voluntary consent to participate, patient confidentiality of laboratory results, and standard, free-of-charge treatment for participants diagnosed with brucellosis. Furthermore, the research protocol of this study was approved by the Ethics Committee of the Health Policy Research Centre affiliated with Shiraz University of Medical Sciences. Finally, the authors assert that all procedures related to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

RESULTS

Of the 578 selected samples, 536 were included in this study, which reflects a 93% participation rate. The median age of interviewees was 30 (range 1–96) years. The female (316, 59%) to male (219, 41%) ratio was 1.4. The median family size of the participants was five and the median number of livestock per family was 100. A large proportion of the

Table 1. *Characteristics of the nomadic population in the seroepidemiological study of brucellosis in Fars province, Iran*

Characteristic	Item	Total
Demographic	Gender, <i>n</i> (%)	
	Male	219 (41)
	Female	316 (59)
	Age (years)	
	25th centile	18
	50th centile	30
	75th centile	44
	Education, <i>n</i> (%)	
	Illiterate	188 (35.1)
	Primary school	189 (35.3)
	Secondary school	68 (12.7)
	High school	60 (11.2)
	University	23 (4.3)
Livestock	Family size (<i>n</i>)	
	25th centile	4
	50th centile	5
	75th centile	6
	Livestock per family (<i>n</i>)	
	25th centile	60
	50th centile	100
	75th centile	200
	Herds' brucellosis vaccination (%)	
	25th centile	0
	50th centile	100
	75th centile	100
	Herd's abortion in the year prior to this study (<i>n</i>)	
25th centile	1	
50th centile	4	
75th centile	10	
Protective equipment	Accessibility, <i>n</i> (%)	
	Yes	30 (5.6)
	No	506 (94.4)
	Application, <i>n</i> (%)	
Yes	23 (4.3)	
No	513 (95.7)	

participants were illiterate (188, 35.1%) or reported education up to primary-school level (189, 35.3%) (Table 1). Of all participants, 449 (83.7%) reported involvement in animal grazing or dairy production, contact with aborted animal fetuses, or cleaning of corrals. In addition, 132 (24.6%) participants reported that their livestock had not been vaccinated against brucellosis. Furthermore, 282 (52.6%) participants reported a median abortion rate of four animals in their livestock during the year prior to this study. More than 506 (94.4%) responders denied owning or

Table 2. Results of brucellosis serological tests in the studied nomadic population of Fars province, Iran ($n = 536$)

	Symptomatic n (%)	Non-symptomatic n (%)	Total n (%)	χ^2 (P value)
Single test				
RBPT				
Positive	33 (6.1)	13 (2.4)	46 (8.5)	2.5 (0.1)
Negative	292 (54.4)	197 (36.7)	489 (91.2)	
SAT				
$\geq 1:80$	18 (3.3)	8 (1.4)	26 (4.8)	0.8 (0.3)
$< 1:80$	306 (57)	202 (37.6)	508 (94.7)	
2-ME				
$\geq 1:40$	30 (5.5)	14 (2.6)	44 (8.2)	1.7 (0.1)
$< 1:40$	250 (46.6)	182 (33.9)	432 (80.5)	
Combined tests				
SAT $\geq 1:80$ and 2-ME $\geq 1:40$				
Yes	18 (3.3)	8 (1.4)	26 (4.8)	0.8 (0.3)
No	306 (57)	202 (37.6)	508 (94.7)	
SAT negative and Coombs positive*				
Yes	36 (6.7)	22 (4.1)	58 (10.8)	0.05 (0.8)
No	288 (53.7)	188 (35)	476 (88.8)	

RBPT, Rose Bengal plate test; SAT, serum agglutination test; 2-ME, 2-mercaptoethanol agglutination test;

* Coombs test was considered positive by forming a 50% agglutination at a titre of at least threefold higher than the SAT titre $< 1:80$.

using protective equipment, such as masks and gloves, during animal contact (Table 1).

Out of all participants, 325 (60.6%) reported clinical symptoms. In symptomatic persons, the RBPT was positive in 33 (6.1%) cases, the SAT was positive in 18 (3.3%) cases, and the 2-ME was positive in 30 (5.5%) cases. Moreover, 18 (3.3%) patients were positive according to both the SAT and 2-ME, and 36 (6.7%) patients were positive for both the SAT and CWT. In total, active brucellosis was detected in 54 participants, indicating a prevalence of 10% (95% CI 8–12). None of these patients had been previously diagnosed or treated for brucellosis. No significant association was noted between serological results and the presence or absence of clinical symptoms (Table 2).

Univariable analysis showed that the prevalence of brucellosis was associated with the age of patients ($P = 0.01$) and the number of abortions in the livestock ($P = 0.01$) (Table 3). According to logistic regression, however, there were no significant associations between brucellosis and these variables (Table 3). The kappa formula also demonstrated the following agreement ratios: RBPT/SAT: -67% ($P < 0.001$), RBPT/2-ME: -92% ($P < 0.001$), RBPT/active brucellosis: -29% ($P < 0.001$), and SAT/2-ME: -72% ($P < 0.001$). However, there was no significant

agreement between CWT and each of RBPT, SAT or 2-ME tests ($P \geq 0.1$).

DISCUSSION

Brucellosis has been eradicated from Norway (1952), Sweden (1957), Finland (1960), Denmark (1962), and England (1981) as well as The Netherlands, Austria, Japan, and Bulgaria (1985) [9]. However, this disease as a re-emergent zoonotic infectious disease is a major worldwide public health concern that is associated with a high economic burden [11, 12]. Brucellosis has a very broad range of clinical manifestations from asymptomatic infection to a serious debilitating disease with diverse complications; therefore its diagnosis may be challenging [7, 12]. In Iran, brucellosis is an endemic disease, and despite improvement in its control, 16 000 cases of this disease are reported each year [13–16].

The Iranian migratory nomadic population, known as ‘Ashaier’ people, comprise around 1 200 000 individuals (1.7% of the total population of Iran), most of whom practice a pastoralist lifestyle. This population manages > 24 million sheep and goats, and produce 20% of the red meat and a significant proportion of all dairy products in Iran. Fars province, in the southwest of Iran, includes a large proportion (12% or

Table 3. Results of univariable analysis in the study of brucellosis of nomadic population in Fars province, Iran

Determinant	Active brucellosis		Statistics	P value	OR	95% CI
	Yes (n = 54)	No (n = 482)				
Age (years)			MWU = 10 150	0.01		
Median	35.5	29				
25th centile	24	17				
50th centile	35.5	29				
75th centile	52.5	43				
Gender (n)			$\chi^2 = 0.8$	0.3	1.3	0.7–2.3
Male	19	199				
Female	35	280				
Education (n)			$\chi^2 = 6.7$	0.2		
Illiterate	23	164				
Primary school	13	176				
Middle school	10	58				
High school	3	57				
University	2	20				
Family size (n)			MWU = 10 812.5	0.4		
Median	5	5				
25th centile	3	4				
50th centile	5	5				
75th centile	7	6				
Livestock per each family (n)			MWU = 12 120	0.4		
Median	100	100				
25th centile	70	60				
50th centile	100	100				
75th centile	200	200				
Livestock brucellosis vaccination (%)			MWU = 11 479.5	0.2		
Median	100	100				
25th centile	100	0.0				
50th centile	100	100				
75th centile	100	100				
Livestock abortion (n)			MWU = 8946	0.01		
Median	6	4				
25th centile	1.2	1				
50th centile	6	4				
75th centile	15	10				

OR, Odds ratio; CI, confidence interval; MWU, Mann–Whitney *U* test.

15 000 individuals) of this population, who collectively own 4 million livestock, mainly sheep and goats.

In this study, we found that 10% of this migratory nomadic population had active brucellosis, a rate that is far higher than the brucellosis prevalence (15.9/100 000) in the general population [9]. Their nomadic lifestyle and their close and continuous contact with livestock make brucellosis an endemic issue for this population that is challenging to diagnose [17]. Moreover, as this survey revealed, current interventions aimed at controlling brucellosis are insufficient due to delays in the detection of brucellosis patients via routine health system activities, a high abortion rate in livestock, low access and use of

protective equipment, and incomplete brucellosis vaccination of livestock. In addition, we did not find any evidence that brucellosis vaccination of livestock in our region is effective. The health of this population is further complicated as a result of herds imported from neighbouring provinces with undefined brucellosis infection or brucellosis vaccination status, weaknesses in quarantine systems at provincial borders [18], lack of herd insurance coverage, inadequate compensation for killed infected animals [18], the practice of mixing animals who have recently experienced an abortion with healthy animals, drought in the settling areas of nomads, and low levels of education in the nomads [19].

There is some lack of consensus regarding the true incidence of brucellosis throughout Iran. One study conducted in rural areas of Isfahan, in central Iran, revealed that the incidence of brucellosis was 8.2/100 000 persons and that 15- to 20-year-old male subjects were more likely (2.1 times) to have this disease [14]. This finding contrasts with our results; the nomadic participants in our survey exhibited a very high prevalence of brucellosis without a significant gender difference. However, a separate study in the Kurdistan province, west of Iran, found a brucellosis incidence rate of 6–12% in the hunters, butchers, and healthcare workers of that area, which is similar to the findings in our study [20]. In Khuzestan, south-west of Iran, the prevalence of brucellosis was found to be 6.3% in nomads, 1.5 times lower than the rate found in this investigation [21]. On the other hand, a seroprevalence study of nomads in Lorestan province, west of Iran, found that 29.5% and 21.1% of nomads had positive SAT and 2-ME results, respectively [22]. In contrast, 3.3% of our participants tested positive according to the SAT and 5.5% tested positive according to the 2-ME test. Studies on this topic also disagree regarding the association between age and prevalence of brucellosis. While our study found no association between age and prevalence, the 10–19 years age group in the Lorestan study [22] and the 15–44 years age group in the Azna study [8] demonstrated a higher prevalence of brucellosis.

Studies in Azerbaijan and Uganda have proven the efficacy of herd immunization in controlling brucellosis [1, 23]. Evidence also suggests that brucellosis control is not achievable unless at-risk groups are educated about this disease in both humans and animals; moreover, this education must involve a serious commitment on the part of both the public and private sectors [23, 24]. Furthermore, studies indicate that health providers must look beyond index cases, as family members of brucellosis patients may also be in contact with common sources of infection and are therefore at risk of acquisition of this disease [25–27]; as such, familial screening in brucellosis cases may lead to early detection and prevention of brucellosis-related complications [8, 26].

We found a two-thirds agreement ratio between RBPT and SAT and a one-third agreement ratio between RBPT and active brucellosis, indicating that brucellosis screening should not be based only on RBPT results. This finding is concurrent with that obtained in another study conducted in northeast Iran [28]. The lack of a strong association between

clinical symptoms and serological tests, as shown in this study, is supported by results from another study in which a series of patients underwent serological tests after confirmation of acute brucellosis [11].

Overall, our work indicates that routine health service activities cannot control brucellosis in the nomadic population in this region, and that a well-integrated surveillance system is needed. This finding is supported by evidence from other studies [8, 15, 16, 29]. Annual active screening for brucellosis in nomadic populations is recommended before the establishment of a surveillance system. In addition, complete treatment of brucellosis patients, as well as screening for their family members, is an important component of a comprehensive brucellosis control programme. Additionally, we recommend implementing programmes to train nomads about this disease, including its transmission and prevention, as well as programmes that provide protective equipment to at-risk populations, support efforts for regular and complete brucellosis vaccination of livestock, and monitor brucellosis vaccination efficacy within herds. Additionally, one study [22] highlighted the importance of research-based field investigations in the fight against brucellosis. Finally, 5% of the screened population in our study demonstrated serological evidence of brucellosis despite being asymptomatic, and only 1/5 members of this group reported a previous history of treatment for brucellosis. Therefore, this group should be followed for possible development of clinical brucellosis.

Our study has a few limitations. Owing to the administrative gap between human and veterinary services in this area, we could not assess brucellosis in the livestock of brucellosis patients. Future studies that address this issue could provide more evidence regarding the risk factors for this disease in nomadic populations.

In conclusion, brucellosis is a common but neglected disease in the Iranian nomadic population. True brucellosis control will not be possible without a national commitment from all stakeholders, including communities as well as human and veterinary services.

ACKNOWLEDGEMENTS

We thank the nomadic populations for their honest participation as well as the staff of the health centres of Eghlid city, Fars province, Iran for assisting us in performing this study.

DECLARATION OF INTEREST

None.

REFERENCES

1. **Kracalik IT, et al.** Human brucellosis trends: re-emergence and prospects for control using a one health approach in Azerbaijan (1983–2009). *Zoonoses and Public Health* Published online: 25 September 2015. doi:10.1111/zph.12229.
2. **Asiimwe BB, Kansiime C, Rwego IB.** Risk factors for human brucellosis in agro-pastoralist communities of south western Uganda: a case-control study. *BMC Research notes* 2015; **4**: 405.
3. **Eini P, et al.** Depressive disorder among brucellosis patients in Hamadan, Iran: a case-control study. *Life Science Journal* 2012; **9**: 2534–2537.
4. **Dean AS, et al.** Clinical manifestations of human brucellosis: a systematic review and meta-analysis. *PLoS Neglected Tropical Diseases*. Published online: 6 December 2012. doi:10.1371/journal.pntd.0001929.
5. **Abdullayev R, et al.** Analyzing the spatial and temporal distribution of human brucellosis in Azerbaijan (1995–2009) using spatial and spatio-temporal statistics. *BMC Infectious Diseases*. Published online: 8 August 2012. doi: 10.1186/1471-2334-12-185.
6. **Al-Anazi KA, Al-Jasser AM.** Brucellosis: a global re-emerging zoonosis history, epidemiology, microbiology, immunology and genetics. Saudi Arabia: OMICS Group International 2013 (<http://www.esciencecentral.org/ebooks/bacterial-mycotic-infections/brucellosis1.php>). Accessed 13 March 2016.
7. **Pappas G, et al.** The new global map of human brucellosis. *Lancet Infectious Diseases* 2006; **6**: 91–99.
8. **Pakzad I, et al.** Human brucellosis in Iran: Incidence, complication, diagnosis, treatment and prevention. *Journal of Pure and Applied Microbiology* 2013; **7**: 1177–1188.
9. **Zeinali M, Shirzadi MR, Hajrasoliha H.** Iran's national guideline for brucellosis control. Centre of Communicable Diseases Management, Ministry of Health, *Treatment and Medical Education*, 2012, pp. 9–19.
10. **Hajia M, et al.** Comparison of methods for diagnosing brucellosis. *Laboratory Medicine* 2013; **44**: 29–33.
11. **Andriopoulos P, et al.** Prevalence of *Brucella* antibodies on a previously acute brucellosis infected population: sensitivity, specificity and predictive values of Rose Bengal and Wright standard tube agglutination tests. *Infection* 2015; **43**: 325–330.
12. **Goksugur SB, et al.** An interesting case of childhood brucellosis with unusual features. *Acta Clinica Croatica* 2015; **54**: 107–111.
13. **Maleki F, et al.** Epidemiological characteristics of human brucellosis in Shahin Dezh, Western Azarbaijan, Iran, 2008–2012. *Archives of Clinical Infectious Diseases* 2015; **10**: 1–4.
14. **Zeinalian Dastjerdi M, Fadaei Nobari R, Ramazanpour J.** Epidemiological features of human brucellosis in central Iran, 2006–2011. *Public Health*. Published online: December 2012. doi: 10.1016/j.puhe.2012.07.001.
15. **Leylabadlo HE, Bialvaei AZ, Samadi Kafil H.** Brucellosis in Iran: why not eradicated? *Clinical Infectious Diseases* 2015; **61**: 1629–1630.
16. **Dean AS, et al.** Global burden of human brucellosis: a systematic review of disease frequency. *PLoS Neglected Tropical Diseases*. Published online: 25 October 2012. doi:10.1371/journal.pntd.0001865.
17. **Ari MD, et al.** Challenges of establishing the correct diagnosis of outbreaks of acute febrile illnesses in Africa: the case of a likely *Brucella* outbreak among nomadic pastoralists, northeast Kenya, March–July 2005. *American Journal of Tropical Medicine and Hygiene* 2011; **85**: 909–912.
18. **Esmaeili H.** Brucellosis in Islamic republic of Iran. *Journal of Medical Bacteriology* 2014; **3**: 47–57.
19. **Honarvar B, et al.** Pulmonary tuberculosis in migratory nomadic populations: the missing link in Iran's National Tuberculosis Programme. *International Journal of Tuberculosis and Lung Disease* 2014; **18**: 272–276.
20. **Esmaeili S, et al.** Seroepidemiological survey of Q fever and brucellosis in Kurdistan Province, western Iran. *Vector-Borne and Zoonotic Diseases* 2014; **14**: 41–45.
21. **Alavi SM, Rafiei A, Nikkhooi A.** The effect of lifestyle on brucellosis among nomads in Khuzestan province of Iran. *Pakistan Journal of Medical Sciences* 2007; **23**: 358–360.
22. **Chegeni AS, et al.** Seroepidemiology of human brucellosis in nomads in a rural area of Iran. *Asian Pacific Journal of Tropical Disease* 2014; **4**: 333–336.
23. **Kansiime C, et al.** Community perceptions on integrating animal vaccination and health education by veterinary and public health workers in the prevention of brucellosis among pastoral communities of south western Uganda. *PLoS ONE*. Published online: 28 July 2015. doi:10.1371/journal.pone.0132206.
24. **Kansiime C, et al.** Knowledge and perceptions of brucellosis in the pastoral communities adjacent to Lake Mburo National Park, Uganda. *BMC Public Health* 2014; **14**: 242.
25. **Sofian M, et al.** Screening of family members of patients with acute brucellosis in an endemic area of Iran. *Iranian Journal of Microbiology* 2013; **5**: 215–219.
26. **Sanodze L, et al.** Expansion of brucellosis detection in the country of Georgia by screening household members of cases and neighboring community members. *BMC Public Health* 2015; **15**: 459.
27. **Tabak F, et al.** Is family screening necessary in brucellosis? *Infection* 2008; **36**: 575–577.
28. **Sadeghian H, et al.** Epidemiology and the agreement rate of serological tests in human brucellosis in North East of Iran. *Archives of Clinical Infectious Diseases* 2015; **10**: 2.
29. **Joulaei H, et al.** Introduction of a pyramidal model based on primary health care: a paradigm for management of 2009 H1N1 flu pandemic. *Iranian Red Crescent Medical Journal* 2010; **12**: 224–230.