

OSCILLATIONS OF THE MAGNETIC NETWORK

BRUCE W. LITES

High Altitude Observatory, NCAR, P.O. Box 3000, Boulder CO 80307-3000, USA

ROBERT J. RUTTEN

Sterrekundig Instituut, Postbus 80 000, NL-3508 TA Utrecht, The Netherlands

WOLFGANG KALKOFEN

Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge MA 02138, USA

ABSTRACT We present high-quality power spectra of oscillations in the quiet solar chromosphere measured from the Ca II H line. They show unambiguously that the network does not participate in chromospheric three-minute oscillations, but that instead it oscillates in slow wave motions which are neither present in the surrounding non-magnetic chromosphere, nor in the underlying photosphere.

1 INTRODUCTION

Chromospheric oscillations are of great interest in the context of cool-star chromospheric heating. The small-scale dynamics of the solar chromosphere supply the primary testing ground to constrain and identify heating mechanisms, whether magnetic or acoustic in origin (*cf.* Ulmschneider et al. 1991). Proper diagnostics are required, however; in particular, vertical discrimination is required between photospheric and chromospheric layers, and horizontal discrimination between the network and internetwork regimes.

Separation between network and internetwork has been made in oscillation analyses employing the Ca II infrared lines (Lites et al. 1982, Deubner & Fleck 1990) and H α (von Uexküll et al. 1989) but these lines offer limited height discrimination. The source functions of the Ca II infrared lines are controlled by H & K scattering (Uitenbroek 1989) and respond to changes over a wide height range (Mein 1971). H α has a doubly-peaked contribution function (Schoolman 1972).

Ca II H & K offer the cleanest chromospheric diagnostics accessible from the ground. The principal Fourier study of their oscillations is by Cram (1978). He did not distinguish between network and internetwork. We improve on his work by measuring oscillations in the core of Ca II H with such discrimination.




FIGURE I Spatially resolved chromospheric oscillations in Ca II H. Grey scales measure monochromatic intensity (top, bright is large) and Fourier power (other panels, dark is large). *Abcissae*: spatial dimension along spectrograph slit. *Top*: time-averaged intensity profiles. *Second panel*: power in line-center Doppler variations (V_3). *Third panel*: power in line-center intensity variations (I_3). *Bottom*: power in integrated-intensity variations (H-index; width $\Delta\lambda = 90$ pm).

The network areas crossed by the slit are marked by bright H_{2V} & H_{2R} emission in the top panel. Their line-center oscillation characteristics differ strikingly from the internetwork three-minute ($f = 3 - 10$ mHz) oscillations seen in V_3 . Both V_3 and I_3 show large low-frequency power ($f < 3$ mHz) in the network areas. The H-index panel contains bands of enhanced power for $f < 2$ mHz and $f = 3 - 7$ mHz, both without obvious network/internetwork distinction.

2 OBSERVATIONS

Our data consist of a one-hour sequence of Ca II H spectrograms taken at 5 s cadence with the NSO/SP Vacuum Tower Telescope in 1984. Figure 1 shows spectra that are spatially resolved along the spectrograph slit. The top panel displays time-averaged intensity spectra and shows bright network H_{2V} and H_{2R} features around $x = 30$ Mm and weaker emission near $x = 20$ Mm. These network locations stand out very clearly in the H_3 Dopplershift (V_3 , second panel) and intensity (I_3 , third panel) power spectra. The three-minute oscillation is prominently present *only* within the internetwork regions, and *only* in the V_3 panel. In stark contrast, the network areas crossed by the slit have much power in both V_3 and I_3 *only* at low frequencies, below $f = 5$ mHz. The vertical column at $x = 30$ Mm betrays seeing jitter of a small structure across the slit.

The bottom panel (H-index) shows intensity power in a $\Delta\lambda = 0.09$ nm wide band around line center. This index is representative of narrow-band Ca II K filtergrams and of the Mt. Wilson H & K photometer. The addition of inner-wing photons produces strikingly different characteristics compared with the I_3 panel. There are two bands of enhanced power which extend fully along the slit; the presence of network is *not* obvious in either band. Only the $x = 30$ Mm network column is weakly present.

The upper band in the H-index panel extends to higher frequencies than the acoustic oscillations of the photosphere (which cover the $f = 2.5 - 5$ mHz range). It contains the three-minute "whiskers" seen in the inner H & K wings in spatially resolved spectrograms (Beckers & Artzner 1974), which portray upwards-propagating acoustic oscillations (Rutten & Uitenbroek 1991). The low-frequency ($f < 2$ mHz) band of power was observed earlier by Schmieder (1976), Cram (1978) and Deubner & Fleck (1989) and attributed by these authors to gravity waves. It is not present in the Ca II H line-center power spectra (V_3 and I_3 panels).

3 DISCUSSION

Figure 1 shows that the network does not share in the internetwork three-minute velocity oscillation, but that instead it has low-frequency velocity and intensity power of its own which is not present in the internetwork, nor in the underlying photosphere (the latter follows from additional data, presented in a longer paper which we have submitted to ApJ).

Such low-frequency network power has been observed before, but far less cleanly and clearly than here. Damé et al. (1984) noted enhanced power for network areas in their 0.12 nm FWHM K-line filtergrams. Our bottom panel shows that such wide-band observations mix in too much lower-atmosphere signal from the Ca II wings to obtain a clear signature of the I_3 network power.

The Ca II infrared line analyses of Lites et al. (1982) and Deubner & Fleck (1990) suffer similarly from lower height of formation and from lack of height discrimination compared with the H & K cores. For example, all power spectra of Deubner & Fleck (1990) display prominent five-minute peaks and dips at 2–3 mHz that are seen here only in the H-index panel. This confirms that the Ca II infrared lines correspond primarily to the inner wings of H & K in formation

(Rutten & Uitenbroek 1991), with mixed-in chromospheric sensitivity through coupling to the H & K cores with NLTE-enhanced response (Uitenbroek 1989).

The H α observations of von Uexküll et al. (1989) come closest to these results, especially their Fig. 3 e–f in which the brightest, respectively darkest pixels were used to obtain network and internetwork velocity power spectra. However, their intensity power spectra contain a $f = 1$ mHz peak for both network and internetwork which corresponds to the low-frequency band in our H-index panel. It is absent in the I_3 and V_3 panels. Thus, the H α intensity seems sensitive to rather low-lying layers corresponding to H & K inner-wing formation.

What causes the low-frequency network power in V_3 and I_3 ? Von Uexküll et al. argue that it portrays stochastic motions. We believe from time-resolved displays and from new, similar data that these motions are oscillatory in nature. We suspect that these slow undulations represent a key phenomenon to identify the nature of chromospheric energy dissipation in magnetic elements. Obviously—to quote our Colloquium Chairman—we need more data.

4 REFERENCES

- Beckers, J. M., Artzner, G. 1974, *Solar Phys.*, 37, 309
- Cram, L. E. 1978, *A&A*, 70, 345
- Damé, L., Gouttebroze, P., Malherbe, J.-M. 1984, *A&A*, 130, 331
- Deubner, F.-L., Fleck, B. 1989, *A&A*, 213, 423
- Deubner, F.-L., Fleck, B. 1990, *A&A*, 228, 506
- Lites, B. W., Chipman, E. G., White, O. R. 1982, *ApJ*, 253, 367
- Mein, P. 1971, *Solar Phys.*, 20, 3
- Rutten, R. J., Uitenbroek, H. 1991, *Solar Phys.*, 134, 15
- Schmieder, B. 1976, *Solar Phys.*, 47, 435
- Schoolman, S. A. 1972, *Solar Phys.*, 22, 344
- Uitenbroek, H. 1989, *A&A*, 213, 360
- Ulmschneider, P., Priest, E. R., Rosner, R. (eds.) 1991, *Mechanisms of Chromospheric and Coronal Heating*, Proc. Int. Heidelberg Conference, Springer-Verlag, Berlin
- von Uexküll, M., Kneer, F., Malherbe, J. M., Mein, P. 1989, *A&A*, 208, 290