

INCREASED FREQUENCY OF SDB AND PLMS IS ASSOCIATED WITH LOWER REM-SLEEP AMOUNT IN COMMON CHILD PSYCHOPATHOLOGY AND NORMALLY DEVELOPING CHILDREN

R. Kirov¹, H. Uebel², B. Albrecht², L. Hecke², T. Banaschewski³, A. Rothenberger², *Sleep in Child Psychiatric Disorders*

¹Institute of Neurobiology, Bulgarian Academy of Sciences, Sofia, Bulgaria, ²Clinic for Child and Adolescent Psychiatry, University of Goettingen, Goettingen, ³Clinic for Child and Adolescent Psychiatry, Central Institute of Mental Health, Mannheim, Germany


Introduction: Sleep problems in children with common psychiatric disorders present a considerable challenge for clinicians in developing effective diagnosis and treatment strategies. Whilst sleep-disordered breathing (SDB) and periodic leg movements in sleep (PLMS) are very frequent in children with attention-deficit/hyperactivity disorder (ADHD) which can deviate sleep architecture, their co-existence in Tic disorder (TD) and ADHD/TD co-morbidity is less well understood.

Objectives: To investigate the frequency of SDB and PLMS across children with ADHD, TD and ADHD/TD co-morbidity compared with healthy peers.

Aims: We asked whether and how the frequency of SDB and PLMS relates to sleep architecture.

Methods: Twenty-four children with ADHD, 21 with TD, 21 with ADHD/TD co-morbidity and 22 healthy controls underwent a two-night polysomnography supplemented by monitoring of apnea-hypopnea (AH) and PLMS events per hour of total sleep time.

Results: Compared with controls, only ADHD children displayed a significantly higher AH and PLMS indices. Yet correlation analyses showed significant and negative association between AH and PLMS indices and rapid eye movement (REM) sleep amount in all, the ADHD, the TD (Fig. 1), the co-morbid, and the control (Fig. 2) groups. No such associations with the other sleep stages were found for all the groups.



[Figure 1]