

Using logistic regression analyses, the patient's fundamental attitude to longer-term medication and his satisfaction with the current medication on discharge were identified as prognostic factors. Relatives, friends, colleagues and the media were found to have no influence on the course of medication compliance. Nor did the compliance rate differ significantly between individual preparations or groups of preparations.

The results show that field studies with long-term follow-up periods are an important supplement to the generally short-term clinical studies when dealing with the question of medication compliance.

P45.05

Who says what? – Statements by schizophrenics and their psychiatrists on the current neuroleptic medication

G. Längle*, M. Köster, H. Waschulewski. *Tübingen University, Department of Psychiatry and Psychotherapy, Germany*

Cooperation based on mutual trust between patients and psychiatrists is a central precondition for successful neuroleptic relapse prevention in schizophrenic outpatients. On the one hand, scientific investigations into treatment compliance are often based on the assumption that the doctors giving the treatment cannot reliably estimate the compliance of their patients. On the other hand, statements made by patients with respect to their medication are seen as unreliable.

These two assumptions were subjected to verification within the framework of a follow-up field study of an unselected cohort of schizophrenic patients (N=169) six months after their discharge from hospital. Subject to the consent of the patients (N=81 / 66), the doctors treating them were interviewed in parallel (response rate 99%).

In the paired comparison, high and significant conformity ($\kappa=0.729$; $p<0.001$) was recorded between the statements made by the doctors and by the patients with respect to different groups of psychotherapeutic agents (e.g. anti-depressants, neuroleptics). The same applied to various groups of preparations (e.g. typical / atypical). With respect to statements on the dose of the preparation, the conformity rate between doctors and patients was 77%.

In contrast, there were major discrepancies in the statements made with respect to the acceptance of supplementary care offers and to re-hospitalisations. According to the present results within the framework of follow-up studies, reliable information on medication, but not on other aspects of psychosocial care, is to be obtained from both patients and doctors.

P45.06

Self-control and frontal lobe function in psychosis

J.-Å. Jansson¹*, B. Gervind², S. Warkentin³. ¹*Psykiatriska Rehabiliteringsenheten, Eslöv, Lund University hospital*; ²*Verksamhetsområde Psykiatri, Eslöv*; ³*Department of Clinical Physiology, MAS, Sweden*

Studies have shown that the experience of self-control in psychiatric patients is related to treatment outcome. In addition, numerous studies have reported frontal lobe dysfunction in groups of psychotic patients. It is however, still unknown to what extent a frontal lobe dysfunction might be related to a subjective experience of self-control in such patients.

Methods: The patient group consisted of nineteen subjects with schizoaffective psychosis or schizophrenia (mean age:35 (24–48 years)). Clinical ratings of self-control were made by Structural

Analysis of Social Behaviour (SASB). Brain function was assessed by regional cerebral blood flow.

Results: Patients who had relatively low level of subjective self-control (median cut) had relatively lower frontal lobe function, compared to those with higher self-control. The degree of self-control was also significantly correlated with level of perceived external support and involvement.

Conclusion: These observations suggest the possibility to identify subgroups of patients who differ in the degree to which they perceive external support in their rehabilitation process, and that such perceptual difference are related to self-control and brain function.

P45.07

Cannabis use and the expression of psychosis vulnerability in daily life

H. Verdoux*, C. Gindre, F. Sorbara, M. Tournier, J. Swendsen. *Department of Psychiatry, University of Bordeaux, France*

Objective: This study investigated in a non-clinical population the interaction between cannabis use and psychosis vulnerability in their effects on psychotic experiences in daily life.

Methods: Subjects (n=79) with high or low levels of cannabis use were selected among a sample of 685 undergraduate university students. Experience Sampling Method was used to collect information on substance use and psychotic experiences in daily life. Vulnerability to develop psychosis was measured using a clinical interview assessing the level of psychotic symptoms.

Results: The acute effects of cannabis are modified by the subject's level of vulnerability for psychosis. Subjects with high vulnerability for psychosis are more likely to report unusual perceptions as well as feelings of thought influence than subjects with low vulnerability for psychosis, and they are less likely to experience enhanced feelings of pleasure associated with cannabis. There is no evidence that use of cannabis is increased following occurrence of psychotic experiences as would be expected by the self-medication model.

Conclusion: Cannabis use interacts with psychosis vulnerability in their effects on experience of psychosis in daily life. The public health impact of the widespread use of cannabis may be considerable.

P45.08

Cannabis use and dimensions of psychosis in a non-clinical population of female subjects

H. Verdoux¹*, F. Sorbara¹, C. Gindre¹, J. Swendsen¹, J. van Os². ¹*Department of Psychiatry, University of Bordeaux, France* ²*Department of Psychiatry, University of Maastricht, The Netherlands*

Objective: The aim of the present study was to explore the pattern of associations between cannabis use and dimensions of psychosis in a non-clinical population of female subjects.

Method: The Community Assessment of Psychic Experiences (CAPE), a 42-item self-report questionnaire that evolved from the Peters et al. Delusions Inventory (PDI) was used to measure dimensions of psychosis in a sample of undergraduate female students (n=571). The participants were also asked to complete a self-report questionnaire collecting information on substance use.

Results: Three correlated dimensions of positive, negative and depressive experiences were identified using principal components factor analysis. Frequency of cannabis use was independently