

UPPSALA NATURAL RADIOCARBON MEASUREMENTS IX

INGRID U. OLSSON, SHAWKY EL-GAMMAL*,
and YETER GÖKSU**

Fysiska Institutionen, Uppsala Universitet, Uppsala, Sweden

The following list covers some old measurements not included in previous lists and most of the samples measured at the Uppsala C¹⁴ laboratory since the last list (Radiocarbon, 1967, v. 9, p. 454-470); samples utilized for determining the increase of the C¹⁴/C¹² ratio due to explosion of nuclear devices are omitted.

The technique used is mainly the same as previously described by Olsson (1958) with changes given in Radiocarbon, 1967, v. 9, p. 454. Pre-treatment of samples is mainly that which has been used earlier (Radiocarbon: 1964, v. 6, p. 291; 1967, v. 9, p. 454). Since shell fragments usually are different in size, the fraction of shell samples, given in per cent as a mean value, does not give the fraction of individual shells.

The reference sample is 95‰ of the C¹⁴/C¹² ratio of the NBS oxalic-acid standard. Any corrections for apparent water ages are thus not included here, but are discussed in papers dealing with the samples. Corrections for deviations from the normal C¹³/C¹² ratio (−25.0‰ in the PDB scale) are applied for the unknown samples. Our six oxalic-acid samples have not shown any significant difference in their C¹³/C¹² ratio. Our secondary standard, oxalic-acid I, was measured by Craig (1961) and it had a δC¹³ value of −18.97‰. This value agrees with the value to which the international oxalic-acid standard should be standardized. When δC¹³ is assumed, an error of 3‰ in the C¹³/C¹² ratio is used.

The value 5570 yr has been used for the half-life of C¹⁴, except in one case of geochemical interest (Table 1) where 5730 yr was used as stated in the head. Since the samples in Table 1 all are rather young, only one sample will get a slightly changed activity value if the results are calculated with the half-life 5570 yr.

Results in this list are given B.P. (before 1950) or as an excess, Δ, over the reference sample:

$$\Delta = \delta C^{14} - (2\delta C^{13} + 50) \left(1 + \frac{\delta C^{14}}{1000}\right)$$

where δC¹⁴ is the age-corrected C¹⁴ deviation from the reference sample in per mil in the year A.D. 1950, and δC¹³ is the deviation from PDB standard in per mil. Errors include standard deviation (σ) of counted particles and errors in corrections due to the C¹³/C¹² ratio, filling pressure, temperature, working voltage, barometric pressure etc. as described by Olsson (1965, 1966). When measured activity is lower than zero, 2 σ has been used for calculation of minimum age. When it is between zero and 2 σ, net activity is increased by 2 σ for calculation of minimum age. Since the results are presented as physical measurements, no terms are included for the error in the half-life or previous C¹⁴/C¹² variations.

* On leave from Univ. of Assuit, Assuit, Egypt.

** On leave from Middle East Technical University, Ankara, Turkey.

These errors are included in special papers dealing with the samples and the interpretation of the results.

Several samples had to be diluted with CO₂ from an old source to bring them to normal working pressure of the counters. This pressure has been about 2600 mm Hg for samples with numbers lower than U-1000, proportional counter 1, and about 1600 mm Hg for samples with numbers from U-2000, proportional counter 4.

ACKNOWLEDGMENTS

Descriptions of samples are based on information provided by those who collected and submitted them. Sincere thanks are due the contributors who helped in the preparation of the text. Special thanks are also due Dr. R. Ryhage and his co-workers for making the C¹³/C¹² determinations; Prof. K. Siegbahn, who has made it possible to do this work at the institute; and Statens Naturvetenskapliga Forskningsråd, which has given the laboratory financial support. The authors are indebted to Miss Ditte Ekwall, Miss Birgitta Wallin, Mr. Martin Klasson, Miss Carina Ericsson and Mrs. Maud Söderman for their work in the laboratory and to the following students who assisted: Torsten Jonsson, Gunnar Andersson, Thyra Erikson, Maj-Lis Persson, Maj-Liz Larsson, Anders Edin, Ulla Lindahl and Rolf Jonsson. One of us (S. E.-G.) acknowledges a fellowship from Sept. 1967 to Aug. 1968 and one (Y.G.) a fellowship until the end of Feb. 1967 from Swedish International Development Authority through the International Seminar for Research and Education in Physics.

SAMPLE DESCRIPTIONS

I. GEOLOGIC SAMPLES

A. Deep-sea cores

Indian Ocean series

Foraminifera tests from deep-sea core 156 (07° 14' N Lat, 52° 49' E Long), Indian Ocean, depth 5107 m. Coll. 1948 by Swedish *Albatross* Expedition (Pettersson); subm. by Eric Olausson, Inst. Oceanog. Univ. Gothenburg, Sweden. This core is rich in coccoliths except in a few horizons.

+5400

39,500

−3300

U-677. Core 156, 431.5 to 438.5 cm, > 62 μ , b

37,500 B.C.

$\delta C^{13} = -0.5\%$

Core 156, depth 431.5 to 438.5 cm, > 62 μ . Sample from coccolith-poor horizon. *Comment:* sample ultrasonically washed in boiled acidified distilled water, sieved, pH ca. 3, and dried before subm. to Uppsala. Activity of U-677 was slightly lower than that of U-678, but outer fraction was measured for a shorter time and statistical error was higher. As the 2 fractions had some activity within limit of errors, true age may be much higher than 40,000 yr (Olsson *et al.*, 1968). Innermost 60% was used.

U-678. Core 156, 431.5 to 438.5 cm, > 62 μ , a **33,500**
 $\delta C^{13} = +2.3\%$

Same sample as U-677; outermost 40% was used.

Red Sea series

Foraminifera tests from deep-sea core 162 (11° 57' N Lat, 44° 18' E Long), Red Sea, depth 883 m. Coll. 1948 by Swedish *Albatross* Expedition (Pettersson); subm. by Eric Olausson. Samples discussed by Olausson and Olsson (1969, Varve stratigraphy in a core from the Gulf of Aden: Palaeoecology, in press). For previous dates in this series, see Radiocarbon, 1967, v. 9, p. 455-456. *Comment*: difference between the 2 fractions is ascribed to contamination during long storage (Olsson *et al.*, 1968).

U-2061. Core 162, 306 to 319.5 cm, > 65 μ , b **4620 \pm 150**
2670 B.C.
 $\delta C^{13} = -3.5\%$

Core 162, depth 306 to 319.5 cm, > 65 μ . Sample from level where "warm" types (*Globigerinella aequilateralis*, *Globigerinoides rubra* and *G. sacculifera*) start (Todd, 1958). *Comment*: sample ultrasonically washed in boiled distilled acidified water, sieved, pH 3 to 4, and dried before subm. to Uppsala. Innermost 55% was used.

U-2062. Core 162, 306 to 319.5 cm, > 65 μ , a **3610 \pm 100**
1660 B.C.
 $\delta C^{13} = -0.7\%$

Same as U-2061; outermost 45% was used. Diluted.

Pacific Ocean core series

Foraminifera from core MSN 138 G (00° 15' S Lat, 147° 34' W Long). Coll. 1961 by the Monsoon Expedition. Samples were separated in different size fractions by sieving by boiled acidified distilled water (pH4) and leached in HCl to give different fractions for studies of contamination such as was seen in cores from Mediterranean (Eriksson and Olsson, 1963; Olsson and Eriksson, 1965; Olsson *et al.*, 1968). Pretreated and subm. by K. Gösta Eriksson, Inst. Geol., Univ. Gothenburg, Sweden.

U-2114. MSN 138 G, 0 to 5 cm, > 62 μ , b **4450 \pm 160**
2500 B.C.
 $\delta C^{13} = +1.7\%$

Core MSN 138 G, depth 0 to 5 cm, > 62 μ . *Comment*: innermost 50% was used. Diluted.

U-681. MSN 138 G, 0 to 5 cm, 4 to 45 μ , b **4540 \pm 70**
2590 B.C.
 $\delta C^{13} = -1.4\%$

Core MSN 138 G, depth 0 to 5 cm, 4 to 45 μ . *Comment*: innermost 40% was used.

U-683. MSN 138 G, 0 to 5 cm, 4 to 45 μ , a **4550 \pm 100**
2600 B.C.
 $\delta C^{13} = +2.6\%$

Core MSN 138 G, depth 0 to 5 cm, 4 to 45 μ . *Comment*: outermost 60% was used.

		+1500
		6900
		-1300
U-2096.	MSN 138 G, 5 to 10 cm, > 62μ, b	4900 B.C.
		$\delta C^{13} = -5.8\text{‰}$
	Core MSN 138 G, depth 5 to 10 cm, > 62 μ . <i>Comment:</i> innermost 5% was used. Diluted.	
		4430 \pm 130
U-2097.	MSN 138 G, 5 to 10 cm, > 62μ, a	2480 B.C.
		$\delta C^{13} = -0.3\text{‰}$
	Core MSN 138 G, depth 5 to 10 cm, > 62 μ . <i>Comment:</i> outermost 95% was used. Diluted.	
		5480 \pm 100
U-2099.	MSN 138 G, 5 to 10 cm, 4 to 45μ, b	3530 B.C.
		$\delta C^{13} = -2.1\text{‰}$
	Core MSN 138 G, depth 5 to 10 cm, 4 to 45 μ . <i>Comment:</i> innermost 30% was used, but fractionation might have failed.	
		5230 \pm 60
U-679.	MSN 138 G, 5 to 10 cm, 4 to 45μ, a	3280 B.C.
		$\delta C^{13} = -7.6\text{‰}$
	Core MSN 138 G, depth 5 to 10 cm, 4 to 45 μ . <i>Comment:</i> outermost 70% was used.	
		9580 \pm 270
U-2100.	MSN 138 G, 15 to 20 cm, > 62μ	7630 B.C.
		δC^{13} assumed +1.6‰
	Core MSN 138 G, depth 15 to 20 cm, > 62 μ . <i>Comment:</i> whole sample was used. Diluted.	
		9730 \pm 310
U-2098.	MSN 138 G, 15 to 20 cm, 4 to 45μ, b	7780 B.C.
		$\delta C^{13} = -3.4\text{‰}$
	Core MSN 138 G, depth 15 to 20 cm, 4 to 45 μ . <i>Comment:</i> innermost 25% was used. Diluted.	
		9750 \pm 100
U-680.	MSN 138 G, 15 to 20 cm, 4 to 45μ, a	7800 B.C.
		$\delta C^{13} = -3.5\text{‰}$
	Core MSN 138 G, depth 15 to 20 cm, 4 to 45 μ . <i>Comment:</i> outermost 75% was used.	

General Comment: only one sample, U-2097, shows significant deviation from other fractions of same level in core.

B. Asia

Sogho-nor series

Plant remains from Lake Sogho-nor (42° N Lat, 101° E Long). Samples coll. by Hörner; subm. by E. Norin, Inst. Geol., Univ. Uppsala, Sweden. Samples from strand-line now 8.6 m above present level. *Comment:* oasis of Etsina was flourishing ca. 600 yr ago; results show that water

level then was ca. 8 m higher than now and lake much larger. Described by Norin (1966).

U-283. Sogho-nor 270530 **500 ± 300**
A.D. 1450
 $\delta C^{13} = -16.0\%$

Aquatic plants, mainly, coll. in highest ancient beach ridge, 7.5 m above water level. Sample SW of lake. Coll. 1930. *Comment:* diluted.

U-284. Sogho-nor 150333:1 **670 ± 110**
A.D. 1280
 $\delta C^{13} = -14.7\%$

Aquatic plants, mainly, coll. from ancient beach, probably 8 m above level of lake. Sample SE of lake. Coll. 1933. *Comment:* diluted.

U-273. Sogho-nor 150333:2 **660 ± 120**
A.D. 1290
 $\delta C^{13} = -15.7\%$

Same sample and pretreatment as U-284, but new combustion. *Comment:* diluted.

C. Central Europe

U-2036. Elmau **>34,600**
 $\delta C^{13} = -25.6\%$

Tree root from adit Elmau (ca. 47° 25' N Lat, 13° 05' E Long), Mühlbach/Hochkönig, Salzburg, Austria. Trees were growing on slope of former valley, now filled by gravel and breccia ca. 80 cm thick. Coll. by M. Maczek, Kupferbergbau Mitterberg, Austria.

D. Lithuanian S.S.R.

River Ula basin series

Wood from Rudnja (54° N Lat, 25° E Long), Ula R. basin, 70 km S Vilnius, Lithuania. Material from this area has been dated previously, Mo-302, Zervinos (moss): 16,260 ± 640; Mo-339, Rudnja (peat): 12,715 ± 315; Mo-341, Pamärkis (peat): 11,500 ± 430; Mo-340, Pamärkis (wood): 12,260 ± 160; Vs-4, Zervinos (peat): 18,350 ± 950; Vs-5, Manciangire (peat): 17,340 ± 840; TA-124, Zervinos (wood): 11,930 ± 110; TA-125, Zervinos (moss): 12,160 ± 120; TA-191, Zervinos (moss): 12,650 ± 130; TA-190, Rudnja (wood): 11,530 ± 120 (Simm, pers. commun.). Samples are deposits submerged under fluvioglacial and eolian sand; they are suggested either to be interstadial or of late-glacial origin. Wood of same tree trunk used for TA-190, subm. as interlaboratory check by H. Simm, Zool. Bot. Inst., Eesti NSV Teaduste Akad., Tartu, Estonia S.S.R.

U-675. Rudnja, 2 **11,970 ± 180**
10,020 B.C.
 δC^{13} assumed -24.9%

Wood from Rudnja, 0.39 to 0.93 m below surface. Sample coll. in aleuritic sand with peat and wood.

+460

12,080

-430

U-2107. Rudnja, 1**10,130 B.C.** δC^{13} assumed -25.0‰

Wood from same sample as was used for U-675, but new pre-treatment.

*E. Iceland***Shoreline series**

Shell and wood from Iceland to determine chronology of Late- and Post-glacial shorelines. Coll. 1966 and subm. by Thorleifur Einarsson, Univ. Research Inst., Reykjavik, Iceland.

12,290 ± 160**U-641. Melar, 3-67, b****10,340 B.C.** $\delta C^{13} = +0.2‰$

One valve (*Pecten islandicus*) from Melar (64° 25' N Long, 22° 01' W Long), Melasveit, Iceland, alt 5 m (Bárdarson, 1923). Sample from glaciomarine silt, which seems to interfinger with an end moraine of Alltanes Readvance, which probably is of Older Dryas age. *Comment*: similar sample, U-2019, dated: 11,620 ± 240 B.P. (Radiocarbon, 1967, v. 9, p. 458, where preliminary result was given). Innermost 60% was used.

12,610 ± 200**U-2054. Melar, 3-67, a****10,660 B.C.** $\delta C^{13} = +0.2‰$

Shell layer surrounding part used for U-641. *Comment*: outermost 40% was used.

6940 ± 110**U-640. Hólahólar, 5-67****4990 B.C.** $\delta C^{13} = -24.0‰$

Juniperus communis from Hólahólar (65° 20' N Lat, 18° 10' W Long), Eyjafjörður, Iceland, alt 60 m. Sample from rockslide, cut through by the Eyjafjardará R. Sample supposed to be older than Hekla ash layer H₅. Pollen Zone A or B from N Iceland (Einarsson, 1961).

3140 ± 140**U-2088. Baejará, 2-67****1190 B.C.** $\delta C^{13} = 0.0‰$

Astarte borealis (L.) from Baejará (65° 18' N Lat, 21° 11' W Long), Hrótafjörður, Iceland, alt 3 m. Sample from top of gravel bar, 2.5 m higher than present one, 35 m inland and covered by ash layer H₄, assumed 4000 B.P. (Bárdarson, 1910; Thorarinsson, 1955). *Comment*: innermost 90% was used. Diluted.

1140 ± 70**U-2082. Tjarnargata 4, 6-67****A.D. 810** $\delta C^{13} = -26.8‰$

Wood shavings from Tjarnargata 4, Reykjavik (64° 09' N Lat, 21° 57' W Long), Iceland. Sample in peat with wood fragments and bones (*Sus scrofa*, domesticus, *Alca impennis*, and *Trichechis rosmarus*). Coll.

1944 by Finnur Gudmundsson, Mus. Nat. Hist., Reykjavik, Iceland and subm. by Th. Einarsson. *Comment*: another sample from old cultural layer nearby was dated previously: K-940, 1340 ± 100 (Radiocarbon, 1966, v. 8, p. 232). Diluted.

U-632. Ellidaár 650609 **4630 ± 90**
2680 B.C.

$$\delta C^{13} = -26.6\text{‰}$$

Charred pieces of birch from Ellidaár (64° 08' N Lat, 21° 50' W Long), Reykjavik, Iceland. Sample underlying lava, Leitahraun, and overlying *Parvocaricetum* peat. Coll. 1965 and subm. by Jón Jónsson, State Electricity Authority, Reykjavik, Iceland. Lava described by Hospers (1953). *Comment*: peat nearby was dated previously: C-749, 5300 ± 340 years (Libby, 1955).

U-2051. Dýrholaós **1240 ± 100**
A.D. 710

$$\delta C^{13} = -23.7\text{‰}$$

Driftwood from Dýrholaós (63° 20' N Lat, 19° 05' W Long), Skaftafellssýsla, Iceland. Sample from a log, 3 m long, diam. 25 to 30 cm, imbedded in peat, now at a lagoon. Due to scarcity of timber on Iceland, logs at this locality presumed to have been imbedded before colonization. Thickness of peat also implies that sea level of the time was several m lower than now. Driftwood deposited at extreme highwater. Coll. and subm. by Jón Jónsson.

F. Spitsbergen, Björnöya, and Shetland

Vestspitsbergen series

Driftwood, shells, and whale bones from Vestspitsbergen coll. for determination of land uplift. Alt. given above mean sea level. Previous determinations in this series are given in Radiocarbon: 1960, v. 2, p. 115-116; 1961, v. 3, p. 82-83; 1964, v. 6, p. 296-298; 1965, v. 7, p. 317-318; 1967, v. 9, p. 456-457. Quaternary geology and land uplift is discussed by Birkenmajer (1960), Feyling-Hanssen (1955), Feyling-Hanssen and Olsson (1959-1960). Samples from Hornsund area coll. 1966 and subm. by Krzysztof Birkenmajer, Polish Acad. Sci., Krakow, Poland and those from Isfjorden area coll. 1960 by D. H. Maling. *Comment*: bone treatment with EDTA is described in Radiocarbon, 1967, v. 9, p. 456.

U-619. Rettkvalbogen E, 1 st, Q4a **1080 ± 70**
A.D. 870

$$\delta C^{13} = -23.7\text{‰}$$

Driftwood from Rettkvalbogen E (77° 00' N Lat, 15° 32' 30" E Long), Hornsund, Spitsbergen, alt. 5.5 m. Sample from log, 10 m long, 40 cm diam., partly buried in moss, 1st terrace.

U-2048. Rettkvalbogen E, 1 st, Q5a, R **1120 ± 80**
A.D. 830

$$\delta C^{13} = -19.5\text{‰}$$

Balaena mysticetus L. from Rettkvalbogen E (77° 00' N Lat, 15°

32' 30" E Long), Hornsund, Spitsbergen, alt 5.5 m. Sample from almost complete upper left jaw, 5.6 m long, 0.45 m wide in its rear edge, partly buried in moss, 1st terrace. *Comment*: sample treated with acetone before EDTA extraction.

U-620. Rettkvalbogen E, 1 st, Q5a, A **770 ± 70**
A.D. 1180
 $\delta C^{13} = -24.3\%$

Acetone extract from sample used for U-2048.

U-2079. Fuglebekken, 2a₁, Q6a, b **9210 ± 180**
7260 B.C.
 δC^{13} assumed -2.1%

Fragments (*Mya truncata* and *Astarte borealis*) from Fuglebekken (77° 00' 30" N Lat, 15° 34' E Long), Hornsund, Spitsbergen, alt 5 m. Sample in sandy gravel, Horizon "a" of 2a₁ Mya Terrace. *Comment*: innermost 33% was used.

U-2069. Fuglebekken, 2a₁, Q6a, a **9150 ± 160**
7200 B.C.
 $\delta C^{13} = -2.1\%$

Shell layer surrounding part used for U-2079. *Comment*: layer corresponds to 37% of shells; 30% we removed by washing.

U-682. Marmorneset, 2a₂, Q1a, b **7410 ± 90**
5460 B.C.
 $\delta C^{13} = +0.3\%$

Shell fragments, mostly *Balanus*, from W part of Marmorneset (77° 00' N Lat, 15° 29' E Long), Hornsund, Spitsbergen, alt 8 m. Sample in shingle and subrounded gravel, with sandy matrix, Horizon "f" of 2a₂ Saxicava- Mya Terrace. *Comment*: innermost 55% was used.

U-2065. Marmorneset, 2a₂, Q1a, a **7620 ± 130**
5670 B.C.
 $\delta C^{13} = +0.2\%$

Shell layer surrounding part used for U-648. *Comment*: layer corresponds to 20% of shells; 25% was removed by washing.

U-665. Isbjørnhamna, 2a₂, Q3a, b **9620 ± 110**
7670 B.C.
 $\delta C^{13} = +0.2\%$

Fragments (*Mya truncata* and *Saxicava arctica*) from Isbjørnhamna (77° 00' N Lat, 15° 33' 30" E Long), Wilczekodden, Hornsund, Spitsbergen, alt. 7.5 m. Sample overlay sand and gravel but underlay sand, Horizon "d" of 2a₂ Terrace, Saxicava-Mya Terrace. *Comment*: innermost 50% was used.

U-666. Isbjørnhamna, 2a₂, Q3a, a **9560 ± 140**
7610 B.C.
 δC^{13} assumed $+0.2\%$

Shell layer surrounding part used for U-665. *Comment*: layer corresponds to 30% of shells; 20% was removed by washing.

U-703. Isbjørnhamna, 2a₂, Q2a, R1 **9380 ± 140**
7430 B.C.
 $\delta C^{13} = -17.1\%$

Whale bone (probably *Balaena mysticetus* L.) from Isbjørnhamna (77° 00' N Lat, 15° 33' 30" E Long), 82 m SW Polish Sta., Hornsund, Spitsbergen, alt 8 m. Sample from lower jaw, 1.9 m long, 17 cm diam, 2/3 buried in tundra: mosses, lichens, *Saxifraga*, *Salix polaris*, and grass, 2a₂ Terrace. *Comment*: EDTA-treated after acetone extraction.

U-2130. Isbjørnhamna, 2a₂, Q2a, R2 **9840 ± 230**
7890 B.C.
 $\delta C^{13} = -18.2\%$

Gas from another combustion but from same pretreatment as for sample U-703. Diluted.

U-2131. Isbjørnhamna, 2a₂, Q2a, W **8610 ± 170**
660 B.C.
 $\delta C^{13} = -19.4\%$

Wrong fraction from EDTA-treatment of sample U-703 and U-2130.

U-2077. Ekholmrika 6020 b, R **9650 ± 130**
7700 B.C.
 $\delta C^{13} = -19.4\%$

Whale rib from Ekholmrika (78° 35' N Lat, 16° 38' E Long), Billefjorden, Spitsbergen, alt ca. 50 m. Sample protruding from surface and buried 1 m in coarse gravel. Location mapped by Balchin (1941). *Comment*: sample treated with acetone before EDTA-treatment.

U-2066. Ekholmrika 6020 b, inorganic $\Delta = 1270 \pm 36\%$
 δC^{13} assumed -5.2%

Same rib as U-2077 but inorganic fraction, CO₂ liberated at HCl-treatment under vacuum, assumed to give false result. *Comment*: diluted. Sample contaminated by CO₂ of air in lab.

U-2076. Ekholmrika 6020 b, HCl-NaOH **+440**
3500
-410
1550 B.C.
 $\delta C^{13} = -24.8\%$

Insoluble remains from HCl treatment giving U-2066. Remains treated with 0.5% NaOH, washed, made acid and dried before combustion. *Comment*: diluted.

U-2121. Ekholmrika 6020 b, HCl **5990 ± 380**
4040 B.C.
 δC^{13} assumed -19.0%

Insoluble remains of same rib as U-2066, 2076, and 2077, after treatment of crushed bones with 1-N HCl until pH1 and washing with distilled water before drying. *Comment*: diluted.

U-2106. Ekholmrika 6021 b, R **9390 ± 180**
7440 B.C.
 $\delta C^{13} = -18.4\text{‰}$

Whale rib from Ekholmrika (78° 35' N Lat, 16° 38' E Long), Billefjorden, Spitsbergen, alt ca. 50 m. Sample protruding ca. 0.75 m from surface and easily broken.

Nordautlandet series

Wood and shells for determination of land uplift. Previous samples in this series are given by Olsson and Blake (1961-1962) and in Radiocarbon, 1959, v. 1, p. 90-91; 1960, v. 2, p. 116-121; 1964, v. 6, p. 298-300. Coll. and subm. by W. Blake, Jr., Geol. Surv. of Canada, Ottawa, Ontario, Canada, except for K. Sparre-Sveanor samples.

U-618. Kvalrosshalvøya 174-66 **6700 ± 70**
4750 B.C.
 $\delta C^{13} = -23.7\text{‰}$

Picea or *Larix* from Kvalrosshalvøya (79° 59' N Lat, 18° 35' E Long), Nordautlandet, Spitsbergen, alt 9.8 m. Sample partly buried in beach shingle with pumice. Subm. to check suggestion (Olsson and Blake, 1961-1962) that U-33 and U-34 dated 9.8 and 2.0 m levels, respectively, instead of the contrary. U-618 is from same log as U-33 (6910 ± 110). Coll. 1966.

U-2058. Lindhagenbukta 71-66, b **410 ± 90**
A.D. 1540
 $\delta C^{13} = -0.6\text{‰}$

Macoma calcarea from Lindhagenbukta (80° 17' N Lat, 20° 42' E Long), Nordautlandet, Spitsbergen, alt 0 to 2 m. Sample from assemblage of well-preserved shells, still hinged and with periostracum nearly intact. Sample supposed to give apparent age of water; shells assumed to be only a few yr old as they had not been crushed by rafted ice. Coll. 1966. *Comment*: apparent age of water 420 yr, $\Delta = -50 \pm 10\text{‰}$, if sample assumed from 1960. Innermost 40% was used.

U-2059. Lindhagenbukta 71-66, a **380 ± 90**
A.D. 1570
 $\delta C^{13} = -1.6\text{‰}$

Shell layer surrounding U-2058. *Comment*: apparent age 390 yr; $\Delta = -47 \pm 10\text{‰}$. Outermost 60% of shells was used.

U-646. Lady Franklinfjorden 163-66, b **22,600 ± 500**
20,600 B.C.
 $\delta C^{13} = +5.3\text{‰}$

Fragments (*Hiatella arctica*, *Mya truncata*, and possibly *Macoma* and *Chlamys*) from Kapp Lady (80° 12' N Lat, 18° 42' E Long), Lady Franklinfjorden, Nordautlandet, Spitsbergen, alt ca. 2 m; depth ca. 0.5 m, from till overlain by beach shingle with lenses of organic debris between till and gravel. Surface of shells pitted but sample dated to check

U-263 (Radiocarbon, 1964, v. 6, p. 299) from same place. Both dates thought by senior author to give unrealistically low ages. Coll. 1966. *Comment*: innermost 60% was used.

+900
21,700
-800

U-2060. Lady Franklinfjorden, 163-66, a 19,700 B.C.
 $\delta C^{13} = +5.9\%$

Shell layer surrounding U-646. *Comment*: layer corresponds to 30% of sample. 10% was removed by washing.

U-2095. K. Sparre-Sveanor, c 11,180 ± 190
9230 B.C.
 $\delta C^{13} = +0.3\%$

Mya truncata from K. Sparre-Sveanor (79° 56' N Lat, 18° 15' E Long), Nordaustlandet, Spitsbergen, alt 82 m. Shells partly buried in beach material. Coll. 1966 by M. Grosswald; subm. by G. Hoppe. *Comment*: shells from same collection previously dated, St-2374: 10,670 ± 145. Innermost 35% was used.

U-660. K. Sparre-Sveanor, b 11,150 ± 110
9200 B.C.
 δC^{13} assumed +0.3%

Shell layer surrounding U-2095. *Comment*: layer corresponds to 40% of shells.

U-661. K. Sparre-Sveanor, a 10,370 ± 220
8420 B.C.
 δC^{13} assumed +0.3%

Shell layer surrounding U-2095 and U-660. *Comment*: layer corresponds to 25% of shells.

Lake sediment series; Nordaustlandet, Bjørnøya, and Shetland

Silty mud from Skinkevatnet (74° 30' N Lat, 19° 00' E Long), Bjørnøya, alt 19.3 m; clay-gyttja from Rippfjorden (80° 18' N Lat, 22° 40' E Long), Nordaustlandet, Spitsbergen, alt 30 m; gyttja from tidal lake Lower Loch of Brouster (60° 15' N Lat, 01° 36' W Long), Shetland; graphite from chemical store. Geologic samples supposedly contain old carbon and different separation methods are being tried. Bjørnøya samples coll. 1965, subm. and described by Hannu Hyvärinen (1968); Nordaustlandet sample coll. 1965 and subm. by Hyvärinen, Dept. Geol. and Palaeontol. Univ. Helsinki, Finland; Shetland sample coll. 1964, subm. and described by Gunnar Hoppe (1965), Dept. Physical Geog. Univ. Stockholm, Sweden; chemical treatment is discussed by Olsson in a preliminary report (1968). *Comment*: all samples diluted—in many cases more than can be recommended, so that errors are so large that result is of little geologic interest.

	11,200 ± 500
U-2031. Skinkevatnet, 92.5 to 87.5, H	9200 B.C.
	$\delta C^{13} = -25.4\%$
Humus products extracted from sample, 92.5 to 87.5 cm depth.	+1400
	8300
	-1200
U-2042. Skinkevatnet, 92.5 to 87.5, HNO₃+H₂O₂	6300 B.C.
	$\delta C^{13} = -25.3\%$
Gas obtained by wet combustion of sample, with HNO ₃ + H ₂ O ₂ , after alkali extraction that gave U-2031.	+1200
	8900
	-1000
U-2064. Skinkevatnet, 92.5 to 87.5, INS	7000 B.C.
	δC^{13} assumed -25.4%
Insoluble remains of sample, after wet combustion that gave U-2042.	+1200
	4600
	-1000
U-2050. Skinkevatnet, 42.5 to 37.5, H	2700 B.C.
	$\delta C^{13} = -28.2\%$
Humus products extracted from sample, 42.5 to 37.5 cm depth.	+1100
	7300
	-1000
U-2125. Skinkevatnet, 42.5 to 37.5, KMnO₄	5300 B.C.
	$\delta C^{13} = -22.6\%$
Gas obtained by wet combustion of sample, with H ₂ SO ₄ +KMnO ₄ , after extraction that gave U-2050.	9460 ± 350
U-2128. Skinkevatnet, 42.5 to 37.5, INS	7510 B.C.
	$\delta C^{13} = -21.7\%$
Insoluble remains of sample, after wet combustion that gave U-2125.	+3100
	6000
	-2200
U-2049. Skinkevatnet, 22.5 to 17.5, H	4100 B.C.
	$\delta C^{13} = -23.9\%$
Humus products extracted from sample, 22.5 to 17.5 cm depth.	6390 ± 270
U-2124. Skinkevatnet, 22.5 to 17.5, KMnO₄	4420 B.C.
	$\delta C^{13} = -21.9\%$
Gas obtained by wet combustion of sample, with H ₂ SO ₄ +KMnO ₄ , after extraction that gave U-2049.	

U-2127. Skinkevatnet, 22.5 to 17.5, INS **6870 ± 470**
4920 B.C.

$$\delta C^{13} = -21.7\text{‰}$$

Insoluble remains of sample, after wet combustion that gave U-2124.

U-2110. Rippfjorden 3/23, 60 to 55, H **10,150 ± 450**
8200 B.C.

$$\delta C^{13} = -27.9\text{‰}$$

Humus products extracted from sample, 60.5 to 54.5 cm depth.
Comment: another sample immediately underlying that used for U-2110 was dated previously, St-2532: 10,885 ± 250.

+550

9640

-510

U-2126. Rippfjorden 3/23, 60 to 55, KMnO₄ **7690 B.C.**

$$\delta C^{13} = -28.8\text{‰}$$

Gas obtained by wet combustion of sample, with H₂SO₄+KMnO₄, after extraction that gave U-2110.

U-2129. Rippfjorden 3/23, 60 to 55, INS **>20,500**

$$\delta C^{13} = -25.3\text{‰}$$

Insoluble remains of sample, after wet combustion that gave U-2126.

+840

6300

-720

U-616. Lower Loch of Brouster, Core 20,
343 to 338, H, I **4350 B.C.**

$$\delta C^{13} = -20.0\text{‰}$$

Humus extracted from gyttja, 343 to 338 cm below top of sediment.
Comment: another sample from same core, 5 cm lower, was dated previously (U-576 and U-2007, mean 7650 ± 450; Radiocarbon, 1967, v. 9, p. 460).

+4300

4500

-2800

U-2041. Lower Loch of Brouster, Core 20,
343 to 338, HNO₃+H₂O₂ **2600 B.C.**

$$\delta C^{13} \text{ assumed } -25.0\text{‰}$$

Gas obtained by wet combustion of sample, with HNO₃+H₂O₂, after extraction that gave U-616.

+1300

5800

-1100

U-2067. Lower Loch of Brouster, Core 20,
343 to 338, INS **3900 B.C.:**

$$\delta C^{13} \text{ assumed } -24.9\text{‰}$$

Insoluble remains of sample, after wet combustion that gave U-2041.

U-2123. Graphite 060868, KMnO₄ **>26,400**
 $\delta C^{13} = -23.2\text{‰}$

Graphite from dealer in chemicals, Kebo. *Comment*: gas obtained by wet combustion with H₂SO₄ + KMnO₄.

G. Norway

U-2121. Sandnes 1966, No. 11, R **>27,300**
 $\delta C^{13} = -13.9\text{‰}$

Balaena mysticetus from Sandnes (58° 50' N Lat, 05° 42' E Long), Rogaland, Norway, 10 m deep in glaciomarine clay with *Portlandia arctica* etc., at ca. 17 m depth. Clay overlain by sand and Würm till. Coll. 1966 and subm. by Björn Andersen, Inst. Geol. Univ. Oslo, Blindern, Norway. *Comment*: sample EDTA-treated. Another piece of same bone dated previously after normal HCl treatment: T-641, 28,900 ± 800 (Andersen, written commun.). Diluted.

Skjomen series

Gyttja from 2 tarns at defile at Somarskardet (68° 17' N Lat, 16° 73' E Long), near fault scarp, SW Narvik, Nordland, Norway, alt ca. +600 m. Samples coll. to correlate lateral moraines in area with position of Tromsö-Lyngen substage ice margin. Geology discussed by Dahl (1967, 1968). Coll. 1966 and subm. by Ragnar Dahl, Dept. Physical Geog., Univ. Uppsala, Sweden. *Comment*: differences between humus-products and insoluble remains are, for 2 of the samples, rather large although samples were so small that they had to be diluted; errors, thus, are large. Dilution for U-605 and U-2028 was larger than normally allowed. Taking previous variations of C¹⁴/C¹² into account, large differences will decrease, however.

U-605. Skjomen I:1, 0 to 5, INS **5200 ± 700**
3200 B.C.
 $\delta C^{13} = -28.1\text{‰}$

Gyttja from lower 5 cm of 80 cm thick organic sediments of 1 tarn, 250 m N of threshold, E of and near moraine ridge.

U-2028. Skjomen I:1, 0 to 5, H **+490**
7830
-450
5880 B.C.
 $\delta C^{13} = -27.2\text{‰}$

Humus products extracted from sample used for U-605.

U-2037. Skjomen I:2, 5 to 10, INS **3650 ± 110**
1700 B.C.
 $\delta C^{13} = -22.9\text{‰}$

Gyttja 5 to 10 cm above sand and silt bottom of tarn from which U-605 + U-2028 was coll.

+700

2620

-630

U-2038. Skjomen I:2, 5 to 10, H

670 B.C.

 $\delta C^{13} = -28.7\%$

Humus products extracted from sample used for U-2037.

3210 \pm 100**U-2039. Skjomen I:3, 0 to 5, INS**

1260 B.C.

 $\delta C^{13} = -25.8\%$

Gyttja from lower 5 cm of 30 cm thick organic sediments of tarn, W of moraine ridge near tarn where Sample U-605 was coll.

3390 \pm 230**U-2040. Skjomen I:3, 0 to 5, H**

1440 B.C.

 $\delta C^{13} = -27.8\%$

Humus products extracted from sample used for U-2039.

*H. Sweden*8920 \pm 190**U-2071. Levide, TF 67**

6970 B.C.

 $\delta C^{13} = -25.0\%$ Wood from trunk (*Pinus*) from Levide (57° 15' N Lat, 18° 15' E Long), Gotland, Sweden. Sample from sand and gravel, 2 m below surface, with *Ancylus* shells and several other trunks. *Comment*: 12 outer tree rings of log, containing 50 tree rings, were used. Coll. 1967 and subm. by Trygve Fahlstedt, Dept. Agr. Hydrotechnics, High School of Agric. Ultuna, Uppsala, Sweden. *Comment*: another sample from Levide was dated previously, U-428: 9830 \pm 140 (Radiocarbon, 1964, v. 6, p. 301).**Shell bank series**

Shells from Lindalsskogen, Väjern (58° 22' N Lat, 06° 47' E Long), Bohuslän, Sweden, alt 25 m and at one unspecified place in Bohuslän, Sweden; shells from Väjern coll. 1966 in shell banks subm. by K. Gösta Eriksson, Inst. Geol. Univ. Gothenburg, Sweden. Samples discussed by Eriksson and Olsson (1967).

10,160 \pm 330**U-2029. Väjern 3282 B**

8210 B.C.

 δC^{13} assumed -1.0% Shells (*Balanus balanus*) coll. 0.2 to 1 m below erosion layer at alt. ca. 24 m. *Comment*: innermost 30% was used. Diluted.10,230 \pm 240**U-2030. Väjern 3282 H**

8280 B.C.

 $\delta C^{13} = +1.4\%$ Shells (*Balanus hameri*) 0.2 to 1 m below erosion layer. *Comment*: innermost 55% was used. Diluted.

U-607. Bohuslän **530 ± 80**
A.D. 1420
 $\delta C^{13} = -1.7\text{‰}$

Valve (*Patella vulgata*) from Bohuslän coll. ca. 1935 by Ingrid U. Olsson. Apparent age: 510 yr. $\Delta = -62 \pm 9\text{‰}$. *Comment:* innermost 95% was used. Diluted.

Degerfors series

Gyttja from Svea Älv (Svea R.) area at Degerfors. Coll. 1967 and subm. by Curt Fredén, Inst. Quaternary Geol., Univ. Uppsala, Sweden. Coll. with Livingstone borer. Pollen analyses by Fredén. Survey of previous discussions on Svea R. given by Fredén (1967). Samples described by Olsson *et al.* (in press).

U-2080. Håkanbol 7, INS **8060 ± 220**
6110 B.C.
 $\delta C^{13} = -33.5\text{‰}$

Clay-gyttja with carbon fragments from Håkanbol (59° 10' N Lat, 14° 22' E Long), Degerfors, Närke, Sweden, at depth 1850 cm below surface in kettle hole filled by peat. Zone V (Jessen). Before birch culmination. Assumed older than Svea R. *Comment:* diluted.

U-2081. Håkanbol 7, H **+640**
8860
-600
6910 B.C.
 δC^{13} assumed -25‰

Humus products extracted from sample used for U-2080. *Comment:* diluted.

U-2113. Håkanbol 18, INS **8720 ± 190**
6770 B.C.
 $\delta C^{13} = -32.3\text{‰}$

Clay-gyttja from Håkanbol (59° 10' N Lat, 14° 22' E Long), Degerfors, Närke, Sweden. Sample 1840 cm below surface. Zone V (Jessen). Assumed older than Svea R. *Comment:* diluted.

U-2091. Håkanbol 5, INS **9240 ± 550**
7290 B.C.
 $\delta C^{13} = -33.5\text{‰}$

Clay-gyttja with carbon fragments from Håkanbol (59° 10' N Lat, 14° 22' E Long), Degerfors, Närke, Sweden, at depth 1830 cm in same kettle hole filled by peat as U-2080. Zone V (Jessen). At short-lived birch maximum. Assumed older than Svea R. *Comment:* sample contained more carbon fragments than U-2080. Diluted.

U-2092. Håkanbol 5, H **9450 ± 320**
7500 B.C.
 $\delta C^{13} = -31.8\text{‰}$

Humus products extracted from sample used for U-2091. *Comment:* diluted.

U-2101. Håkanbol 21, INS **9450 ± 210**
7500 B.C.

$$\delta C^{13} = -33.2\text{‰}$$

Clay-gyttja from Håkanbol (59° 10' N Lat, 14° 22' E Long), Degerfors, Närke, Sweden, at depth 1825 cm below surface. Zone V (Jessen). Below start of *Alnus*. Assumed to be beginning of Svea R. *Comment*: diluted.

U-2094. Solbergatjärn 110 A, INS **8600 ± 600**
6650 B.C.

$$\delta C^{13} = -31.7\text{‰}$$

Fine-detritus-gyttja from Solbergatjärn (59° 13' N Lat, 14° 27' E Long), Degerfors, Närke, Sweden, at depth 733 cm below surface. Zone boundary V/VI (Jessen). At beginning of *Alnus*, which supposedly occurred 8500 B.P. according to investigations by Fromm (1938) and Lundqvist (1957). *Comment*: diluted.

U-2102. Solbergatjärn 110 A, H **9270 ± 550**
7320 B.C.

$$\delta C^{13} = -33.0\text{‰}$$

Humus products extracted from sample used for U-2094. *Comment*: diluted.

U-2134. Klippetorpstjärn 125 A, INS **9330 ± 600**
7380 B.C.

$$\delta C^{13} = -30.8\text{‰}$$

Fine-detritus-gyttja from Klippetorpstjärn (59° 12' N Lat, 14° 26' E Long), Degerfors, Närke, Sweden, at depth 638 cm below surface. Zone boundary V/VI (Jessen). At beginning of *Alnus* (cf. U-2094). *Comment*: diluted.

U-2093. Håkanbol 25, INS **8920 ± 160**
6970 B.C.

$$\delta C^{13} = -34.8\text{‰}$$

Fine-detritus-gyttja from Håkanbol (59° 10' N Lat, 14° 22' E Long), Degerfors, Närke, Sweden, at depth 1800 cm below surface. Zone boundary V/VI (Jessen). At beginning of *Alnus* (cf. U-2094 and U-2090, 10 km away). *Comment*: diluted.

U-2108. Håkanbol 25, H **8690 ± 530**
6740 B.C.

$$\delta C^{13} = -33.3\text{‰}$$

Humus products extracted from sample used for U-2093. *Comment*: diluted.

Land uplift series, Central Sweden

Sediments from Central Sweden, coll. from ancient lakes developed by isolation from sea (S. Florin, 1944, 1948, 1963). Subm. by S. Florin, Inst. Quaternary Geol., Univ. Uppsala, Sweden. Pollen analyses by Th. Candolin. Several samples were dated previously (Radiocarbon: 1959, v. 1, p. 94; 1964, v. 6, p. 301-302; 1965, v. 7, p. 322-324; 1967, v. 9, p. 463-464).

U-648. Nedre Mogetorp, 670813, INS **6640 ± 80**
4690 B.C.
 δC^{13} assumed -34.1%

Gyttja with *Carex* from Nedre Mogetorp (59° 00' N Lat, 16° 09' E Long), Södermanland, Sweden, alt 44.8 m, at depth 396 to 400 cm below surface, Littorina time, before LII maximum. Layer corresponds to 400 cm depth below surface in diagram given by Florin (1948). Coll. 1967 by S. Florin and Ingrid U. Olsson.

U-2055. Nedre Mogetorp, 670813, H **5990 ± 140**
4040 B.C.
 $\delta C^{13} = -27.3\%$

Humus products extracted from sample used for U-648 and U-2115.
Comment: diluted.

U-2115. Nedre Mogetorp, 670813, KMnO₄ **>17,600**
 δC^{13} assumed -25.0%

Gas obtained by wet combustion of sample with KMnO₄. *Comment:* high age indicates that sediment contains old carbon, probably graphite, and that age of U-648 in radiocarbon yr is too high. Diluted.

U-700. Överåda, INS **2750 ± 70**
800 B.C.
 $\delta C^{13} = -26.5\%$

Brackish-water gyttja from Överåda (58° 53' N Lat, 17° 30' E Long), Vagnhärad-Trosa, Sweden, alt 27.0 m. Sample, 10 cm long, 150 cm below surface. Sub-Boreal, Zone VIII (Jessen), assoc. with neolithic pitted ware.

U-702. Överåda, H **2080 ± 60**
130 B.C.
 $\delta C^{13} = -26.2\%$

Humus products from sample used for U-700.

Sjödyn series

Gyttja from tarn Sjödyn (59° 55' N Lat, 17° 42' E Long), Vaksala parish, Uppland, Sweden, alt 34.7 m. Drainage threshold at 36.8 m. Sediments described by Sidenvall (1967). Depth to post-glacial clay ca. 6.5 m below water surface, water depth ca. 1.26 m. Livingstone borer used as sampler. Pollen and diatom analyses by Sidenvall. Coll. 1963 and subm. by Jan Sidenvall, Inst. Quaternary Geol. Univ. Uppsala, Sweden.
Comment: samples diluted.

U-2047. Sjödyn, 441 to 434, INS **4580 ± 130**
2630 B.C.
 $\delta C^{13} = -22.8\%$

Clay-gyttja, gray-greenish with olive-greenish bands, 441 to 434 cm below surface at beginning of lagoon stage. Zone VIII (Jessen).

U-2046. Sjödyn 430 to 425, INS **4850 ± 140**
2900 B.C.
 $\delta C^{13} = -26.5\%$

Gyttja, 430 to 425 cm below surface from boundary between lagoon

gyttja and freshwater gyttja, coll. 10 cm higher than U-2047. Zone VIII (Jessen).

U-2074. Sjödyn 230, INS **2600 ± 270**
650 B.C.

δC^{13} assumed -24.6‰

Gyttja with some dy, ca. 230 cm below surface. Zone boundary VII/VIII (Jessen). *Alnus* decreasing and *Betula* increasing. *Carpinus betulus* and *Fagus silvatica* show continuous curves. Sample shows beginning of *Picea* and of agriculture.

U-2075. Sjödyn 230, H **2720 ± 140**
770 B.C.

$\delta C^{13} = -31.8\text{‰}$

Humus products extracted from sample used for U-2074.

Floran series

Gyttja from mire district "Floran" in N Uppland, Sweden, to determine land uplift and to study vegetational history. District described by Ingmar (1963). Previous samples in series are given in Radiocarbon, 1965, v. 7, p. 324-326. Pollen analyses by Thorolf Candolin, diatom analyses by Tord Ingmar. Coll. and subm. by Ingmar, Inst. Plant Ecol. Univ. Uppsala, Sweden.

U-2089. Sörbackenmossen 3, INS **4650 ± 130**
2700 B.C.

$\delta C^{13} = -21.4\text{‰}$

Gyttja from bog Sörbackenmossen (60° 20' N Lat, 17° 42' E Long), Uppland, Sweden, alt 42 m. Sample, 3-cm-thick layer of brackish-water gyttja, immediately below boundary to freshwater gyttja. Sample grayish brown, somewhat minerogenous. Zone VIII (Jessen). Coll. 1966.

U-2073. Sörbackenmossen 3, H **4590 ± 80**
2640 B.C.

$\delta C^{13} = -20.1\text{‰}$

Humus products extracted from sample used for U-2089.

U-663. Sörbackenmossen 2, INS **4660 ± 90**
2710 B.C.

$\delta C^{13} = -22.4\text{‰}$

Gyttja from bog Sörbackenmossen (60° 20' N Lat, 17° 42' E Long), Uppland, Sweden, alt 42 m. Sample, 3-cm-thick layer of freshwater gyttja, immediately above boundary to brackish-water gyttja. Sample grayish brown, hardly minerogenous. Zone VIII (Jessen). Coll. 1966.

U-2109. Skrivarmyren 2, INS **4470 ± 110**
2520 B.C.

$\delta C^{13} = -16.5\text{‰}$

Gyttja from bog Skrivarmyren (60° 20' N Lat, 17° 44' E Long), Uppland, Sweden, drainage threshold at alt 39 m. Sample, 2-cm-thick layer of brackish-water gyttja, just below boundary to freshwater gyttja. Sample dark olive green, somewhat minerogenous and gelatinous. Zone VIII (Jessen). Coll. 1967.

U-664. Skrivarmyren 1, INS **3910 ± 100**
1960 B.C.

$$\delta C^{13} = -21.4\text{‰}$$

Gyttja from bog Skrivarmyren (60° 20' N Lat, 17° 44' E Long), Uppland, Sweden, drainage threshold at alt 39 m. Sample, 2-cm-thick layer of freshwater gyttja, immediately above boundary to brackish-water gyttja. Sample reddish-brown, gelatinous Cyanophyceae-gyttja. Zone VIII (Jessen). Coll. 1967.

U-639. Skrivarmyren 1, H **4350 ± 80**
2400 B.C.

$$\delta C^{13} \text{ assumed } -21.4\text{‰}$$

Humus products extracted from sample used for U-664.

U-2025. Ulvsbo trusk 4, INS **3690 ± 100**
1740 B.C.

$$\delta C^{13} = -24.0\text{‰}$$

Gyttja from tarn Ulvsbo trusk (60° 19' N Lat, 17° 46' E Long), Uppland, Sweden, drainage threshold at alt 35 m. Sample, 4-cm-thick layer of light-brown and slightly minerogenous gyttja around transition from brackish-water gyttja to freshwater gyttja. Zone VIII (Jessen). Coll. 1965.

U-704. Ulvsbo trusk 3, H **3020 ± 220**
1070 B.C.

$$\delta C^{13} = -19.7\text{‰}$$

Humus products extracted from gyttja from tarn Ulvsbo trusk (60° 19' N Lat, 17° 46' E Long), Uppland, Sweden, drainage threshold at alt 35 m. Sample, 2-cm-thick layer of brackish-water gyttja, immediately below boundary to freshwater gyttja. Sample light-brown and slightly minerogenous. Zone VIII (Jessen). Coll. 1964.

U-2024. Ensjön 2, INS **2950 ± 170**
1000 B.C.

$$\delta C^{13} = -21.1\text{‰}$$

Gyttja from lake Ensjön (60° 24' N Lat, 17° 51' E Long), Uppland, Sweden, drainage threshold at alt 27 m. Sample, 3-cm-thick layer of brackish-water gyttja, immediately below boundary to freshwater gyttja. Sample dark olive green and somewhat minerogenous and gelatinous. Zone VIII (Jessen). Coll. 1964 .

U-642. Ensjön 2, H **3660 ± 80**
1710 B.C.

$$\delta C^{13} = -19.6\text{‰}$$

Humus products extracted from sample used for U-2024.

U-537. Västerängen 2, INS 1 **2350 ± 220**
400 B.C.

$$\delta C^{13} = -28.2\text{‰}$$

Gyttja from ditched peatland Västerängen (60° 25' N Lat, 14° 44' E Long), Uppland, Sweden, drainage threshold at alt 20 m. Sample, 1-cm-thick layer, uppermost part of brackish-water gyttja rich in *Phrag-*

mites, underlying gyttja-free *Phragmites* peat. Sample represents isolation from sea in this very shallow basin. Sample underlay marked increase of *Picea* pollen in beginning of Zone IX (Jessen). Coll. 1963.

U-574. Västerängen 2, INS 2 **2610 ± 70**
660 B.C.
 $\delta C^{13} = -26.9\%$

Gyttja from same sample as U-537, but new pretreatment.

U-2078. Kätjåure **1500 ± 180**
A.D. 450
 $\delta C^{13} = -23.4\%$

Peat from Kätjåure (67° 18' N Lat, 17° 00' E Long), Padjelanta, Lappland, Sweden. Sample, 80 cm below surface, from frozen peat. Coll. 1967 and subm. by Anders Rapp, Dept. Physical Geog. Univ. Uppsala, Sweden. *Comment*: diluted.

Latnjajaure series

Diatom gyttja from Latnjajaure (68° 21' N Lat, 18° 37' E Long), Abisko, Lappland, Sweden. Core might contain material washed down from surroundings. Coll. 1967; subm. by Arnold Nauwerck, Inst. Limnology, Univ. Uppsala, Sweden. *Comment*: inconsistency in results might be due to contamination during storage. All samples diluted. Very little humus was recovered.

U-2086. Latnjajaure, 138.5 to 125, INS **6510 ± 230**
4560 B.C.
 $\delta C^{13} = -25.3\%$

Gyttja from 138.5 to 125 cm depth in core.

U-2087. Latnjajaure, 138.5 to 125, H **100 ± 300**
A.D. 1850
 δC^{13} assumed -25.2%

Humus products extracted from sample used for U-2086.

U-2084. Latnjajaure, 65 to 55, INS **10,750 ± 340**
8800 B.C.
 $\delta C^{13} = -28.2\%$

Gyttja from 65 to 55 cm depth in core.

U-2085. Latnjajaure, 65 to 55, H **+1100**
7500
-900
5600 B.C.
 δC^{13} assumed -25.2%

Humus products extracted from sample used for U-2084.

II. ARCHAEOLOGIC SAMPLES

A. Central Europe

U-611. Budva ship **150 ± 50**
A.D. 1800
 $\delta C^{13} = -25.4\%$

Wood from wrecked ship at Budva (42° 17' N Lat, 18° 50' E Long), Montenegro, Yugoslavia. Coll. 1966 by V. Stanišić and subm. by A.

Sliepčević, Institute Ruder Bošković, Zagreb, Yugoslavia. *Comment*: another portion of same sample was dated as a quick check and also gave very low age.

*B. Sweden***Gårdlösa series**

Charcoal from Gårdlösa grave-field No. 2 and No. 3 (55° 34' N Lat, 14° 08' E Long), Smedstorp parish, Skåne, Sweden. Coll. and subm. by Berta Stjernquist, Inst. Scandinavian Antiquities, Univ. Lund, Sweden. Several samples were dated previously (*Radiocarbon*: 1965, v. 7, p. 326-327; 1967, v. 9, p. 465-467).

U-697. Gårdlösa 2, Stone-settings 1967, No. 67:10 **2400 ± 80**
450 B.C.
 $\delta C^{13} = -23.8\text{‰}$

Charcoal found in bottom of hearth, 0.15 to 9.20 m below big stone in setting. Coll. 1967.

U-672. Gårdlösa 2, Stone-settings 1967, **1340 ± 70**
No. 67:32 **A.D. 610**
 $\delta C^{13} = -26.5\text{‰}$

Charcoal from bottom at 0.58 to 0.63 m depth with bones of a dog. Coll. 1967.

U-670. Gårdlösa 3, House XLIX **1170 ± 60**
A.D. 780
 $\delta C^{13} = -22.8\text{‰}$

Charcoal, at depth 0.75 to 0.85 m, below surface with potsherds, metal objects, etc. Coll. 1966. *Comment*: upper 0.2 m of profile cultivated.

U-671. Gårdlösa 3, Grave 36 **2570 ± 80**
620 B.C.
 $\delta C^{13} = -25.4\text{‰}$

Charcoal from 0.4 to 0.8 m depth, from post-hole below cairn, where Grave 36 was located. Coll. 1966.

U-674. Gårdlösa 3, Grave 42 **1300 ± 80**
A.D. 650
 $\delta C^{13} = -26.8\text{‰}$

Charcoal from 0.35 to 0.55 m depth, from layer with charcoal and soot; found with metal objects. Coll. 1965. *Comment*: upper 0.15 m cultivated.

U-673. Sandby 11, Well **1390 ± 70**
A.D. 560
 $\delta C^{13} = -24.0\text{‰}$

Wood from Sandby (54° 26' N Lat, 14° 12' E Long), Borrby parish, Skåne, Sweden. Sample from oak-log, 0.73 m deep covered by 0.30 m stones. A hearth was on log. Coll. 1965. *Comment*: upper 0.2 m cultivated.

Stora Alvaret series

Charcoal from Mysinge 1:9, 1:3 (56° 32' N Lat, 16° 30½' E Long), Öland, Sweden. Iron fibula indicate usage of settlement to ca. A.D. 300.

Charcoal pieces sorted to allow determination of different bush and tree species. Coll. 1966 and subm. by Ebba-Stina Königsson, Inst. Nordic Antiquities, Univ. Uppsala, Sweden.

U-654. Mysinge, F 114:1 **1430 ± 70**
A.D. 520
 $\delta C^{13} = -25.0\%$

Charcoal, of undetermined specie, 0.5 m below surface, overlying bedrock close to wall of one "house," A:4.

U-2063. Mysinge, F 114:2 **1630 ± 100**
A.D. 320
 $\delta C^{13} = -19.0\%$

Rhamnus cathartica from same place and level as U-654.

U-653. Mysinge, F 124:1 **1710 ± 70**
A.D. 240
 $\delta C^{13} = -24.0\%$

Charcoal, of undetermined specie, 0.5 m below surface of "house" A:2, close to sample U-654.

U-649. Mysinge, F 124:2 **1800 ± 80**
A.D. 150
 $\delta C^{13} = -25.6\%$

Betula and *Populus* from same place and level as U-653.

U-662. Mysinge F 124:3 **1640 ± 70**
A.D. 310
 $\delta C^{13} = -23.4\%$

Juniperus and *Taxus* from same place and level as U-653.

Västerås-Äs series

Charcoal from Äs (59° 44' N Lat, 16° 30' E Long), Romfartuna parish, Västmanland, Sweden, undisturbed cultural layer containing pitted pottery. Coll. 1967 and subm. by Lars Löfstrand, Inst. Nordic Antiquities, Univ. Uppsala, Sweden.

U-2112. Västerås-Äs 785/396 **1120 ± 70**
830 B.C.
 $\delta C^{13} = -25.5\%$

Charcoal from lowest part of Layer G-S II, at ca. 15 to 20 cm depth, dating Stå MN pitted pottery culture.

U-701. Västerås-Äs 841/394 **2250 ± 60**
300 B.C.
 $\delta C^{13} = -25.7\%$

Charcoal from hearth in middle of Layer G-S II, at ca. 15 cm depth, dating ornamented pitted pottery of Säter III type.

Berthåga series

Charcoal from dwelling under Iron age grave-field. Site excavated in connection with new grave-field, Berthåga kyrkogård (59° 51' N Lat,

17° 35' E Long), Uppland, Sweden. Coll. and subm. by Astrid Sjöberg, Inst. Nordic Antiquities, Univ. Uppsala, Sweden.

U-656. Berthåga A212, F2 **2770 ± 80**
820 B.C.
 $\delta C^{13} = -23.9\text{‰}$

Charcoal from burned piece of wood, ca. 40 × 10 cm, from Hearth Pit A212, also containing animal tooth and burned clay. Coll. 1967.

U-657. Berthåga A218, F2 **2390 ± 80**
440 B.C.
 $\delta C^{13} = -24.6\text{‰}$

Charcoal from concentration in layer with charcoal and soot at bottom of split-stone mound containing potsherd and burned clay, at ca. 0.5 m depth. Coll. 1967.

U-658. Berthåga A216, F71 **100 ± 70**
A.D. 1850
 $\delta C^{13} = -24.8\text{‰}$

Charcoal from layer with charcoal and soot, at 0.30 to 0.35 m depth. Nearby were house-remains, post-holes, burned clay, animal bones, and potsherds. Coll. 1966.

U-684. Berthåga A216, R214296 **80 ± 110**
A.D. 1870
 δC^{13} assumed -24.9‰

Charcoal from same layer as U-658. Coll. 1966.

U-688. Avasjön **460 ± 50**
A.D. 1490
 $\delta C^{13} = -24.3\text{‰}$

Wood from SE end of Avasjön (64° 19' N Lat, 21° 20' E Long), Löfvånger parish, Västerbotten, Sweden, from board of ship, ca. 10 m long. Part of ship is still in lake. Avasjön supposedly is part of old passage; 4 wrecks are known. Commercial center at Mångbyn was flourishing ca. 450 yr ago. Coll. 1926: subm. by Peter Gustavsson, Inst. Nordic Antiquities, Univ. Uppsala, Sweden.

Yttervik series

Charcoal from Yttervik (64° 40' N Lat, 21° 10' E Long), Bureå parish, Västerbotten, Sweden, from burial cairn. Described by Westin (1962). Coll. 1875 by Hans Hildebrand, Central Office Natl. Antiquities, Stockholm, Sweden; subm. by Hans Christiansson, Inst. Nordic Antiquities, Univ. Uppsala, Sweden. Samples stored, since coll. in Mus. Antiquities, Stockholm, Sweden.

U-2057. Yttervik 5 **640 ± 90**
A.D. 1310
 $\delta C^{13} = -19.0\text{‰}$

Charcoal from Grave 5, a stone-setting. No other artifacts found. Grave 10 m W of old road to Båtvik. Collection No. SHM 7043 B.

U-655. Yttervik 9 **890 ± 70**
A.D. 1060
 $\delta C^{13} = -26.6\%$

Charcoal from Grave 9, a cairn. No other artifacts found. Grave ca. 5 m E of old road to Båtvik. Collection No. SHM 7043 B.

U-2070. Yttervik 12 **1110 ± 100**
A.D. 840
 $\delta C^{13} = -35.7\%$

Charcoal from Grave 12, a cairn. No other artifacts found. Grave ca. 10 m E of old road to Båtvik. Collection No. SHM 7043 B. *Comment:* diluted.

Bjurselet series

Charcoal from Bjurselet (65° 00' N Lat, 21° 04' E Long), Byske parish, Västerbotten, Sweden. From Stone age dwelling-site, alt 53 m, with imported flint axes, scrapers, and potsherds. Subm. by Hans Christiansson. Upper Layer consisted of cultivated soil, Layer I, underlain by sand, Layers II, III, and IV. Preliminary described by Christiansson (1965a, b). Pollen profile from same area and several archaeologic samples were dated previously (Radiocarbon, 1967, v. 9, p. 467-469).

U-647. Bjurselet 210/228 Åkern **2140 ± 80**
190 B.C.
 $\delta C^{13} = -24.7\%$

Charcoal, from field, from lowest layer with brittle burned stones. Coll. 1966 by P. Gustavsson, Inst. Nordic Antiquities, Univ. Uppsala, Sweden.

U-2035. Bjurselet, Hunting Pit 1, Svarvet **930 ± 190**
A.D. 1020
 $\delta C^{13} = -25.8\%$

Charcoal, at 0.9 m depth, from hunting pit, 1.2 m deep. Coll. 1965 by Christiansson. *Comment:* similar pits recorded in Radiocarbon, 1967, v. 9, p. 469. Diluted.

U-2045. Bjurselet 216/119 Svarvet **4310 ± 180**
2360 B.C.
 $\delta C^{13} = -25.1\%$

Wood and charred wood from Layer II in sandy soil, probably never cultivated. Coll. 1965 by H. C. Vorting. *Comment:* diluted.

III. TREE-RING SAMPLES

Northern Sweden series

Wood from N Sweden. Tree felled in 1966, had 75 tree rings. Sample from ca. 1 m above ground. Subm. 1966 by Ing. A. Assarsson, Mo and Domsjö AB, Örnsköldsvik, Sweden, where chemical treatment was made. Wood was pulverized, water was removed by acetone, and pulverized wood was treated by ethanol-benzene (1:2) mixture for 20 hrs and washed with water for 20 hrs. Extract from acetone and ethanol-benzene treat-

ment was evaporated at temp. $<40^{\circ}$ C, dissolved in di-ethyl-ether, and dried. Part of remaining material was treated with Na-chlorite and acetic acid at elevated temp. to give cellulose and holocellulose (Wise *et al.*, 1946). Results will be discussed by Assarsson and Olsson.

General Comment: extract has excess of C^{14} relative to air for rather young tree rings but is depleted for oldest rings. Sample U-598 indicates that this wood is, to some extent, affected by younger material. Difference between the 3 fractions of youngest sample is not significant since relative amount of material produced during 1960-1966 probably varied from year to year and "atomic bomb effect" changed greatly.

TABLE 1

No.	Sample	$\delta C^{13} \text{‰}$	Δ in ‰; $T_{1/2} = 5730$ yr		
			Extract	Extracted wood	Holo-cellulose
U-643	Tree-ring 69-75, xylem	-24.5			620 ± 9
U-644	formed 1960-1966.	-32.0		645 ± 10	
U-645		-27.9	554 ± 8		
U-626	Tree-ring 59-69, xylem	-21.9			-27 ± 7
U-627	formed 1950-1955.	-25.1		-31 ± 6	
U-628		-28.5	119 ± 7		
U-594	Tree-ring 44-49, xylem	-20.7			-28 ± 7
U-595	formed 1935-1941.	-25.0		-21 ± 5	
U-596		-28.2	117 ± 6		
U-597	Tree-ring 26-31, xylem	-23.7			-22 ± 6
U-598	formed 1917-1922, heartwood	-25.2		7 ± 5	
U-599	formed 1960-1965.	-26.8	10 ± 5		
U-629	Tree-ring 0-10, xylem	-23.2			8 ± 6
U-630	formed 1891-1901, heartwood.	-25.7		13 ± 5	
U-631		-26.2	-14 ± 6		

Argentina series

Tree trunk (*Fitzroya cupressoides*) from Parque Nac. Los Alerces (43° S Lat, 71° W Long), Esquel, Chubut, Argentina. Park created 1937. Tree probably felled a few yr earlier. Trunk donated by Intendencia, Parque Nac. Los Alerces through Åke Vinterbäck, Univ. Uppsala, Sweden.

Comment: tree-ring project is performed in collaboration with Hendrik de Waard, J. C. Vogel, and J. C. Lerman, Natuurkundig Lab. Groningen, The Netherlands. Tree rings counted by Horrying, High School of Forestry, Stockholm, Sweden. In the calculations it was assumed that tree was felled A.D. 1934. Samples will be checked for C^{13}/C^{12} ratio in Groningen. Cellulose and lignin prepared acc. to Olson and Broecker (1958) after normal pretreatment of wood.

Lab. no.	Age before		Substance	$\delta C^{13} \text{‰}$	$\delta C^{14} \text{‰}$	$\Delta \text{‰}$
	A.D. 1934	Age B.P. (1950)				
U-509	649-645	663	Lignin	-24.2	-10.5	-12 \pm 5
U-650	644-640	658	Lignin	-23.4	-20.1	-23 \pm 5
U-508	634-630	648	Cellulose	-21.3	-17.3	-25 \pm 5
U-668	629-625	643	Cellulose	-22.9	-16.5	-21 \pm 5
U-667	629-625	643	Xylem	-24.1	-13.1	-15 \pm 5

IV. GEOCHEMICAL SAMPLES

Lake sediment series

Living plants from hard-water lakes coll. to demonstrate influence of dissolved carbonate on C^{14}/C^{12} ratio in plants, components of sediments studied on Öland. Present C^{14}/C^{12} ratio of atmosphere will be given in Radiocarbon, 1970, v. 12. Subm. by Lars-König Königsson, Inst. Quaternary Geol., Univ. Uppsala, Sweden.

U-651. Carex 270666, D

$$\begin{aligned}\delta C^{14} &= +689 \pm 13 \text{‰} \\ \Delta &= +700 \pm 13 \text{‰} \\ \delta C^{13} &= -28.1 \text{‰}\end{aligned}$$

Carex elata from Dröstorps mose (56° 35' N Lat, 16° 32' E Long), Öland, Sweden. Coll. June 27, 1966 by Königsson and B. Fredskild. *Comment:* another portion of this sample was measured by J. M. Punning, TA-222: $\delta C^{14} = 616 \text{‰}$.

U-2083. Floating 270666, D

$$\begin{aligned}\delta C^{14} &= +533 \pm 12 \text{‰} \\ \Delta &= +556 \pm 12 \text{‰} \\ \delta C^{13} &= -32.4 \text{‰}\end{aligned}$$

Floating plants from Dröstorps mose (56° 35' N Lat, 16° 32' E Long), Öland, Sweden. Coll. June 27, 1966 by Königsson and Fredskild.

U-2072. Submerged 270666, D

$$\begin{aligned}\delta C^{14} &= +561 \pm 12 \text{‰} \\ \Delta &= +579 \pm 12 \text{‰} \\ \delta C^{13} &= -30.9 \text{‰}\end{aligned}$$

Submerged plants from Dröstorps mose (56° 35' N Lat, 16° 33' E Long), Öland, Sweden. Coll. June 27, 1966 by Königsson and Fredskild.

U-652. Characeae and submerged 270666, D

$$\begin{aligned}\delta C^{14} &= +468 \pm 9 \text{‰} \\ \Delta &= +457 \pm 9 \text{‰} \\ \delta C^{13} &= -21.4 \text{‰}\end{aligned}$$

Characeae and other submerged plants from Dröstorps mose (56°

35' N Lat, 16° 32' E Long), Öland, Sweden. Coll. June 27, 1966 by Königsson and Fredskild.

U-2068. Characeae 270666, BK

$$\begin{aligned}\delta C^{14} &= +396 \pm 12\% \\ \Delta &= +406 \pm 12\% \\ \delta C^{13} &= -28.7\%\end{aligned}$$

Characeae from Bårby Källa (Lilla Dammen) (56° 31' N Lat, 16° 30' E Long), Öland, Sweden. Coll. June 27, 1966 by Königsson and Fredskild.

U-2118. Algae 020768, MM

$$\begin{aligned}\delta C^{14} &= +327 \pm 10\% \\ \Delta &= +341 \pm 10\% \\ \delta C^{13} &= -30.2\%\end{aligned}$$

Algae from Möckelmosen (56° 32' N Lat, 16° 31' E Long), Öland, Sweden. Coll. July 2, 1968 by Königsson.

Resin sample series

Birch-bark from Nåsten (ca. 59° 49' N Lat, 17° 35' E Long), Uppland, Sweden. Tree felled 1967; bark coll. 1968 by Ingrid U. Olsson. Tree ca. 30 yr old. Resin, used for tightening of vessels and fixing of metal objects to wood, etc. is supposed to derive from birch-bark (Sandermann, 1965). Chemical analysis and infra-red spectra of different extracts of archaeological resin and of resin obtained by heating, at 400°C, of outer parts of birch-bark, agree so well that origins might be identical. In Swedish resin some part of inner bark might have been used. *Comment:* results indicate that resin dates would be systematically somewhat too high—the error dependent on age of tree. Only one pair of sample of resin-charcoal has been dated in Uppsala, U-151 and U-152: 2070 ± 90 and 2240 ± 90, respectively (Deevey *et al.*, 1967, p. 177). All archaeological resin samples (15), dated in Uppsala, have their δC^{13} between -26 and -34‰.

U-2116. Birch-bark, white, 09068, Ac.

$$\begin{aligned}\delta C^{14} &= +189 \pm 11\% \\ \Delta &= +191 \pm 11\% \\ \delta C^{13} &= -25.9\%\end{aligned}$$

Acetone extract from outer layer of birch-bark.

U-698. Birch-bark, white, 090668, INS

$$\begin{aligned}\delta C^{14} &= +319 \pm 10\% \\ \Delta &= +330 \pm 10\% \\ \delta C^{13} &= -29.2\%\end{aligned}$$

Remains of birch-bark after acetone extraction and normal pre-treatment.

REFERENCES

Date lists:

Copenhagen VII	Tauber, 1966
Uppsala I	Olsson, 1959
Uppsala II	Olsson, 1960
Uppsala III	Olsson <i>et al.</i> , 1961
Uppsala IV	Olsson and Kilicci, 1964
Uppsala V	Olsson and Pivanuj, 1965
Uppsala VII	Olsson <i>et al.</i> , 1967

Balchin, W. G. V., 1941, The raised features of Billefjord and Sassenfjord West Spitsbergen: *Geog. Jour.*, v. 97, p. 364-376.

- Bárdarson, G. B., 1910, Traces of changes of climate and level at Húnaflói, N-Iceland: Postglaziale Klimaveränderungen, Internat. Geol. Kongr., Stockholm, p. 347-352.
- 1923, Old sea-deposits in Borgarfjörður and Hvalfjörður: Soc. Sci. Isl., v. 1, 118 p.
- Birkenmajer, Krzysztof, 1960, Raised marine features of the Hornsund area, Vestspitsbergen: Studia Geol. Polon., v. 5, 95 p.
- Christiansson, Hans, 1965a, De arkeologiska undersökningarna vid Bjurselet i Byske: Västerbotten, p. 191-202.
- 1965b, Flint finds in Västerbotten; Hunting, trade or agriculture?, in: Hunting and Fishing, Nordic symposium in Luleå 1962, Norrbottens Mus., Luleå, Sweden, p. 91-110.
- Craig, Harmon, 1961, Mass-spectrometer analyses of radiocarbon standards: Radiocarbon, v. 3, p. 1-3.
- Dahl, Ragnar, 1967, Senglaciala ackumulationsformer och glaciationsförhållanden in Narvik-Skjomenområdet, Norge: Norsk geog. tidskr., v. 21, p. 157-241.
- 1968, Glacial accumulations, drainage and ice recession in the Narvik-Skjomen district, Norway: Norsk geog. tidsskr., v. 22, p. 101-165.
- Deevey, E. S., Flint, R. F., and Rouse, I. (eds.), 1967, Radiocarbon measurements: Comprehensive index, 1950-1965, Yale Univ., New Haven, 221 p.
- Einarsson, Thorleifur, 1961, Pollenanalytische Untersuchungen zur spät- und postglazialen Klimageschichte Islands: Sonderöff. Geol. Inst. Univ. Köln, v. 6, 52 p.
- Eriksson, K. G. and Olsson, I. U., 1963, Some problems in connection with C^{14} dating of tests of Foraminifera: Geol. Inst. Univ. Bull., Uppsala, v. 42, p. 1-13.
- 1967, Några data från en planerad undersökning av en skalbank vid Linddalsskogen, Väjern, Mellersta Bohuslän: Teknik och natur, Studier tillägnade Gunnar Beskow, Akademiförlaget, Göteborg, p. 149-166.
- Feyling-Hanssen, R. W., 1955, Stratigraphy of the marine late-Pleistocene of Billefjorden, Vestspitsbergen: Norsk Polarinst. Skr. 107, 186 p.
- Feyling-Hanssen, R. W. and Olsson, Ingrid, 1959-1960, Five radiocarbon datings of post glacial shorelines in central Spitsbergen: Norsk geog. tidsskr., v. 17, p. 122-131.
- Florin, Sten, 1944, Havsstrandens förskjutningar och bebyggelseutvecklingen i östra Mellansverige under senkvartär tid. I. Allmän översikt: G. F. F., v. 66, p. 551-634.
- 1948, Havsstrandens förskjutningar och bebyggelseutvecklingen i östra Mellansverige under senkvartär tid. II. De baltiska strandbildningarna och stenåldersboplatsen vid Dammstugan nära Katrineholm: G. F. F., v. 70, p. 17-202.
- 1963, Bodenschwankungen i Schweden während des Spätquartärs: Baltica, v. 1, p. 233-264.
- Fredén, Curt, 1967, A historical review of the Ancylus Lake and the Svea River: G. F. F., v. 89, p. 239-267.
- Fromm, Erik, 1938, Geochronologisch datierte Pollendiagramme und Diatoméenanalysen aus Angermanland: G. F. F., v. 60, p. 365-381.
- Hoppe, Gunnar, 1965, Submarine peat in the Shetland Islands: Geog. Annales, v. 47, ser. A, p. 195-203.
- Hospers, J., 1953, Reversals of the main geomagnetic field I: Akad. Sci. Amsterdam, Proc. ser. B, v. 56, p. 467-476.
- Hyvärinen, Hannu, 1968, Late-quaternaly sediment cores from lakes on Bjørnøya: Geog. Annales, v. 50, ser. A, p. 235-245.
- Ingmar, Tord, 1963, Från havsvik till mosse: Sveriges Natur, Årsbok, v. 54, p. 155-177.
- Libby, W. F., 1955, Radiocarbon dating, 2nd ed.: Chicago, Univ. of Chicago Press, 175 p.
- Lundqvist, Gösta, 1957, C^{14} -analyser i svensk kvartärgeologi 1955-57 (with English summary): Sveriges geol. undersökning, ser. C, no. 557, v. 51, no. 8, 25 p.
- Norin, Erik, 1966, Edsengol-oasen i Gobiöknen: G. F. F., v. 88, p. 340-350.
- Olson, E. A. and Broecker, W. S., 1958, Sample contamination and reliability of radiocarbon dates: N.Y. Acad. Sci. Trans. ser. II, v. 20, p. 593-604.
- Olsson, Ingrid, 1958, A C^{14} dating station using the CO_2 proportional counting method: Arkiv f. Fysik, v. 13, p. 37-60.
- 1959, Uppsala natural radiocarbon measurements I: Am. Jour. Sci. Radiocarbon Suppl., v. 1, p. 87-102.
- 1960, Uppsala natural radiocarbon measurements II: Am. Jour. Sci. Radiocarbon Suppl., v. 2, p. 112-128.
- Olsson, Ingrid, Cazeneuve, Horacio, Gustavsson, John, and Karlén, Ingvar, 1961, Uppsala natural radiocarbon measurements III: Radiocarbon, v. 3, p. 81-85.

- Olsson, I. U., 1965, Computer calculations of C¹⁴ determinations: Proc., 6th internatl. conf. radiocarbon and tritium dating, Pullman, Washington, June 7-11, 1965, p. 383-392.
- 1966, Computer calculations of C¹⁴ determinations: Uppsala Univ. Inst. Physics Rep., UUIP-477, 11 p.
- 1968, The radiocarbon analyses of lake sediment samples from Björnøya: Geog. Annales, v. 50, ser. A, p. 246-247.
- Olsson, I. U. and Blake, Weston, Jr., 1961-1962, Problems of radiocarbon dating of raised beaches, based on experience in Spitsbergen: Norsk geog. tidsskr., v. 18, p. 47-64.
- Olsson, I. U. and Eriksson, K. G., 1965, Remarks on C¹⁴-dating of shell material in sea sediments: Progr. in Oceanog., v. 3, p. 253-266.
- Olsson, I. U., Göksu, Yeter, and Stenberg, Allan, 1968, Further investigations of storing and treatment of foraminifera and mollusks for C¹⁴-dating: G. F. F., v. 90, p. 417-426.
- Olsson, I. U. and Kilicci, Serap, 1964, Uppsala natural radiocarbon measurements IV: Radiocarbon, v. 6, p. 291-307.
- Olsson, I. U. and Piyanuj, Piya, 1965, Uppsala natural radiocarbon measurements V: Radiocarbon, v. 7, p. 315-330.
- Olsson, I. U., Stenberg, Allan, and Göksu, Yeter, 1967, Uppsala natural radiocarbon measurements VII: Radiocarbon, v. 9, p. 454-470.
- Sandermann, Wilhelm, 1965, Untersuchung vorgeschichtlicher "Gräberharze" und Kitte: Techn. Beitr. Archäol., v. 2, p. 58-73.
- Sidenvall, Jan, 1967, Senkvartär vegetationsutveckling i trakten av Sjärdyn, Vaksala socken, Uppland: Inst. Quaternary Geol., Uppsala, 16 p.
- Sjöberg, Astrid, 1969, A Bronze Age site at Berthåga: Tor, 1967/1968, v. 12, p. 243-247.
- Tauber, Henrik, 1966, Copenhagen radiocarbon dates VIII: Radiocarbon, v. 8, p. 213-234.
- Thorarinsson, Sigurdur, 1955, The *Nucella* shore line at Húnaflói in the light of tephrochronological and radiocarbon datings: Náttúrufræðingurinn, v. 25, p. 172-186.
- Todd, Ruth, 1958, Foraminifera from Western Mediterranean deep-sea cores: Rept. Swedish Deep-Sea Exped., v. 8, no. 3, p. 167-217.
- Westin, Gunnar, 1962, Övre Norrlands forntid. Övre Norrlands historia I. (Umeå), p. 1-122.
- Wise, L. E., Murphy, Maxine, and D'Addieco, A. A., 1946, Chlorite holocellulose, its fractionation and bearing on summative wood analysis and on studies on the hemicelluloses: Paper Trade Jour., v. 122, p. 35-43.