

- 3 Woolhandler S, Himmelstein DU. Competition in a publicly funded healthcare system. *BMJ* 2007; **335**: 1126–9.
- 4 Fulop N, Protosaltis G, Hutchings A, King A, Allen P, Normand C, et al. Process and impact of mergers of NHS trusts: multicentre case study and management cost analysis. *BMJ* 2002; **325**: 246.

David Dodwell, Consultant in Assertive Outreach, Cambridgeshire & Peterborough NHS Foundation Trust, Peterborough, email: david.dodwell@cpft.nhs.uk

doi: 10.1192/pb.36.12.476

Newer is not automatically better

It is ungenerous and unjustified for Helen Killaspy¹ to accuse George Lodge² of nostalgia and wearing rose-tinted spectacles just because she disagrees with him. Newer is not automatically better. We have had altogether too much frenetic reorganisation of mental health services where every change, no matter how hare-brained, is hailed 'an innovation'. Nobody waits to see whether it makes any difference, never mind delivers an improvement. It is whether an idea is right or not that matters, not how long it has been around. Similarly, it is disingenuous of her to claim that the service changes she describes were 'informed by research'.

New developments arise from a mixture of creative thinking and professional ambition, and there is nothing wrong with that. By the very nature of the beast, evidence comes later. We need the new services in place to research them rigorously or make judgements from mature experience. The National Service Framework is a case in point. Only one of the new teams imposed had any evidence for it at the time, and assertive outreach teams' international evidence was unravelling in the UK context as they were being rolled out.³

There was not a single randomised controlled trial of crisis teams until Johnson's excellent, but still unrepeated, 2005 study.⁴ The only two randomised controlled trials of early intervention teams also came later, and neither found a significant advantage in their declared primary outcomes. A more measured position is probably justified.

Continuity of care can be a complex concept to define⁵ but it is not that difficult to recognise. We can all grasp the importance of being treated by familiar individuals who know our situation and illness, of not being passed on, and not having to repeat our history to an endless stream of new staff whom we then have to learn to trust. Everybody who is asked, patients, staff or families, insists that they value continuity. I know I do.

Whatever else mental illnesses are, they are experienced, expressed and treated in relationships. George Lodge is right that these relationships have been given altogether too low a priority in recent planning and strategy. Our decade of fragmentation may have contributed some improved understanding of process, but undoubtedly at a cost of simple humanity and attention to the unique individuals for whom the whole edifice exists. Helen Killaspy is right that we have a progressive discipline, responsive to an expanding evidence base. That does not mean that every change is improvement, nor that more specialised services (with their inevitable fragmentation of care) are necessarily better for patients.

- 1 Killaspy H. Importance of specialisation in psychiatric services. Commentary on . . . How did we let it come to this? *Psychiatrist* 2012; **36**: 364–5.
- 2 Lodge G. How did we let it come to this? A plea for the principle of continuity of care. *Psychiatrist* 2012; **36**: 361–3.

- 3 Burns T, Creed F, Fahy T, Thompson S, Tyrer P, White I. Intensive versus standard case management for severe psychotic illness: a randomised trial (UK 700 Group). *Lancet* 1999; **353**: 2185–9.
- 4 Johnson S, Nolan F, Pilling S, Sandor A, Hoult J, McKenzie N, et al. Randomised controlled trial of acute mental healthcare by a crisis resolution team: the north Islington crisis study. *BMJ* 2005; **331**: 599.
- 5 Burns T, Catty J, White S, Clement S, Ellis G, Jones IR, et al. Continuity of care in mental health: understanding and measuring a complex phenomenon. *Psychol Med* 2009; **39**: 313–23.

Tom Burns, Professor of Social Psychiatry, Oxford University, UK, email: tom.burns@psych.ox.ac.uk

doi: 10.1192/pb.36.12.477

Author's response: Dr MacMillan (p. 475, this issue) is quite right to point out the importance of appropriate access to in-patient beds as a critical component of mental health services. That mental health systems should provide a balance of in-patient beds and community services tailored to the mental health needs and resources of the local community being served is something all mental health practitioners across the world can probably agree on. My commentary did not suggest that increased specialisation means we should do away with in-patient services, it simply stated the fact that investment in specialist community mental health teams (particularly crisis teams) through the National Service Framework for Mental Health was associated with a reduced need for in-patient admissions. Where I believe Dr Lodge and I also agree is on the need for continued investment in mental health rehabilitation services to prevent the inappropriate use of out-of-area placements for the small number of people with particularly complex and long-term psychoses.^{1,2}

Professor Burns' response (see letter above) states: 'It is ungenerous and unjustified for Helen Killaspy to accuse George Lodge of nostalgia and wearing rose-tinted spectacles just because she disagrees with him. Newer is not automatically better.' This accusation is not only unjust and ungenerous to those who have been working without feeling conflicted in both specialist and generalist services for many years, but it is without basis in fact. My commentary made clear, evidence-based justification for my view. I included reference to the lack of evidence for the effectiveness of assertive community treatment in the UK context that probably influenced subsequent disinvestment in this model. However, our research group, while contributing to such findings, simultaneously participated in a multicentred international study which suggested that assertive community treatment in the UK may have not performed as effectively as in Australia owing to lack of implementation of critical components that Professor Burns' own team identified through meta-analyses.^{3,4} His further accusation that I was 'disingenuous' is a little ironic given his lack of reference to the robust international evidence on which investment in the new specialist teams was made, not to mention the expanding evidence base for early intervention services.

Dr Dodwell's response (pp. 476–7, this issue) accuses me of dismissing evidence on therapeutic alliance, yet I did not mention it. It is a truism to say that the therapeutic alliance is important. Who would argue against the importance of being treated with humanity and respect in the therapeutic encounter? However, therapeutic alliance is not the same as continuity of care, which was, after all, the focus of Dr Lodge's piece.