

important in maintaining healthy habits and lifestyle in line with what is taught in a nursing degree.

*Disclosure of interest* The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.1253>

#### EV0924

### Integrated identification of new substantial gadget addiction: With selfie-mania phenomenon model

I. Sosin<sup>1,\*</sup>, Y. Chuev<sup>2</sup>, O. Goncharova<sup>2</sup>

<sup>1</sup> Kharkov medical academy of postgraduate education, Narcology, Kharkov, Ukraine

<sup>2</sup> Kharkiv Medical Academy of Postgraduate Education, Narcology, Kharkiv, Ukraine

\* Corresponding author.

*Introduction* Modern selfie-infatuation extent went far beyond fashion and subculture causing distinct tendency to non-chemical addiction state transformation requiring evidential scientific identification.

*Aims and objectives* To develop clinical-psychological and classification basis for new type of gadget addiction (selfie-mania), prevention and approaches to mental correction and rehabilitation.

*Method* Selfie attitude testing of 157 respondents-students, and internet resources topical data analysis were done.

*Results* Behaviour models were rubricated, design and content, causes, clinical-psychological description, selfie-phenomenon prevalence were clarified. It allows to state selfie-addiction specific key symptoms, conforming to ICD-10 diagnostic criteria for addiction states:

- psychological and emotional supercomfort feeling during realizing selfie-interventions (substantial analogue of euphoria phenomenon caused by psychoactive substances in narcology);
- desire for permanent updating and layouting in social networks selfie-portraits, selfie-positions (impulse control disorder analogue);
- selfie-modifying need causing adrenaline extreme and life threat (analogues: compulsive craving, megadoses, overdoses, amnesias);
- obsessive craving to increase daily number of selfie-shots (tolerance syndrome analogue);
- constant foreshortening change of selfie-interventions (analogue: experimentation with different psychoactive substances, searching behaviour);
- selfie-destruction psychopathological consequences (neurotism, mental and behavioural disorders, group pattern of behavioural selfie-deviations);
- formation of associated comorbid chemical and other substantial addiction forms;
- selfie-deprivation syndrome (analogue: alcohol or drug withdrawal syndrome);
- interdisciplinary range of problems (professional sphere of study for sociologists, psychologists, neurologists, psychiatrists, narcologists, psychotherapists, sexopathologists, lawyers).

*Conclusions* Selfie-addiction problem recognition is necessary on the level of inclusion to ICD, that will allow to develop scientific, legal and clinical base for integrated prevention, rehabilitation and treatment.

*Disclosure of interest* The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.1254>

#### EV0925

### Selfie as a method of perception of the virtual environment

M. Sulaimanova<sup>1</sup>, R. Sulaimanov<sup>2,\*</sup>

<sup>1</sup> Kyrgyz-Slavic University, Medical Psychology- Psychiatry- and Psychotherapy, Bishkek, Kyrgyz Republic

<sup>2</sup> Osnabrück University, Institute of Cognitive Science, Osnabrück, Germany

\* Corresponding author.

Nowadays selfie is one of the significant communication methods in the conditions of the developing internet space.

*Goal* The study of psychological mechanisms motivating perception of internet space via selfie.

*Objectives* The identification of:

- The personal-social necessity in selfie, as a method of the virtual world perception;
- The factors of personal-social necessity of the virtual communication via selfie;
- The personal features, which contribute to self-presentation in virtual space via selfie.

*Research methods* One hundred and thirty-four respondents aging between 16–28 (1<sup>st</sup> group: 68 respondents fond of selfie (50.8%), 2<sup>nd</sup> group: 66 respondents not fond of selfie (49.2%)) were studied using standardized scales of anxiety (Spilberger-Hanin), depression (Zung), ego-defense mechanisms (EGMs) (Plutchik-Kellerman-Konte), and life quality (WHO, 1991).

*Results* Desire of high evaluation dominates in 47.8% of respondents ( $r = 0.3$ ;  $P < 0.01$ ). Fifty point seven percent of the 1<sup>st</sup> group and 13.4% in 2<sup>nd</sup> group have heightened anxiety ( $r = 0.3$ ;  $P < 0.01$ ). Forty-three point two percent in 1<sup>st</sup> group and 23.9% in 2<sup>nd</sup> group don't have depression.

Subdepressive level dominates in 19.4% of 2<sup>nd</sup> group ( $r = 0.36$ ;  $P < 0.01$ ), against 11.9% of 1<sup>st</sup> group.

In 1<sup>st</sup> group dominate: denial (17.9%), projection (16.4%), regression (7.5%); in 2<sup>nd</sup> group: denial (17.9%), intellectualization (13.4%), projection (7.5%) ( $P < 0.05$ ). Forty-four point eight percent in 1<sup>st</sup> group and 40.3% in 2<sup>nd</sup> group have a high level of EGM stress ( $P < 0.05$ ). Forty-six point eight percent in 1<sup>st</sup> group have high level of subjective life quality, 0% had low one; in 2<sup>nd</sup> group: 26.9% have high level, 13.4% low one.

*Conclusions* Personal-social necessity of virtual space perception via selfie reflects the severity of narcissistic tendencies. Selfie in conditions of illusory reality increases the subjective level of life quality. Isolation of narcissistic personality from communication through selfie suggests subjective decrease of life quality. Effects of the selfie should be considered in the development of rational methods of use of internet-technologies and in the psycho-correction of people with low life quality.

*Disclosure of interest* The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.1255>

#### EV0926

### Cross-national trainee collaboration-The EFPT research experience in psychiatry

S. Tomori<sup>1,\*</sup>, T. Mogren<sup>2</sup>, M. Asztalos<sup>3</sup>, G. Sampogna<sup>4</sup>, M. Borovcanin<sup>5</sup>, G. Erzin<sup>6</sup>, J. Veiera<sup>7</sup>, R. Tipa<sup>8</sup>, A. Gurcan<sup>9</sup>, S. Naughton<sup>10</sup>, E.F.O.P.T. EFPT<sup>11</sup>

<sup>1</sup> University Hospital Center "Mother Teresa", Pediatric Service, Tirana, Albania

<sup>2</sup> Allmänpsykiatriska kliniken Falun/Säter, Allmänpsykiatriska kliniken Falun/Säter, Säter, Sweden

<sup>3</sup> Aalborg University Hospital, Psychiatry, Aalborg, Denmark

<sup>4</sup> University of Naples SUN, Department of Psychiatry, Naples, Italy